

Simon Daniel Robinson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1179373/publications.pdf>

Version: 2024-02-01

67
papers

3,411
citations

136950

32
h-index

149698

56
g-index

72
all docs

72
docs citations

72
times ranked

5122
citing authors

#	ARTICLE	IF	CITATIONS
1	Anterior temporal lobe degeneration produces widespread network-driven dysfunction. <i>Brain</i> , 2013, 136, 2979-2991.	7.6	184
2	Amygdala activation and facial expressions: Explicit emotion discrimination versus implicit emotion processing. <i>Neuropsychologia</i> , 2007, 45, 2369-2377.	1.6	171
3	Altered reward processing in the nucleus accumbens and mesial prefrontal cortex of patients with posttraumatic stress disorder. <i>Neuropsychologia</i> , 2008, 46, 2836-2844.	1.6	169
4	Amygdala activity to fear and anger in healthy young males is associated with testosterone. <i>Psychoneuroendocrinology</i> , 2009, 34, 687-693.	2.7	166
5	Facial emotion recognition and amygdala activation are associated with menstrual cycle phase. <i>Psychoneuroendocrinology</i> , 2008, 33, 1031-1040.	2.7	156
6	A resting state network in the motor control circuit of the basal ganglia. <i>BMC Neuroscience</i> , 2009, 10, 137.	1.9	134
7	The influence of brain iron and myelin on magnetic susceptibility and effective transverse relaxation - A biochemical and histological validation study. <i>NeuroImage</i> , 2018, 179, 117-133.	4.2	129
8	An illustrated comparison of processing methods for phase MRI and QSM: removal of background field contributions from sources outside the region of interest. <i>NMR in Biomedicine</i> , 2017, 30, e3604.	2.8	124
9	An illustrated comparison of processing methods for MR phase imaging and QSM: combining array coil signals and phase unwrapping. <i>NMR in Biomedicine</i> , 2017, 30, e3601.	2.8	124
10	The selection of intended actions and the observation of others' actions: A time-resolved fMRI study. <i>NeuroImage</i> , 2006, 29, 1294-1302.	4.2	123
11	Key clinical benefits of neuroimaging at 7 T. <i>NeuroImage</i> , 2018, 168, 477-489.	4.2	113
12	Amygdala activation at 3T in response to human and avatar facial expressions of emotions. <i>Journal of Neuroscience Methods</i> , 2007, 161, 126-133.	2.5	110
13	Clinical fMRI: Evidence for a 7T benefit over 3T. <i>NeuroImage</i> , 2011, 57, 1015-1021.	4.2	110
14	A comparison of PET imaging characteristics of various copper radioisotopes. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2005, 32, 1473-1480.	6.4	82
15	Comparison of Routine Brain Imaging at 3 T and 7 T. <i>Investigative Radiology</i> , 2016, 51, 469-482.	6.2	82
16	Combining phase images from multi-channel RF coils using 3D phase offset maps derived from a dual-echo scan. <i>Magnetic Resonance in Medicine</i> , 2011, 65, 1638-1648.	3.0	81
17	General and specific responsiveness of the amygdala during explicit emotion recognition in females and males. <i>BMC Neuroscience</i> , 2009, 10, 91.	1.9	76
18	Computationally Efficient Combination of Multi-channel Phase Data From Multi-echo Acquisitions (ASPIRE). <i>Magnetic Resonance in Medicine</i> , 2018, 79, 2996-3006.	3.0	72

#	ARTICLE	IF	CITATIONS
19	A Neuronal Basis for Task-Negative Responses in the Human Brain. <i>Cerebral Cortex</i> , 2011, 21, 821-830.	2.9	71
20	Quantitative Sodium MR Imaging at 7 T: Initial Results and Comparison with Diffusion-weighted Imaging in Patients with Breast Tumors. <i>Radiology</i> , 2016, 280, 39-48.	7.3	69
21	Robust field map generation using a triple-echo acquisition. <i>Journal of Magnetic Resonance Imaging</i> , 2004, 20, 730-734.	3.4	59
22	A method for the dynamic correction of B ₀ -related distortions in single-echo EPI at 7 T. <i>NeuroImage</i> , 2018, 168, 321-331.	4.2	57
23	Clinical applications at ultrahigh field (7T). Where does it make the difference?. <i>NMR in Biomedicine</i> , 2016, 29, 1316-1334.	2.8	56
24	Comparing localized and nonlocalized dynamic ³¹ P magnetic resonance spectroscopy in exercising muscle at 7T. <i>Magnetic Resonance in Medicine</i> , 2012, 68, 1713-1723.	3.0	55
25	Amygdala activation during recognition of emotions in a foreign ethnic group is associated with duration of stay. <i>Social Neuroscience</i> , 2009, 4, 294-307.	1.3	50
26	Combining phase images from array coils using a short echo time reference scan (COMPOSER). <i>Magnetic Resonance in Medicine</i> , 2017, 77, 318-327.	3.0	49
27	Phase unwrapping with a rapid opensource minimum spanning tree algorithm (ROMEIO). <i>Magnetic Resonance in Medicine</i> , 2021, 85, 2294-2308.	3.0	48
28	A method for unwrapping highly wrapped multi-echo phase images at very high field: UMPIRE. <i>Magnetic Resonance in Medicine</i> , 2014, 72, 80-92.	3.0	46
29	The impact of EPI voxel size on SNR and BOLD sensitivity in the anterior medio-temporal lobe: a comparative group study of deactivation of the Default Mode. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2008, 21, 279-290.	2.0	45
30	Distinct Neural Substrates for Semantic Knowledge and Naming in the Temporoparietal Network. <i>Cerebral Cortex</i> , 2012, 22, 2217-2226.	2.9	45
31	<i>B₀</i> mapping with multi-channel RF coils at high field. <i>Magnetic Resonance in Medicine</i> , 2011, 66, 976-988.	3.0	44
32	Culture but not gender modulates amygdala activation during explicit emotion recognition. <i>BMC Neuroscience</i> , 2012, 13, 54.	1.9	35
33	Dynamic PCr and pH imaging of human calf muscles during exercise and recovery using ³¹ P gradient-echo MRI at 7 Tesla. <i>Magnetic Resonance in Medicine</i> , 2016, 75, 2324-2331.	3.0	31
34	The traveling heads 2.0: Multicenter reproducibility of quantitative imaging methods at 7 Tesla. <i>NeuroImage</i> , 2021, 232, 117910.	4.2	31
35	The traveling heads: multicenter brain imaging at 7 Tesla. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2016, 29, 399-415.	2.0	26
36	Imaging the changing role of feedback during learning in decision-making. <i>NeuroImage</i> , 2007, 37, 1474-1486.	4.2	24

#	ARTICLE	IF	CITATIONS
37	Intra-session and inter-subject variability of 3D FID-MRSI using single-echo volumetric EPI navigators at 3T. <i>Magnetic Resonance in Medicine</i> , 2020, 83, 1920-1929.	3.0	23
38	The influence of spatial resolution on the spectral quality and quantification accuracy of whole-brain MRSI at 1.5T, 3T, 7T, and 9.4T. <i>Magnetic Resonance in Medicine</i> , 2019, 82, 551-565.	3.0	22
39	Correcting dynamic distortions in 7T echo planar imaging using a jittered echo time sequence. <i>Magnetic Resonance in Medicine</i> , 2016, 76, 1388-1399.	3.0	20
40	A comparison of static and dynamic B_0 mapping methods for correction of CEST MRI in the presence of temporal B_0 field variations. <i>Magnetic Resonance in Medicine</i> , 2019, 82, 633-646.	3.0	19
41	Multiparametric Quantitative Brain MRI in Neurological and Hepatic Forms of Wilson's Disease. <i>Journal of Magnetic Resonance Imaging</i> , 2020, 51, 1829-1835.	3.4	19
42	Improved susceptibility weighted imaging at ultra-high field using bipolar multi-echo acquisition and optimized image processing: CLEAR-SWI. <i>NeuroImage</i> , 2021, 237, 118175.	4.2	19
43	The challenge of bias-free coil combination for quantitative susceptibility mapping at ultra-high field. <i>Magnetic Resonance in Medicine</i> , 2018, 79, 97-107.	3.0	17
44	Applying Independent Component Analysis to Clinical fMRI at 7T. <i>Frontiers in Human Neuroscience</i> , 2013, 7, 496.	2.0	16
45	The clinical relevance of distortion correction in presurgical fMRI at 7 T. <i>NeuroImage</i> , 2018, 168, 490-498.	4.2	16
46	Brain Iron and Metabolic Abnormalities in C19orf12 Mutation Carriers: A 7.0 Tesla MRI Study in Mitochondrial Membrane Protein-associated Neurodegeneration. <i>Movement Disorders</i> , 2020, 35, 142-150.	3.9	16
47	fMRI of the Emotions: Towards an Improved Understanding of Amygdala Function. <i>Current Medical Imaging</i> , 2005, 1, 115-129.	0.8	14
48	Microvessels may Confound the "Swallow Tail Sign" in Normal Aged Midbrains: A Postmortem 7 T SWI-MRI Study. <i>Journal of Neuroimaging</i> , 2019, 29, 65-69.	2.0	14
49	Beware detrending: Optimal preprocessing pipeline for low-frequency fluctuation analysis. <i>Human Brain Mapping</i> , 2019, 40, 1571-1582.	3.6	14
50	Reinforcement and Punishment Shape the Learning Dynamics in fMRI Neurofeedback. <i>Frontiers in Human Neuroscience</i> , 2020, 14, 304.	2.0	14
51	In vivo phase imaging of human epiphyseal cartilage at 7 T. <i>Magnetic Resonance in Medicine</i> , 2018, 79, 2149-2155.	3.0	12
52	Comparing the Microvascular Specificity of the 3- and 7-T BOLD Response Using ICA and Susceptibility-Weighted Imaging. <i>Frontiers in Human Neuroscience</i> , 2013, 7, 474.	2.0	11
53	Improving sensitivity, specificity, and reproducibility of individual brainstem activation. <i>Brain Structure and Function</i> , 2019, 224, 2823-2838.	2.3	11
54	QSMxT: Robust masking and artifact reduction for quantitative susceptibility mapping. <i>Magnetic Resonance in Medicine</i> , 2022, 87, 1289-1300.	3.0	11

#	ARTICLE	IF	CITATIONS
55	ICA of fMRI Studies: New Approaches and Cutting Edge Applications. <i>Frontiers in Human Neuroscience</i> , 2013, 7, 724.	2.0	10
56	Vessel architecture in human knee cartilage in children: an in vivo susceptibility-weighted imaging study at 7 T. <i>European Radiology</i> , 2018, 28, 3384-3392.	4.5	8
57	Simultaneous Multiple Resonance Frequency imaging (SMURF): Fat-water imaging using multi-band principles. <i>Magnetic Resonance in Medicine</i> , 2021, 85, 1379-1396.	3.0	8
58	fMRI of Emotion. <i>Neuromethods</i> , 2009, , 411-456.	0.3	8
59	Quantitative susceptibility mapping of the head and neck using SMURF fat-water imaging with chemical shift and relaxation rate corrections. <i>Magnetic Resonance in Medicine</i> , 2022, 87, 1461-1479.	3.0	8
60	Phase-based masking for quantitative susceptibility mapping of the human brain at 9.4T. <i>Magnetic Resonance in Medicine</i> , 2022, 88, 2267-2276.	3.0	7
61	Improving the clinical potential of ultra-high field fMRI using a model-free analysis method based on response consistency. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2016, 29, 435-449.	2.0	6
62	Robust presurgical functional MRI at 7 T using response consistency. <i>Human Brain Mapping</i> , 2017, 38, 3163-3174.	3.6	5
63	Multi-echo GRE imaging of knee cartilage. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 45, 1502-1513.	3.4	4
64	Post Mortem Validation of MRI-Identified Veins on the Surface of the Cerebral Cortex as Potential Landmarks for Neurosurgery. <i>Frontiers in Neuroscience</i> , 2017, 11, 355.	2.8	4
65	The Impact of Echo Time Shifts and Temporal Signal Fluctuations on BOLD Sensitivity in Presurgical Planning at 7 T. <i>Investigative Radiology</i> , 2019, 54, 340-348.	6.2	3
66	fMRI of Emotion. <i>Neuromethods</i> , 2016, , 451-494.	0.3	1
67	Feasibility of Hepatic Fat Quantification Using Proton Density Fat Fraction by Multi-Echo Chemical-Shift-Encoded MRI at 7T. <i>Frontiers in Physics</i> , 2021, 9, 665562.	2.1	0