

# Mogens Groenvold

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11861854/publications.pdf>

Version: 2024-02-01

58  
papers

3,871  
citations

186265

28  
h-index

149698

56  
g-index

58  
all docs

58  
docs citations

58  
times ranked

4626  
citing authors

#	ARTICLE	IF	CITATIONS
1	The development of the EORTC QLQ-C15-PAL: A shortened questionnaire for cancer patients in palliative care. <i>European Journal of Cancer</i> , 2006, 42, 55-64.	2.8	502
2	Validation of the EORTC QLQ-C30 quality of life questionnaire through combined qualitative and quantitative assessment of patient-observer agreement. <i>Journal of Clinical Epidemiology</i> , 1997, 50, 441-450.	5.0	388
3	Replication and validation of higher order models demonstrated that a summary score for the EORTC QLQ-C30 is robust. <i>Journal of Clinical Epidemiology</i> , 2016, 69, 79-88.	5.0	306
4	Tests of Data Quality, Scaling Assumptions, and Reliability of the Danish SF-36. <i>Journal of Clinical Epidemiology</i> , 1998, 51, 1001-1011.	5.0	223
5	Quality of life in patients with benign thyroid disorders. A review. <i>European Journal of Endocrinology</i> , 2006, 154, 501-510.	3.7	184
6	Health related quality of life in a nationally representative sample of haematological patients. <i>European Journal of Haematology</i> , 2009, 83, 139-148.	2.2	176
7	Validity and reliability of the novel thyroid-specific quality of life questionnaire, ThyPRO. <i>European Journal of Endocrinology</i> , 2010, 162, 161-167.	3.7	163
8	Does the Medical Record Cover the Symptoms Experienced by Cancer Patients Receiving Palliative Care? A Comparison of the Record and Patient Self-Rating. <i>Journal of Pain and Symptom Management</i> , 2001, 21, 189-196.	1.2	128
9	The Thyroid-Related Quality of Life Measure ThyPRO Has Good Responsiveness and Ability to Detect Relevant Treatment Effects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, 3708-3717.	3.6	123
10	Assessing health-related quality of life in palliative care: Comparing patient and physician assessments. <i>European Journal of Cancer</i> , 2006, 42, 1159-1166.	2.8	120
11	Self-assessment in cancer patients referred to palliative care. <i>Cancer</i> , 2002, 94, 512-520.	4.1	119
12	Validation of a questionnaire for self-assessment of sexual function and vaginal changes after gynaecological cancer. <i>Psycho-Oncology</i> , 2004, 13, 577-592.	2.3	106
13	Anxiety and Depression in Breast Cancer Patients at Low Risk of Recurrence Compared with the General Population. <i>Journal of Clinical Epidemiology</i> , 1999, 52, 523-530.	5.0	88
14	Establishing construct validity for the thyroid-specific patient reported outcome measure (ThyPRO): an initial examination. <i>Quality of Life Research</i> , 2009, 18, 483-496.	3.1	84
15	Development of a Short Version of the Thyroid-Related Patient-Reported Outcome ThyPRO. <i>Thyroid</i> , 2015, 25, 1069-1079.	4.5	82
16	Which Domains of Thyroid-Related Quality of Life Are Most Relevant? Patients and Clinicians Provide Complementary Perspectives. <i>Thyroid</i> , 2007, 17, 647-654.	4.5	78
17	Improving a newly developed patient-reported outcome for thyroid patients, using cognitive interviewing. <i>Quality of Life Research</i> , 2008, 17, 1009-1017.	3.1	65
18	The chronic autoimmune thyroiditis quality of life selenium trial (CATALYST): study protocol for a randomized controlled trial. <i>Trials</i> , 2014, 15, 115.	1.6	60

#	ARTICLE	IF	CITATIONS
19	Is Thyroid Autoimmunity per se a Determinant of Quality of Life in Patients with Autoimmune Hypothyroidism?. <i>European Thyroid Journal</i> , 2012, 1, 186-192.	2.4	59
20	Test for item bias in a quality of life questionnaire. <i>Journal of Clinical Epidemiology</i> , 1995, 48, 805-816.	5.0	56
21	Quality-of-Life Impairments Persist Six Months After Treatment of Graves' Hyperthyroidism and Toxic Nodular Goiter: A Prospective Cohort Study. <i>Thyroid</i> , 2016, 26, 1010-1018.	4.5	55
22	Quality of Life in Patients with Benign Nontoxic Goiter: Impact of Disease and Treatment Response, and Comparison with the General Population. <i>Thyroid</i> , 2015, 25, 284-291.	4.5	54
23	Comparing higher order models for the EORTC QLQ-C30. <i>Quality of Life Research</i> , 2012, 21, 1607-1617.	3.1	52
24	Cross-cultural validity of the thyroid-specific quality-of-life patient-reported outcome measure, ThyPRO. <i>Quality of Life Research</i> , 2015, 24, 769-780.	3.1	52
25	Confirmatory factor analysis of the thyroid-related quality of life questionnaire ThyPRO. <i>Health and Quality of Life Outcomes</i> , 2014, 12, 126.	2.4	41
26	Item response theory was used to shorten EORTC QLQ-C30 scales for use in palliative care. <i>Journal of Clinical Epidemiology</i> , 2006, 59, 36-44.	5.0	40
27	Development and Initial Validation of the Three-Levels-of-Needs Questionnaire for Self-Assessment of Palliative Needs in Patients with Cancer. <i>Journal of Pain and Symptom Management</i> , 2011, 41, 1025-1039.	1.2	35
28	Is admittance to specialised palliative care among cancer patients related to sex, age and cancer diagnosis? A nation-wide study from the Danish Palliative Care Database (DPD). <i>BMC Palliative Care</i> , 2017, 16, 21.	1.8	31
29	Are different groups of cancer patients offered rehabilitation to the same extent? A report from the population-based study "The Cancer Patient's World". <i>Supportive Care in Cancer</i> , 2012, 20, 1089-1100.	2.2	30
30	Few items in the thyroid-related quality of life instrument ThyPRO exhibited differential item functioning. <i>Quality of Life Research</i> , 2014, 23, 327-338.	3.1	29
31	Breast cancer patients on adjuvant chemotherapy report a wide range of problems not identified by health-care staff. <i>Breast Cancer Research and Treatment</i> , 2007, 103, 185-195.	2.5	27
32	Chemotherapy versus ovarian ablation as adjuvant therapy for breast cancer: impact on health-related quality of life in a randomized trial. <i>Breast Cancer Research and Treatment</i> , 2006, 98, 275-284.	2.5	22
33	Danish Palliative Care Database. <i>Clinical Epidemiology</i> , 2016, Volume 8, 637-643.	3.0	21
34	Prevalence and possible predictors of sexual dysfunction and self-reported needs related to the sexual life of advanced cancer patients. <i>Acta Oncologica</i> , 2019, 58, 769-775.	1.8	21
35	A new model of early, integrated palliative care: palliative rehabilitation for newly diagnosed patients with non-resectable cancer. <i>Supportive Care in Cancer</i> , 2019, 27, 3291-3300.	2.2	20
36	Admittance to specialized palliative care (SPC) of patients with an assessed need: a study from the Danish palliative care database (DPD). <i>Acta Oncologica</i> , 2017, 56, 1210-1217.	1.8	18

#	ARTICLE	IF	CITATIONS
37	Validation of the European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire Core 30 Summary Score in Patients With Hematologic Malignancies. <i>Value in Health</i> , 2019, 22, 1303-1310.	0.3	18
38	Validation of a questionnaire for self-rating of urological and gynaecological morbidity after treatment of gynaecological cancer. <i>Radiotherapy and Oncology</i> , 2002, 65, 29-38.	0.6	17
39	A parallel-group randomized clinical trial of individually tailored, multidisciplinary, palliative rehabilitation for patients with newly diagnosed advanced cancer: the Pal-Rehab study protocol. <i>BMC Cancer</i> , 2017, 17, 560.	2.6	17
40	Determining minimal important change for the thyroid-related quality of life questionnaire ThyPRO. <i>Endocrine Connections</i> , 2021, 10, 316-324.	1.9	17
41	Age, cancer site and gender associations with symptoms and problems in specialised palliative care: a large, nationwide, register-based study. <i>BMJ Supportive and Palliative Care</i> , 2019, , bmjspcare-2019-001880.	1.6	16
42	Quality of life in patients with diffuse large B-cell lymphoma treated with dose-dense chemotherapy is only affected temporarily. <i>Leukemia and Lymphoma</i> , 2011, 52, 400-408.	1.3	15
43	Measuring health-related quality of life in colorectal cancer patients: systematic review of measurement properties of the EORTC QLQ-CR29. <i>Supportive Care in Cancer</i> , 2019, 27, 2395-2412.	2.2	13
44	Exploratory analyses of the Danish Palliative Care Trial (DanPaCT): a randomized trial of early specialized palliative care plus standard care versus standard care in advanced cancer patients. <i>Supportive Care in Cancer</i> , 2020, 28, 2145-2155.	2.2	13
45	Quality of life and symptoms in patients with malignant diseases admitted to a comprehensive cancer centre. <i>Supportive Care in Cancer</i> , 2014, 22, 1843-1849.	2.2	11
46	Comparing three different approaches to the measurement of needs concerning fatigue in patients with advanced cancer. <i>Quality of Life Research</i> , 2015, 24, 2231-2238.	3.1	11
47	Patients with Benign Thyroid Diseases Experience an Impaired Sex Life. <i>Thyroid</i> , 2018, 28, 1261-1269.	4.5	10
48	Which symptoms and problems do advanced cancer patients admitted to specialized palliative care report in addition to those included in the EORTC QLQ-C15-PAL? A register-based national study. <i>Supportive Care in Cancer</i> , 2020, 28, 1725-1735.	2.2	10
49	Using method triangulation to validate a new instrument (CPWQ-com) assessing cancer patients' satisfaction with communication. <i>Cancer Epidemiology</i> , 2012, 36, 29-35.	1.9	8
50	Satisfaction with information provided to Danish cancer patients: Validation and survey results. <i>Patient Education and Counseling</i> , 2013, 93, 239-247.	2.2	7
51	Adaptation and cross-cultural validation of the Spanish version of the Thyroid-Related Quality-of-Life Patient-Reported Outcome questionnaire. <i>Endocrinología, Diabetes Y Nutrición</i> , 2018, 65, 500-507.	0.3	7
52	Nausea in advanced cancer: relationships between intensity, burden, and the need for help. <i>Supportive Care in Cancer</i> , 2019, 27, 265-273.	2.2	7
53	Using mixed methods to assess how cancer patients' needs in relation to their relatives are met in the Danish health care system: a report from the population-based study 'The Cancer Patient's World'. <i>Supportive Care in Cancer</i> , 2012, 20, 3211-3220.	2.2	6
54	How does pain experience relate to the need for pain relief? A secondary exploratory analysis in a large sample of cancer patients. <i>Supportive Care in Cancer</i> , 2016, 24, 4187-4195.	2.2	4

#	ARTICLE	IF	CITATIONS
55	Change in health-related quality of life over 1 month in cancer patients with high initial levels of symptoms and problems. <i>Quality of Life Research</i> , 2016, 25, 2669-2674.	3.1	4
56	Investigating the response scale of the EORTC QLQ-C30 in German cancer patients and a population survey. <i>Health and Quality of Life Outcomes</i> , 2021, 19, 235.	2.4	2
57	Adaptation and cross-cultural validation of the Spanish version of the Thyroid-Related Quality-of-Life Patient-Reported Outcome questionnaire. <i>Endocrinología y Nutrición (English Ed)</i> , 2018, 65, 500-507.	0.2	0
58	Should analyses of large, national palliative care data sets with patient reported outcomes (PROs) be restricted to services with high patient participation? A register-based study. <i>BMC Palliative Care</i> , 2020, 19, 89.	1.8	0