

# Bandit Thinkhamrop

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11905388/publications.pdf>

Version: 2024-02-01

15  
papers

674  
citations

687363

13  
h-index

996975

15  
g-index

15  
all docs

15  
docs citations

15  
times ranked

579  
citing authors

#	ARTICLE	IF	CITATIONS
1	Advanced periductal fibrosis from infection with the carcinogenic human liver fluke <i>Opisthorchis viverrini</i> correlates with elevated levels of interleukin-6. <i>Hepatology</i> , 2009, 50, 1273-1281.	7.3	145
2	Ultrasonography assessment of hepatobiliary abnormalities in 3359 subjects with <i>Opisthorchis viverrini</i> infection in endemic areas of Thailand. <i>Parasitology International</i> , 2012, 61, 208-211.	1.3	102
3	Elevated Plasma IL-6 Associates with Increased Risk of Advanced Fibrosis and Cholangiocarcinoma in Individuals Infected by <i>Opisthorchis viverrini</i> . <i>PLoS Neglected Tropical Diseases</i> , 2012, 6, e1654.	3.0	96
4	Cohort profile: cholangiocarcinoma screening and care program (CASCAP). <i>BMC Cancer</i> , 2015, 15, 459.	2.6	93
5	A Comprehensive Public Health Conceptual Framework and Strategy to Effectively Combat Cholangiocarcinoma in Thailand. <i>PLoS Neglected Tropical Diseases</i> , 2016, 10, e0004293.	3.0	51
6	Current Perspectives on Opisthorchiasis Control and Cholangiocarcinoma Detection in Southeast Asia. <i>Frontiers in Medicine</i> , 2018, 5, 117.	2.6	51
7	Outcome of curative resection for perihilar cholangiocarcinoma in Northeast Thailand. <i>World Journal of Gastrointestinal Oncology</i> , 2015, 7, 503.	2.0	24
8	Association between Diabetes Mellitus and Fatty Liver Based on Ultrasonography Screening in the World's Highest Cholangiocarcinoma Incidence Region, Northeast Thailand. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 16, 3931-3936.	1.2	19
9	Association between periductal fibrosis and bile duct dilatation among a population at high risk of cholangiocarcinoma: a cross-sectional study of cholangiocarcinoma screening in Northeast Thailand. <i>BMJ Open</i> , 2019, 9, e023217.	1.9	17
10	Survival after surgery among patients with cholangiocarcinoma in Northeast Thailand according to anatomical and morphological classification. <i>BMC Cancer</i> , 2021, 21, 497.	2.6	16
11	Changing patterns of prevalence in <i>Opisthorchis viverrini</i> sensu lato infection in children and adolescents in northeast Thailand. <i>Acta Tropica</i> , 2016, 164, 469-472.	2.0	15
12	Teleconsultation ultrasonography: a new weapon to combat cholangiocarcinoma. <i>ESMO Open</i> , 2017, 2, e000231.	4.5	15
13	The Socioeconomic Burden of Cholangiocarcinoma Associated With <i>Opisthorchis viverrini</i> Sensu Lato Infection in Northeast Thailand. <i>Advances in Parasitology</i> , 2018, 102, 141-163.	3.2	13
14	A comparison of the proportion of early stage cholangiocarcinoma found in an ultrasound-screening program compared to walk-in patients. <i>Hpb</i> , 2020, 22, 874-883.	0.3	11
15	Spatial analysis of hepatobiliary abnormalities in a population at high-risk of cholangiocarcinoma in Thailand. <i>Scientific Reports</i> , 2020, 10, 16855.	3.3	6