## V?ronica C Arancibia

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1193624/publications.pdf

Version: 2024-02-01

394421 434195 65 1,182 19 31 citations g-index h-index papers 65 65 65 1624 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	A electrochemical biosensor for As(III) detection based on the catalytic activity of Alcaligenes faecalis immobilized on a gold nanoparticle–modified screen–printed carbon electrode. Talanta, 2021, 223, 121702.	5.5	25
2	Determination of molybdenum(VI) via adsorptive stripping voltammetry using an exâ€'situ bismuth screenâ€'printed carbon electrode. Microchemical Journal, 2020, 154, 104589.	4.5	7
3	Development of a fast and sensitive method for the determination of As(III) at trace levels in urine by differential pulse anodic voltammetry using a simple graphene screen–printed electrode. Microchemical Journal, 2020, 159, 105393.	4.5	10
4	Determination of Pb(II) and Cd(II) via anodic stripping voltammetry using an in-situ bismuth film electrode. Increasing the sensitivity of the method by the presence of Alizarin Red S. Microchemical Journal, 2020, 159, 105373.	4.5	15
5	Determination of a natural $(17\hat{1}^2$ -estradiol) and a synthetic $(17\hat{1}\pm$ -ethinylestradiol) hormones in pharmaceutical formulations and urine by adsorptive stripping voltammetry. Sensors and Actuators B: Chemical, 2019, 297, 126728.	7.8	17
6	Development of a microcomposite with single-walled carbon nanotubes and Nd2O3 for determination of paracetamol in pharmaceutical dosage by adsorptive voltammetry. Journal of Pharmaceutical Analysis, 2019, 9, 62-69.	5.3	35
7	Determination of Se(IV) concentration via cathodic stripping voltammetry in the presence of Cu(II) ions and ammonium diethyl dithiophosphate. Analytica Chimica Acta, 2019, 1048, 22-30.	5.4	6
8	Fast and highly sensitive method for molybdenum(VI) determination by catalytic adsorptive stripping voltammetry. Sensors and Actuators B: Chemical, 2018, 258, 612-620.	7.8	5
9	Electrocomposite Developed with Chitosan and Ionic Liquids Using Screen-Printed Carbon Electrodes Useful to Detect Rutin in Tropical Fruits. Sensors, 2018, 18, 2934.	3.8	2
10	Detection of SO <sub>2</sub> derivatives using a new chalco-coumarin derivative in cationic micellar media: application to real samples. RSC Advances, 2018, 8, 31261-31266.	3.6	11
11	A new strategy for the modification of a carbon paste electrode with carrageenan hydrogel for a sensitive and selective determination of arsenic in natural waters. Talanta, 2018, 187, 259-264.	5.5	12
12	New fluorescent turn-off probes for highly sensitive and selective detection of SO2 derivatives in a micellar media. Sensors and Actuators B: Chemical, 2017, 238, 578-587.	7.8	33
13	Adsorptive Stripping Voltammetric Determination of Amaranth and Tartrazine in Drinks and Gelatins Using a Screen-Printed Carbon Electrode. Sensors, 2017, 17, 2665.	3.8	21
14	Adsorptive Stripping Voltammetric Determination of Morin in Tea Infusions and Chocolate Drinks on a Gold Electrode. Effect of Cetylpyridinium Bromide on the Sensitivity of the Method. International Journal of Electrochemical Science, 2017, , 9408-9417.	1.3	12
15	ANTIPROLIFERATIVE ACTIVITY OF NEW 6-BROMINE DERIVATIVES OF 7-ANILINO-1-ARYLISOQUINOLINEQUINONES. Journal of the Chilean Chemical Society, 2016, 61, 3191-3194.	1.2	6
16	Determination of Sb(III) using an ex-situ bismuth screen-printed carbon electrode by adsorptive stripping voltammetry. Talanta, 2016, 155, 21-27.	5.5	33
17	Determination of Sudan I in drinks containing Sunset yellow by adsorptive stripping voltammetry. Food Chemistry, 2016, 212, 807-813.	8.2	24
18	Determination of arsenic in the presence of copper by adsorptive stripping voltammetry using pyrrolidine dithiocarbamate or diethyl dithiophosphate as chelating-adsorbent agents. Effect of CPB on the sensitivity of the method. Microchemical Journal, 2016, 126, 70-75.	4.5	8

#	Article	IF	Citations
19	Highly sensitive determination of vanadium (V) by catalytic adsorptive stripping voltammetry. Substituent effect on sensitivity III. Sensors and Actuators B: Chemical, 2016, 224, 772-779.	7.8	13
20	A Comparative Study of 8-Hydroxyquinoline and 8-Hydroxyquinoline-5-sulfonic Acid for Antimony(III) Determination by AdSV. Substituent Effect on Sensitivity II. Electroanalysis, 2015, 27, 1262-1267.	2.9	6
21	Levels of copper in sweeteners, sugar, tea, coffee and mate infusions. Determination by adsorptive stripping voltammetry in the presence of α-lipoic acid. Microchemical Journal, 2015, 119, 11-16.	4.5	19
22	Synthesis, Half-Wave Potentials and Antiproliferative Activity of 1-Aryl-substituted Aminoisoquinolinequinones. Molecules, 2014, 19, 726-739.	3.8	10
23	Simultaneous determination of Pb2+, Cd2+ and Zn2+ by adsorptive stripping voltammetry using Clioquinol as a chelating-adsorbent agent. Journal of Electroanalytical Chemistry, 2014, 729, 9-14.	3.8	24
24	Simultaneous Determination of Antimony(III) and Molybdenum(VI) by Adsorptive Stripping Voltammetry Using Quercetin as Complexing Agent. Electroanalysis, 2013, 25, 439-447.	2.9	12
25	Ex situ prepared nafion-coated antimony film electrode for adsorptive stripping voltammetry of model metal ions in the presence of pyrogallol red. Sensors and Actuators B: Chemical, 2013, 182, 368-373.	7.8	20
26	Synthesis and structural characterization of new 2-bromo-1,3-bis(triazol-1-ylmethyl)benzene ligands. Study of their behavior as complexing agents for determination of nickel(II) by adsorptive stripping voltammetry. Journal of Coordination Chemistry, 2013, 66, 592-601.	2.2	3
27	High sensitivity adsorptive stripping voltammetric method for antimony(III) determination in the presence of quercetin-5′-sulfonic acid. Substituent effect on sensitivity. Sensors and Actuators B: Chemical, 2013, 185, 560-567.	7.8	24
28	Synthesis and in Vitro Antiproliferative Activity of New Phenylaminoisoquinolinequinones against Cancer Cell Lines. Molecules, 2013, 18, 721-734.	3.8	8
29	Synthesis, characterization, and electrochemical study of diiron organometallic derivatives of 2,6-dibutyl-4,8-dimethyl-1,5-dihydro- <i>s</i> -indacene. Canadian Journal of Chemistry, 2013, 91, 727-731.	1.1	1
30	Nafion–mercury coated film electrode for the adsorptive stripping voltammetric determination of lead and cadmium in the presence of pyrogallol red. Talanta, 2012, 99, 119-124.	5.5	30
31	Adsorptive stripping voltammetry of nickel with 1-nitroso-2-napthol using a bismuth film electrode. Talanta, 2011, 85, 2316-2319.	5.5	23
32	Stripping voltammetric determination of cadmium in sea water using a carbon paste electrode modified with alginic acid from brown algae. Journal of the Brazilian Chemical Society, 2010, 21, 1688-1691.	0.6	7
33	Studies on quinones. Part 45: Novel 7-aminoisoquinoline-5,8-quinone derivatives with antitumor properties on cancer cell lines. Bioorganic and Medicinal Chemistry, 2009, 17, 2894-2901.	3.0	46
34	Determination of lead in the presence of morin- $5\hat{a}\in^2$ -sulfonic acid and sodium dodecyl sulfate by adsorptive stripping voltammetry. Talanta, 2009, 80, 184-188.	5.5	28
35	Determination of iron in water samples by adsorptive stripping voltammetry with a bismuth film electrode in the presence of 1-(2-piridylazo)-2-naphthol. Talanta, 2008, 75, 973-977.	5.5	50
36	Determination of aluminium in water samples by adsorptive cathodic stripping voltammetry in the presence of pyrogallol red and a quaternary ammonium salt. Talanta, 2007, 73, 546-552.	5.5	44

#	Article	IF	CITATIONS
37	Stoichiometry and conditional stability constants of Cu(II) or Zn(II) clioquinol complexes; implications for Alzheimer's and Huntington's disease therapy. NeuroToxicology, 2007, 28, 445-449.	3.0	75
38	Synthesis, structure and properties of ruthenium(II) complexes with quinolinedione derivatives as chelate ligands: Crystal structure of $[Ru(CO)2Cl2(6-methoxybenzo[g]quinoline-5,10-dione)]$ . Polyhedron, 2007, 26, 5527-5532.	2.2	13
39	Extraction of arsenic as the diethyl dithiophosphate complex with supercritical fluid and quantitation by cathodic stripping voltammetry. Talanta, 2006, 68, 1567-1573.	5.5	17
40	Evaluation of Powdered Infant Formula Milk as Chelating Agent for Copper under Simulated Gastric Conditions of a Baby's Stomach. Analytical Sciences, 2006, 22, 1197-1200.	1.6	3
41	Synthesis, characterization and structure of diiron organometallic derivatives of 2,9-dimethyl-1,10-dihydro-dicyclopenta[a,h]naphthalene. Journal of Organometallic Chemistry, 2005, 690, 1340-1349.	1.8	11
42	DETERMINATION OF PARTITION COEFFICIENT OF BENZO[b]THIOPHENES BY REVERSED-PHASE HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY. Heterocyclic Communications, 2005, 11, .	1.2	0
43	Supercritical fluid extraction of cadmium as Cd–oxine complex from human hair. Analytica Chimica Acta, 2004, 502, 189-194.	5.4	10
44	Quantitative extraction of sulfonamides in meats by supercritical methanol-modified carbon dioxide: A foray into real-world sampling. Journal of Separation Science, 2003, 26, 1710-1716.	2.5	17
45	Determination of chromium in urine samples by complexation–supercritical fluid extraction and liquid or gas chromatography. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2003, 785, 303-309.	2.3	78
46	Characterization of copper in uterine fluids of patients who use the copper T-380A intrauterine device. Clinica Chimica Acta, 2003, 332, 69-78.	1.1	50
47	Extraction of nitrofurantoin and its toxic metabolite from urine by supercritical fluids. Quantitation by high performance liquid chromatography with UV detection. Talanta, 2003, 61, 377-383.	5.5	38
48	Dithioethers as ligands in pentamethylcyclopentadienylrhodium(III) and iridium(III) complexes. Crystal structures of $[\{(\hat{l}\cdot 5-C5Me5)Rh\}2(\hat{l}\cdot 4-Cl)2\{\hat{l}\cdot 4-(MeS)2CH2\}](BF4)2\hat{A}\cdot H2O$ and $[(\hat{l}\cdot 5-C5Me5)IrCl2\{\hat{l}\cdot 1-(PhS)2CH2\}]$ . Journal of Organometallic Chemistry, 2001, 620, 256-262.	1.8	13
49	Synthesis and characterization of [Cp*Fe-dicyclopenta(a,f)naphthalene-FeCp*] and [Cp*Fe-dicyclopenta(a,f)naphthalene-FeCp*] BF4â°. Journal of Organometallic Chemistry, 2001, 620, 32-38.	1.8	11
50	SYNTHESIS AND ELECTROCHEMICAL PROPERTIES OF CYCLOPENTADIENYLIRON(II) COMPLEXES WITH BIS(DIPHENYLPHOSPHINO)AMINE AS LIGAND. Journal of Coordination Chemistry, 2001, 54, 389-400.	2.2	3
51	Selective oxidants for organometallic compounds containing a stabilising anion of highly reactive Chemistry, 2000, 601, 126-132.	1.8	76
52	Synthesis, Spectroscopic Characterization, and Crystal Structure of the Bimetallic Complex [Ni2( $\hat{l}$ /4-CO)(CO)2( $\hat{l}$ /4-NH(PPh2)2)2]. Inorganic Chemistry, 2000, 39, 1650-1654.	4.0	13
53	HEXAMETHYLBENZENERUTHENIUM(II) COMPLEXES CONTAINING BIS(DIPHENYLPHOSPHINE)AMINE AND THEIR SULPHUR OR SELENIUM DERIVATIVES AS LIGANDS. Journal of the Chilean Chemical Society, 2000, 45, .	0.1	1
54	DI AND TRIMETALLIC PtII, PtIV COMPLEXES CONTAINING DIMETHYLPHOSPHONATE GROUPS S BRIDGING LIGANDS. Journal of the Chilean Chemical Society, 1999, 44, .	0.1	0

#	Article	IF	CITATIONS
55	Title is missing!. Transition Metal Chemistry, 1997, 22, 150-155.	1.4	10
56	Tris(diphenylthiophosphinoyl)methanide as tripod ligand in rhodium(III), iridium(III) and ruthenium(II) complexes. Crystal structures of [(η5-C5Me5)Ir{η3-(SPPh2)3C-S,S′,S″}]BF4 and [(η6-MeC6H4Pri)Ru{η3-(SPPh2)3C-S,S′,S″}]BPh4. Journal of Organometallic Chemistry, 1997, 545-546, 50	1.8 7-517.	10
57	Synthesis and electrochemical properties of p-cymene-ruthenium(II) complexes with (EPPh2)2CHR (Eî—»S,) Tj ET [(Î-6-MeC6H4Pri)Ru{Î-3-(SPPh2)2CMe-C,S,S′}]PF6. Inorganica Chimica Acta, 1997, 255, 221-227.	Qq1 1 0.7 2.4	84314 rgBT 14
58	Redox chemistry of mononuclear manganese(II), binuclear manganese(III) and binuclear mixed manganese(II)-manganese(III) complexes with 3-aminopyrazine-2-carboxylate in dimethylsulphoxide. Transition Metal Chemistry, 1995, 20, 179-184.	1.4	5
59	Redox chemistry of 3,4-dihydroxy-2-benzoic acids, its oxidation products and their interaction with manganese(II) and manganese(III). Polyhedron, 1995, 14, 2933-2936.	2.2	13
60	A polarographic, voltammetric and spectroscopic study of 2-mercaptonicotinic acid and its chromium (III) complex. Analytica Chimica Acta, 1994, 298, 91-98.	5.4	9
61	Redox chemistry and spectroscopy of the manganese-quinizarine-thiosalicylic acid mixed-ligand complex in aprotic medium. Polyhedron, 1993, 12, 1733-1738.	2.2	4
62	Voltammetric and spectroscopic studies of the behaviour of manganese(II) and manganese(III) complexes with isoquinolinecarboxylic acid in aprotic medium. Polyhedron, 1993, 12, 1739-1744.	2.2	2
63	Zinc(II) complexes with the reduction products of 2,3-dimethoxy-1,4-naphthoquinone in dimethyl sulphoxide redox chemistry and spectroscopy. Polyhedron, 1993, 12, 1745-1750.	2.2	1
64	Redox chemistry of 1,4-dihydroxy-9,10-anthraquinone (quinizarine) and its manganese(II) complexes in dimethylsulphoxide. Polyhedron, 1991, 10, 1929-1937.	2.2	16
65	Manganese complexes with 2-hydroxy-3(3-methyl-2-butenyl)-1,4-naphthoquinone (Lapachol). Redox chemistry and spectroscopy in dimethylsulphoxide. Polyhedron, 1989, 8, 1407-1412.	2.2	19