

Inger Sandlie

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11940658/publications.pdf>

Version: 2024-02-01

121
papers

5,872
citations

71102

41
h-index

82547

72
g-index

126
all docs

126
docs citations

126
times ranked

6201
citing authors

#	ARTICLE	IF	CITATIONS
1	Antibody variable sequences have a pronounced effect on cellular transport and plasma half-life. <i>IScience</i> , 2022, 25, 103746.	4.1	23
2	Potent TRIM21 and complement-dependent intracellular antiviral immunity requires the IgG3 hinge. <i>Science Immunology</i> , 2022, 7, eabj1640.	11.9	14
3	The neonatal Fc receptor in mucosal immune regulation. <i>Scandinavian Journal of Immunology</i> , 2021, 93, e13017.	2.7	8
4	Antibody-mediated delivery of T-cell epitopes to antigen-presenting cells induce strong CD4 and CD8 T-cell responses. <i>Vaccine</i> , 2021, 39, 1583-1592.	3.8	0
5	A high-affinity human TCR-like antibody detects celiac disease gluten peptide-MHC complexes and inhibits T cell activation. <i>Science Immunology</i> , 2021, 6, .	11.9	15
6	Extended plasma half-life of albumin-binding domain fused human IgA upon pH-dependent albumin engagement of human FcRn <i>in vitro</i> and <i>in vivo</i> . <i>MAbs</i> , 2021, 13, 1893888.	5.2	16
7	A <i>TRAV26</i> -encoded recognition motif focuses the biased T cell response in celiac disease. <i>European Journal of Immunology</i> , 2020, 50, 142-145.	2.9	2
8	An engineered human albumin enhances half-life and transmucosal delivery when fused to protein-based biologics. <i>Science Translational Medicine</i> , 2020, 12, .	12.4	37
9	An intact C-terminal end of albumin is required for its long half-life in humans. <i>Communications Biology</i> , 2020, 3, 181.	4.4	40
10	FcRn is a CD32a coreceptor that determines susceptibility to IgG immune complex-driven autoimmunity. <i>Journal of Experimental Medicine</i> , 2020, 217, .	8.5	24
11	The Neonatal Fc Receptor (FcRn): A Misnomer?. <i>Frontiers in Immunology</i> , 2019, 10, 1540.	4.8	271
12	TRIM21-From Intracellular Immunity to Therapy. <i>Frontiers in Immunology</i> , 2019, 10, 2049.	4.8	85
13	Targeting the MHC Ligandome by Use of TCR-Like Antibodies. <i>Antibodies</i> , 2019, 8, 32.	2.5	36
14	Complement C4 Prevents Viral Infection through Capsid Inactivation. <i>Cell Host and Microbe</i> , 2019, 25, 617-629.e7.	11.0	53
15	Plasma Cells Are the Most Abundant Gluten Peptide MHC-expressing Cells in Inflamed Intestinal Tissues From Patients With Celiac Disease. <i>Gastroenterology</i> , 2019, 156, 1428-1439.e10.	1.3	61
16	Binding to nanopatterned antigens is dominated by the spatial tolerance of antibodies. <i>Nature Nanotechnology</i> , 2019, 14, 184-190.	31.5	134
17	A human endothelial cell-based recycling assay for screening of FcRn targeted molecules. <i>Nature Communications</i> , 2018, 9, 621.	12.8	59
18	Animal models for evaluation of albumin-based therapeutics. <i>Current Opinion in Chemical Engineering</i> , 2018, 19, 68-76.	7.8	13

#	ARTICLE	IF	CITATIONS
19	Human and mouse albumin bind their respective neonatal Fc receptors differently. <i>Scientific Reports</i> , 2018, 8, 14648.	3.3	42
20	Soluble T-cell receptor design influences functional yield in an E. coli chaperone-assisted expression system. <i>PLoS ONE</i> , 2018, 13, e0195868.	2.5	13
21	Hepatic FcRn regulates albumin homeostasis and susceptibility to liver injury. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E2862-E2871.	7.1	84
22	A TCR β framework α -centered codon shapes a biased T cell repertoire through direct MHC and CDR3 β interactions. <i>JCI Insight</i> , 2017, 2, .	5.0	15
23	Antibody-antigen kinetics constrain intracellular humoral immunity. <i>Scientific Reports</i> , 2016, 6, 37457.	3.3	27
24	Multivalent pIX phage display selects for distinct and improved antibody properties. <i>Scientific Reports</i> , 2016, 6, 39066.	3.3	14
25	TRIM21 Immune Signaling Is More Sensitive to Antibody Affinity Than Its Neutralization Activity. <i>Journal of Immunology</i> , 2016, 196, 3452-3459.	0.8	34
26	Enhanced FcRn-dependent transepithelial delivery of IgG by Fc-engineering and polymerization. <i>Journal of Controlled Release</i> , 2016, 223, 42-52.	9.9	25
27	The Influence of FcRn on Albumin-Fused and Targeted Drugs. , 2016, , 179-208.		1
28	<scp>TRIM</scp>21: a cytosolic Fc receptor with broad antibody isotype specificity. <i>Immunological Reviews</i> , 2015, 268, 328-339.	6.0	78
29	The role of albumin receptors in regulation of albumin homeostasis: Implications for drug delivery. <i>Journal of Controlled Release</i> , 2015, 211, 144-162.	9.9	152
30	Fc Engineering of Human IgG1 for Altered Binding to the Neonatal Fc Receptor Affects Fc Effector Functions. <i>Journal of Immunology</i> , 2015, 194, 5497-5508.	0.8	56
31	Phage Display Engineered T Cell Receptors as Tools for the Study of Tumor Peptide α -MHC Interactions. <i>Frontiers in Oncology</i> , 2015, 4, 378.	2.8	4
32	Developing the IVIG biomimetic, Hexa-Fc, for drug and vaccine applications. <i>Scientific Reports</i> , 2015, 5, 9526.	3.3	33
33	Eculizumab treatment during pregnancy does not affect the complement system activity of the newborn. <i>Immunobiology</i> , 2015, 220, 452-459.	1.9	90
34	Phage Display and Selection of Protein Ligands. , 2015, , 115-134.		0
35	Interaction with Both Domain I and III of Albumin Is Required for Optimal pH-dependent Binding to the Neonatal Fc Receptor (FcRn). <i>Journal of Biological Chemistry</i> , 2014, 289, 34583-34594.	3.4	36
36	Dissection of the Neonatal Fc Receptor (FcRn)-Albumin Interface Using Mutagenesis and Anti-FcRn Albumin-blocking Antibodies. <i>Journal of Biological Chemistry</i> , 2014, 289, 17228-17239.	3.4	38

#	ARTICLE	IF	CITATIONS
37	Extending Serum Half-life of Albumin by Engineering Neonatal Fc Receptor (FcRn) Binding. <i>Journal of Biological Chemistry</i> , 2014, 289, 13492-13502.	3.4	132
38	Unraveling the Interaction between FcRn and Albumin: Opportunities for Design of Albumin-Based Therapeutics. <i>Frontiers in Immunology</i> , 2014, 5, 682.	4.8	188
39	Chaperone-assisted thermostability engineering of a soluble T cell receptor using phage display. <i>Scientific Reports</i> , 2013, 3, 1162.	3.3	23
40	Selection of Nanobodies that Target Human Neonatal Fc Receptor. <i>Scientific Reports</i> , 2013, 3, 1118.	3.3	9
41	Single-chain Variable Fragment Albumin Fusions Bind the Neonatal Fc Receptor (FcRn) in a Species-dependent Manner. <i>Journal of Biological Chemistry</i> , 2013, 288, 24277-24285.	3.4	55
42	A series of anti-CEA/anti-DOTA bispecific antibody formats evaluated for pre-targeting: comparison of tumor uptake and blood clearance. <i>Protein Engineering, Design and Selection</i> , 2013, 26, 187-193.	2.1	30
43	Effective Phagocytosis of Low Her2 Tumor Cell Lines with Engineered, Aglycosylated IgG Displaying High Fcγ3R1a Affinity and Selectivity. <i>ACS Chemical Biology</i> , 2013, 8, 368-375.	3.4	61
44	Chimeric Anti-CD14 IGG2/4 Hybrid Antibodies for Therapeutic Intervention in Pig and Human Models of Inflammation. <i>Journal of Immunology</i> , 2013, 191, 4769-4777.	0.8	34
45	Maternofetal transplacental transport of recombinant IgG antibodies lacking effector functions. <i>Blood</i> , 2013, 122, 1174-1181.	1.4	43
46	CD40/APC-specific antibodies with three T-cell epitopes loaded in the constant domains induce CD4+ T-cell responses. <i>Protein Engineering, Design and Selection</i> , 2012, 25, 89-96.	2.1	3
47	DeltaPhage™ a novel helper phage for high-valence pIX phagemid display. <i>Nucleic Acids Research</i> , 2012, 40, e120-e120.	14.5	13
48	Anti-carcinoembryonic Antigen Single-chain Variable Fragment Antibody Variants Bind Mouse and Human Neonatal Fc Receptor with Different Affinities That Reveal Distinct Cross-species Differences in Serum Half-life. <i>Journal of Biological Chemistry</i> , 2012, 287, 22927-22937.	3.4	30
49	Targeted DNA vaccines for enhanced induction of idio-type-specific B and T cells. <i>Frontiers in Oncology</i> , 2012, 2, 154.	2.8	23
50	Next generation phage display by use of pVII and pIX as display scaffolds. <i>Methods</i> , 2012, 58, 40-46.	3.8	28
51	Structure-based mutagenesis reveals the albumin-binding site of the neonatal Fc receptor. <i>Nature Communications</i> , 2012, 3, 610.	12.8	160
52	Competition for FcRn-mediated transport gives rise to short half-life of human IgG3 and offers therapeutic potential. <i>Nature Communications</i> , 2011, 2, 599.	12.8	220
53	Posttranslational Modification of Gluten Shapes TCR Usage in Celiac Disease. <i>Journal of Immunology</i> , 2011, 187, 3064-3071.	0.8	92
54	Neonatal Fc receptor for IgG (FcRn) regulates cross-presentation of IgG immune complexes by CD8^αCD11b⁺ dendritic cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 9927-9932.	7.1	187

#	ARTICLE	IF	CITATIONS
55	Polymeric human Fc-fusion proteins with modified effector functions. <i>Scientific Reports</i> , 2011, 1, 124.	3.3	68
56	Expanding the Versatility of Phage Display I: Efficient Display of Peptide-Tags on Protein VII of the Filamentous Phage. <i>PLoS ONE</i> , 2011, 6, e14702.	2.5	20
57	Expanding the Versatility of Phage Display II: Improved Affinity Selection of Folded Domains on Protein VII and IX of the Filamentous Phage. <i>PLoS ONE</i> , 2011, 6, e17433.	2.5	30
58	FcRn binding properties of an abnormal truncated analbuminemic albumin variant. <i>Clinical Biochemistry</i> , 2010, 43, 367-372.	1.9	29
59	Stabilizing mutations increase secretion of functional soluble TCR-Ig fusion proteins. <i>BMC Biotechnology</i> , 2010, 10, 61.	3.3	8
60	Periplasmic expression of soluble single chain T cell receptors is rescued by the chaperone FkpA. <i>BMC Biotechnology</i> , 2010, 10, 8.	3.3	22
61	Aglycosylated IgG variants expressed in bacteria that selectively bind Fc γ RI potentiate tumor cell killing by monocyte-dendritic cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 604-609.	7.1	146
62	Cross-species Binding Analyses of Mouse and Human Neonatal Fc Receptor Show Dramatic Differences in Immunoglobulin G and Albumin Binding. <i>Journal of Biological Chemistry</i> , 2010, 285, 4826-4836.	3.4	165
63	Engineering of the Fc Region for Improved PK (FcRn Interaction). , 2010, , 411-430.		2
64	The Versatile MHC Class I-related FcRn Protects IgG and Albumin from Degradation: Implications for Development of New Diagnostics and Therapeutics. <i>Drug Metabolism and Pharmacokinetics</i> , 2009, 24, 318-332.	2.2	107
65	Identification of a High Affinity Fc γ RIIA-binding Peptide That Distinguishes Fc γ RIIA from Fc γ RIIB and Exploits Fc γ RIIA-mediated Phagocytosis and Degradation. <i>Journal of Biological Chemistry</i> , 2009, 284, 1126-1135.	3.4	8
66	Structural requirements for the interaction of human IgM and IgA with the human Fc γ 1/4 receptor. <i>European Journal of Immunology</i> , 2009, 39, 1147-1156.	2.9	63
67	Ligand binding and antigenic properties of a human neonatal Fc receptor with mutation of two unpaired cysteine residues. <i>FEBS Journal</i> , 2008, 275, 4097-4110.	4.7	30
68	In vitro assessment of recombinant, mutant immunoglobulin ϵ G anti ϵ D devoid of hemolytic activity for treatment of ongoing hemolytic disease of the fetus and newborn. <i>Transfusion</i> , 2008, 48, 12-19.	1.6	13
69	A strategy for bacterial production of a soluble functional human neonatal Fc receptor. <i>Journal of Immunological Methods</i> , 2008, 331, 39-49.	1.4	28
70	Identification of Residues in the C γ 1/4 Domain of Polymeric IgM Essential for Interaction with <i>Plasmodium falciparum</i> Erythrocyte Membrane Protein 1 (PfEMP1). <i>Journal of Immunology</i> , 2008, 181, 1988-2000.	0.8	55
71	Processing of an Antigenic Sequence from IgG Constant Domains for Presentation by MHC Class II. <i>Journal of Immunology</i> , 2008, 181, 7062-7072.	0.8	6
72	Recombinant antibodies for delivery of antigen: a single loop between \hat{A} -strands in the constant region can accommodate long, complex and tandem T cell epitopes. <i>International Immunology</i> , 2008, 20, 295-306.	4.0	4

#	ARTICLE	IF	CITATIONS
73	Reliable titration of filamentous bacteriophages independent of pIII fusion moiety and genome size by using trypsin to restore wild-type pIII phenotype. <i>BioTechniques</i> , 2008, 44, 551-554.	1.8	16
74	A Receptor-Mediated Mechanism to Support Clinical Observation of Altered Albumin Variants. <i>Clinical Chemistry</i> , 2007, 53, 2216-2216.	3.2	14
75	Solution Conformation of Wild-Type and Mutant IgG3 and IgG4 Immunoglobulins Using Crystallography: Possible Implications for Complement Activation. <i>Biophysical Journal</i> , 2007, 93, 3733-3744.	0.5	59
76	Functional in vitro studies of recombinant human immunoglobulin G and immunoglobulin A anti-D. <i>Transfusion</i> , 2007, 47, 306-315.	1.6	14
77	In vitro functional test of two subclasses of an anti-RhD antibody produced by transient expression in COS cells. <i>Apmis</i> , 2006, 114, 345-351.	2.0	5
78	A mutant human IgG molecule with only one C1q binding site can activate complement and induce lysis of target cells. <i>European Journal of Immunology</i> , 2006, 36, 129-138.	2.9	11
79	The conserved histidine 166 residue of the human neonatal Fc receptor heavy chain is critical for the pH-dependent binding to albumin. <i>European Journal of Immunology</i> , 2006, 36, 3044-3051.	2.9	108
80	Identification of a Polymeric Ig Receptor Binding Phage-displayed Peptide That Exploits Epithelial Transcytosis without Dimeric IgA Competition. <i>Journal of Biological Chemistry</i> , 2006, 281, 7075-7081.	3.4	14
81	DNA Vaccines Increase Immunogenicity of Idiotypic Tumor Antigen by Targeting Novel Fusion Proteins to Antigen-Presenting Cells. <i>Molecular Therapy</i> , 2006, 13, 776-785.	8.2	68
82	Prolonged and increased expression of soluble Fc receptors, IgG and a TCR-Ig fusion protein by transiently transfected adherent 293E cells. <i>Journal of Immunological Methods</i> , 2005, 298, 93-104.	1.4	76
83	Induction of central T cell tolerance: Recombinant antibodies deliver peptides for deletion of antigen-specific CD4+8+ thymocytes. <i>European Journal of Immunology</i> , 2005, 35, 3142-3152.	2.9	7
84	Human CD14 is an efficient target for recombinant immunoglobulin vaccine constructs that deliver T cell epitopes. <i>Journal of Leukocyte Biology</i> , 2005, 77, 303-310.	3.3	6
85	Differential Segmental Flexibility and Reach Dictate the Antigen Binding Mode of Chimeric IgD and IgM: Implications for the Function of the B Cell Receptor. <i>Journal of Immunology</i> , 2004, 172, 2925-2934.	0.8	45
86	Monoclonal Antibodies Produced by Muscle after Plasmid Injection and Electroporation. <i>Molecular Therapy</i> , 2004, 9, 328-336.	8.2	63
87	Therapeutic antibodies for human diseases at the dawn of the twenty-first century. <i>Nature Reviews Drug Discovery</i> , 2003, 2, 52-62.	46.4	468
88	The Carboxyl-terminal Domains of IgA and IgM Direct Isotype-specific Polymerization and Interaction with the Polymeric Immunoglobulin Receptor. <i>Journal of Biological Chemistry</i> , 2002, 277, 42755-42762.	3.4	58
89	Efficient Delivery of T Cell Epitopes to APC by Use of MHC Class II-Specific Trophoblasts. <i>Journal of Immunology</i> , 2002, 168, 2154-2162.	0.8	38
90	Antibody-mediated neutralization of cytomegalovirus: modulation of efficacy induced through the IgG constant region. <i>Molecular Immunology</i> , 2002, 38, 833-840.	2.2	13

#	ARTICLE	IF	CITATIONS
91	Balanced expression of single subunits in a multisubunit protein, achieved by cell fusion of individual transfectants. <i>FEBS Journal</i> , 2002, 269, 3205-3210.	0.2	0
92	â€˜Troy-bodiesâ€™™: antibodies as vector proteins for T cell epitopes. <i>New Biotechnology</i> , 2001, 18, 109-116.	2.7	8
93	Recombinant chimeric OKT3 scFv IgM antibodies mediate immune suppression while reducing T cell activation in vitro. <i>European Journal of Immunology</i> , 2001, 31, 94-106.	2.9	16
94	â€œTroy-bodiesâ€ Recombinant Antibodies that Target T Cell Epitopes to Antigen Presenting Cells. <i>International Reviews of Immunology</i> , 2001, 20, 647-673.	3.3	3
95	T Cell Recognition of the Dominant I-Akâ€™Restricted Hen Egg Lysozyme Epitope. <i>Journal of Experimental Medicine</i> , 2001, 193, 1239-1246.	8.5	37
96	Recombinant antibodies as carrier proteins for sub-unit vaccines: influence of mode of fusion on protein production and T-cell activation. <i>Journal of Immunological Methods</i> , 2000, 245, 119-131.	1.4	20
97	Structural requirements for incorporation of J chain into human IgM and IgA. <i>International Immunology</i> , 2000, 12, 19-27.	4.0	56
98	Lysine 322 in the human IgG3 CH2 domain is crucial for antibody dependent complement activation. <i>Molecular Immunology</i> , 2000, 37, 995-1004.	2.2	85
99	Antibodies engineered with IgD specificity efficiently deliver integrated T-cell epitopes for antigen presentation by B cells. <i>Nature Biotechnology</i> , 1999, 17, 670-675.	17.5	51
100	Complement-mediated lysis of cultured osteosarcoma cell lines using chimeric mouse/human TP-1 IgG1 and IgG3 antibodies. <i>Cancer Immunology, Immunotherapy</i> , 1999, 48, 411-418.	4.2	6
101	Recombinant expression of polymeric IgA: incorporation of J chain and secretory component of human origin. <i>European Journal of Immunology</i> , 1999, 29, 1701-1708.	2.9	41
102	Recombinant expression of polymeric IgA: incorporation of J chain and secretory component of human origin. , 1999, 29, 1701.		1
103	The influence of the hinge region length in binding of human IgG to human FcÎ³ receptors. <i>Human Immunology</i> , 1998, 59, 720-727.	2.4	40
104	IgM secretory tailpiece drives multimerisation of bivalent scFv fragments in eukaryotic cells. <i>Immunotechnology: an International Journal of Immunological Engineering</i> , 1998, 4, 141-153.	2.4	19
105	Immunoglobulin as a vehicle for foreign antigenic peptides immunogenic to T cells. <i>Molecular Immunology</i> , 1997, 34, 1167-1176.	2.2	31
106	Versatile vectors for transient and stable expression of recombinant antibody molecules in mammalian cells. <i>Journal of Immunological Methods</i> , 1997, 204, 77-87.	1.4	121
107	Abundant Tyrosine Residues in the Antigen Binding Site in Anti-Osteosarcoma Monoclonal Antibodies Tp-1 and Tp-3: Application to radiolabeling. <i>Acta OncolÃ³gica</i> , 1996, 35, 297-301.	1.8	16
108	The structural requirements for complement activation by IgG: does it hinge on the hinge?. <i>Trends in Immunology</i> , 1995, 16, 85-90.	7.5	140

#	ARTICLE	IF	CITATIONS
109	Cloning and sequencing of V genes from anti-osteosarcoma monoclonal antibodies TP-1 and TP-3: Location of lysine residues and implications for radiolabeling. <i>Nuclear Medicine and Biology</i> , 1995, 22, 765-771.	0.6	21
110	Human IgG isotype-specific amino acid residues affecting complement-mediated cell lysis and phagocytosis. <i>European Journal of Immunology</i> , 1994, 24, 2542-2547.	2.9	40
111	The extended hinge region of IgG3 is not required for high phagocytic capacity mediated by Fc γ 3 receptors, but the heavy chains must be disulfide bonded. <i>European Journal of Immunology</i> , 1993, 23, 1546-1551.	2.9	15
112	Activation of complement by an IgG molecule without a genetic hinge. <i>Nature</i> , 1993, 363, 628-630.	27.8	39
113	Human IgG3 can adopt the disulfide bond pattern characteristic for IgG1 without resembling it in complement mediated cell lysis. <i>Molecular Immunology</i> , 1993, 30, 1419-1425.	2.2	9
114	Antibody dependent cell-mediated cytotoxicity induced by chimeric mouse-human IgG subclasses and IgG3 antibodies with altered hinge region. <i>Molecular Immunology</i> , 1992, 29, 319-326.	2.2	64
115	Chimeric mouse human IgG3 antibodies with an IgG4-like hinge region induce complement-mediated lysis more efficiently than IgG3 with normal hing. <i>European Journal of Immunology</i> , 1991, 21, 2379-2384.	2.9	31
116	C1q binding to chimeric monoclonal IgG3 antibodies consisting of mouse variable regions and human constant regions with shortened hinge containing 15 to 47 amino acids. <i>European Journal of Immunology</i> , 1989, 19, 1599-1603.	2.9	36
117	Mechanism of caffeine-induced inhibition of DNA synthesis in <i>Escherichia coli</i> . <i>FEBS Letters</i> , 1983, 151, 237-242.	2.8	6
118	Effect of caffeine on nucleotide pools in <i>Escherichia coli</i> . <i>Chemico-Biological Interactions</i> , 1982, 40, 141-148.	4.0	3
119	The effect of caffeine on cell growth and metabolism of thymidine in <i>Escherichia coli</i> . <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1980, 73, 29-41.	1.0	39
120	Mechanism of inhibition of thymidine kinase from <i>Escherichia coli</i> by caffeine. <i>FEBS Letters</i> , 1980, 110, 223-226.	2.8	13
121	Extending Antibody Fragment Half-Lives with Albumin. , 0, , 293-310.		0