

# Richard Van Dyck

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12114889/publications.pdf>

Version: 2024-02-01

123  
papers

12,950  
citations

25034

57  
h-index

24258

110  
g-index

128  
all docs

128  
docs citations

128  
times ranked

11847  
citing authors

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | Prediction of treatment discontinuation and recovery from Borderline Personality Disorder: Results from an RCT comparing Schema Therapy and Transference Focused Psychotherapy. <i>Behaviour Research and Therapy</i> , 2015, 74, 60-71. | 3.1  | 55        |
| 2  | Salivary cortisol levels and the 2-year course of depressive and anxiety disorders. <i>Psychoneuroendocrinology</i> , 2013, 38, 1494-1502.   | 2.7  | 110       |
| 3  | Depressive and anxiety disorders on-the-job: The importance of job characteristics for good work functioning in persons with depressive and anxiety disorders. <i>Psychiatry Research</i> , 2012, 200, 382-388.                          | 3.3  | 42        |
| 4  | Sociodemographic and psychiatric determinants of attrition in the Netherlands Study of Depression and Anxiety (NESDA). <i>Comprehensive Psychiatry</i> , 2012, 53, 63-70.  | 3.1  | 150       |
| 5  | The effectiveness of long-term psychoanalytic psychotherapy – A meta-analysis of randomized controlled trials. <i>Clinical Psychology Review</i> , 2012, 32, 81-92.  | 11.4 | 58        |
| 6  | Two-year course of depressive and anxiety disorders: Results from the Netherlands Study of Depression and Anxiety (NESDA). <i>Journal of Affective Disorders</i> , 2011, 133, 76-85.   | 4.1  | 278       |
| 7  | Salivary Cortisol Levels in Persons With and Without Different Anxiety Disorders. <i>Psychosomatic Medicine</i> , 2010, 72, 340-347.   | 2.0  | 238       |
| 8  | Parental history of depression or anxiety and the cortisol awakening response. <i>British Journal of Psychiatry</i> , 2010, 197, 180-185.  | 2.8  | 75        |
| 9  | Do Automatic Self-Associations Relate to Suicidal Ideation?. <i>Journal of Psychopathology and Behavioral Assessment</i> , 2010, 32, 428-437.  | 1.2  | 30        |
| 10 | Treatment of Depression in Patients from Ethnic Minority Groups in the Netherlands. <i>Transcultural Psychiatry</i> , 2010, 47, 473-490.   | 1.6  | 14        |
| 11 | Stimulated Gene Expression Profiles as a Blood Marker of Major Depressive Disorder. <i>Biological Psychiatry</i> , 2010, 68, 179-186.  | 1.3  | 97        |
| 12 | Longitudinal Evidence for Unfavorable Effects of Antidepressants on Heart Rate Variability. <i>Biological Psychiatry</i> , 2010, 68, 861-868.  | 1.3  | 184       |
| 13 | Major Depressive Disorder and Hypothalamic-Pituitary-Adrenal Axis Activity. <i>Archives of General Psychiatry</i> , 2009, 66, 617.   | 12.3 | 641       |
| 14 | Depression Is Associated With Decreased Blood Pressure, but Antidepressant Use Increases the Risk for Hypertension. <i>Hypertension</i> , 2009, 53, 631-638.   | 2.7  | 257       |
| 15 | Associations between sociodemographic, sampling and health factors and various salivary cortisol indicators in a large sample without psychopathology. <i>Psychoneuroendocrinology</i> , 2009, 34, 1109-1120.                            | 2.7  | 159       |
| 16 | Implementation of outpatient schema therapy for borderline personality disorder: study design. <i>BMC Psychiatry</i> , 2009, 9, 64.  | 2.6  | 21        |
| 17 | Three preparatory studies for promoting implementation of outpatient schema therapy for borderline personality disorder in general mental health care. <i>Behaviour Research and Therapy</i> , 2009, 47, 938-945.                        | 3.1  | 18        |
| 18 | Implementation of outpatient schema therapy for borderline personality disorder with versus without crisis support by the therapist outside office hours: A randomized trial. <i>Behaviour Research and Therapy</i> , 2009, 47, 961-973. | 3.1  | 213       |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 19 | Measuring obsessive-compulsive symptoms: Padua Inventory-Revised vs. Yale-Brown Obsessive Compulsive Scale. <i>Journal of Anxiety Disorders</i> , 2009, 23, 830-835.  | 3.2  | 79        |
| 20 | Association between Anxiety Disorders and Heart Rate Variability in The Netherlands Study of Depression and Anxiety (NESDA). <i>Psychosomatic Medicine</i> , 2009, 71, 508-518.   | 2.0  | 164       |
| 21 | Long-term effectiveness and prediction of treatment outcome in cognitive behavioral therapy and sertraline for late-life anxiety disorders. <i>International Psychogeriatrics</i> , 2009, 21, 1148-1159.                | 1.0  | 35        |
| 22 | The Netherlands Study of Depression and Anxiety (NESDA): rationale, objectives and methods. <i>International Journal of Methods in Psychiatric Research</i> , 2008, 17, 121-140.  | 2.1  | 791       |
| 23 | Comorbid depression, but not comorbid anxiety disorders, predicts poor outcome in anxiety disorders. <i>Depression and Anxiety</i> , 2008, 25, 408-415.   | 4.1  | 61        |
| 24 | Genome-wide association of major depression: description of samples for the GAIN Major Depressive Disorder Study: NTR and NESDA biobank projects. <i>European Journal of Human Genetics</i> , 2008, 16, 335-342.        | 2.8  | 145       |
| 25 | Out-patient psychotherapy for borderline personality disorder: Cost-effectiveness of schema-focused therapy v. transference-focused psychotherapy. <i>British Journal of Psychiatry</i> , 2008, 192, 450-457.           | 2.8  | 142       |
| 26 | Association Between Major Depressive Disorder and Heart Rate Variability in the Netherlands Study of Depression and Anxiety (NESDA). <i>Archives of General Psychiatry</i> , 2008, 65, 1358.                            | 12.3 | 306       |
| 27 | Treatment of Panic Disorder and/or Generalized Anxiety Disorder With A Guided Self-Help Manual. <i>Clinical Case Studies</i> , 2007, 6, 277-291.  | 0.8  | 4         |
| 28 | Which Panic Disorder Patients Benefit from Which Treatment: Cognitive Therapy or Antidepressants?. <i>Psychotherapy and Psychosomatics</i> , 2007, 76, 154-161.   | 8.8  | 25        |
| 29 | Combination Treatment for Acute Depression Is Superior Only when Psychotherapy Is Added to Medication. <i>Psychotherapy and Psychosomatics</i> , 2007, 76, 289-297.   | 8.8  | 52        |
| 30 | Anxiety Disorders: Treatable regardless of the Severity of Comorbid Alcohol Dependence. <i>European Addiction Research</i> , 2007, 13, 109-115.   | 2.4  | 18        |
| 31 | Neural correlates of enhanced working-memory performance in dissociative disorder: a functional MRI study. <i>Psychological Medicine</i> , 2007, 37, 235-245.   | 4.5  | 73        |
| 32 | Negative words enhance recognition in nonclinical high dissociators: An fMRI study. <i>NeuroImage</i> , 2007, 37, 323-334.  | 4.2  | 17        |
| 33 | The therapeutic alliance in schema-focused therapy and transference-focused psychotherapy for borderline personality disorder.. <i>Journal of Consulting and Clinical Psychology</i> , 2007, 75, 104-115.               | 2.0  | 150       |
| 34 | Interpersonal psychotherapy (IPT) for late-life depression in general practice: uptake and satisfaction by patients, therapists and physicians. <i>BMC Family Practice</i> , 2007, 8, 52.                               | 2.9  | 15        |
| 35 | Clinical effectiveness of usual care with or without antidepressant medication for primary care patients with minor or mild-major depression: a randomized equivalence trial. <i>BMC Medicine</i> , 2007, 5, 36.        | 5.5  | 37        |
| 36 | Randomised Controlled Trial of a Collaborative Care Model with Psychiatric Consultation for Persistent Medically Unexplained Symptoms in General Practice. <i>Psychotherapy and Psychosomatics</i> , 2006, 75, 282-289. | 8.8  | 118       |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 37 | Interpersonal Psychotherapy for Elderly Patients in Primary Care. <i>American Journal of Geriatric Psychiatry</i> , 2006, 14, 777-786.   | 1.2  | 84        |
| 38 | A Randomized, Controlled Trial of the Effectiveness of Cognitive Behavioral Therapy and Sertraline versus a Waitlist Control Group for Anxiety Disorders in Older Adults. <i>American Journal of Geriatric Psychiatry</i> , 2006, 14, 255-263. | 1.2  | 80        |
| 39 | Glucocorticoids and Relapse of Major Depression (Dexamethasone/Corticotropin-Releasing Hormone) Tj ETQq1 1 0,784314 rgBT /Over   | 1.3  | 148       |
| 40 | Behavioral and cognitive group treatment for fear of flying: A randomized controlled trial. <i>Journal of Behavior Therapy and Experimental Psychiatry</i> , 2006, 37, 358-371.  | 1.2  | 16        |
| 41 | Administering the MADRS by telephone or face-to-face: a validity study. <i>Annals of General Psychiatry</i> , 2006, 5, 3.  | 2.7  | 39        |
| 42 | Interictal Depression, Anxiety, Personality Traits, and Psychological Dissociation in Patients with Temporal Lobe Epilepsy (TLE) and Extra-TLE. <i>Epilepsia</i> , 2006, 47, 2092-2103.  | 5.1  | 108       |
| 43 | Prediction of treatment response by HPA-axis and glucocorticoid receptor polymorphisms in major depression. <i>Psychoneuroendocrinology</i> , 2006, 31, 1154-1163.   | 2.7  | 103       |
| 44 | Well-being, physical functioning, and use of health services in the elderly with PTSD and subthreshold PTSD. <i>International Journal of Geriatric Psychiatry</i> , 2006, 21, 180-188.   | 2.7  | 88        |
| 45 | COMORBID PHOBIC DISORDERS DO NOT INFLUENCE OUTCOME OF ALCOHOL DEPENDENCE TREATMENT. RESULTS OF A NATURALISTIC FOLLOW-UP STUDY. <i>Alcohol and Alcoholism</i> , 2006, 41, 168-173.  | 1.6  | 26        |
| 46 | Thyrotropin, but not a polymorphism in type II deiodinase, predicts response to paroxetine in major depression. <i>European Journal of Endocrinology</i> , 2006, 154, 819-825.   | 3.7  | 18        |
| 47 | Outpatient Psychotherapy for Borderline Personality Disorder. <i>Archives of General Psychiatry</i> , 2006, 63, 649.   | 12.3 | 1,062     |
| 48 | Familial Clustering of Major Depression and Anxiety Disorders in Australian and Dutch Twins and Siblings. <i>Twin Research and Human Genetics</i> , 2005, 8, 609-615.  | 0.6  | 60        |
| 49 | The Effectiveness of Anxiety Treatment on Alcohol-Dependent Patients with a Comorbid Phobic Disorder: A Randomized Controlled Trial. <i>Alcoholism: Clinical and Experimental Research</i> , 2005, 29, 794-800.                                | 2.4  | 74        |
| 50 | Frontal-Striatal Dysfunction During Planning in Obsessive-Compulsive Disorder. <i>Archives of General Psychiatry</i> , 2005, 62, 301.  | 12.3 | 351       |
| 51 | Neurophysiological correlates of increased verbal working memory in high-dissociative participants: a functional MRI study. <i>Psychological Medicine</i> , 2005, 35, 175-185.   | 4.5  | 55        |
| 52 | Thyroid and adrenal axis in major depression: a controlled study in outpatients. <i>European Journal of Endocrinology</i> , 2005, 152, 185-191.  | 3.7  | 88        |
| 53 | Efficacy of self-help manuals for anxiety disorders in primary care: a systematic review. <i>Family Practice</i> , 2005, 22, 192-196.  | 1.9  | 55        |
| 54 | Treatment of anxiety disorders in primary care practice: a randomised controlled trial. <i>British Journal of General Practice</i> , 2005, 55, 763-9.  | 1.4  | 41        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Triiodothyronine Addition to Paroxetine in the Treatment of Major Depressive Disorder. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004, 89, 6271-6276.   | 3.6 | 51        |
| 56 | A patient-doctor relationship questionnaire (PDRQ-9) in primary care: development and psychometric evaluation. <i>General Hospital Psychiatry</i> , 2004, 26, 115-120.  | 2.4 | 164       |
| 57 | Patients' preferences in the treatment of depressive disorder in primary care. <i>General Hospital Psychiatry</i> , 2004, 26, 184-189.  | 2.4 | 349       |
| 58 | Dissociative style and individual differences in verbal working memory span. <i>Consciousness and Cognition</i> , 2004, 13, 821-828.  | 1.5 | 46        |
| 59 | Trauma and Dissociation in Conversion Disorder and Chronic Pelvic Pain. <i>International Journal of Psychiatry in Medicine</i> , 2004, 34, 305-318.   | 1.8 | 40        |
| 60 | Frontostriatal system in planning complexity: a parametric functional magnetic resonance version of tower of london task. <i>NeuroImage</i> , 2003, 18, 367-374.  | 4.2 | 208       |
| 61 | Attention as a characteristic of nonclinical dissociation: an event-related potential study. <i>NeuroImage</i> , 2003, 19, 376-390.   | 4.2 | 39        |
| 62 | A Randomized Controlled Clinical Trial of a Hypnosis-Based Treatment for Patients with Conversion Disorder, Motor Type. <i>International Journal of Clinical and Experimental Hypnosis</i> , 2003, 51, 29-50.                             | 1.8 | 148       |
| 63 | Personality pathology and cognitive-behavioral treatment of fear of flying. <i>Journal of Behavior Therapy and Experimental Psychiatry</i> , 2003, 34, 171-189.   | 1.2 | 11        |
| 64 | DO COMORBID ANXIETY DISORDERS IN ALCOHOL-DEPENDENT PATIENTS NEED SPECIFIC TREATMENT TO PREVENT RELAPSE?. <i>Alcohol and Alcoholism</i> , 2003, 38, 255-262.   | 1.6 | 31        |
| 65 | Directed forgetting between, but not within, dissociative personality states.. <i>Journal of Abnormal Psychology</i> , 2003, 112, 237-243.  | 1.9 | 65        |
| 66 | Prevalence and Risk Factors of Posttraumatic Stress Disorder in Older Adults. <i>Psychotherapy and Psychosomatics</i> , 2003, 72, 333-342.  | 8.8 | 141       |
| 67 | A Randomised Controlled Clinical Trial on the Additional Effect of Hypnosis in a Comprehensive Treatment Programme for In-Patients with Conversion Disorder of the Motor Type. <i>Psychotherapy and Psychosomatics</i> , 2002, 71, 66-76. | 8.8 | 170       |
| 68 | The Relationship between Dissociative Proneness and Alexithymia. <i>Psychotherapy and Psychosomatics</i> , 2002, 71, 104-111.   | 8.8 | 74        |
| 69 | Locus of Control Orientation in Panic Disorder and the Differential Effects of Treatment. <i>Psychotherapy and Psychosomatics</i> , 2002, 71, 85-89.  | 8.8 | 23        |
| 70 | Relevance of Assessment of Cognitions during Panic Attacks in the Treatment of Panic Disorder. <i>Psychotherapy and Psychosomatics</i> , 2002, 71, 158-161.   | 8.8 | 15        |
| 71 | Multicomponent standardized treatment programs for fear of flying: Description and effectiveness. <i>Cognitive and Behavioral Practice</i> , 2002, 9, 138-149.  | 1.5 | 32        |
| 72 | THE INFLUENCE ON TREATMENT GAIN OF COMORBID AVOIDANT PERSONALITY DISORDER IN PATIENTS WITH SOCIAL PHOBIA. <i>Journal of Nervous and Mental Disease</i> , 2002, 190, 41-43.  | 1.0 | 24        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | Cognitive therapy versus moclobemide in social phobia: a controlled study. <i>Clinical Psychology and Psychotherapy</i> , 2001, 8, 263-273.   | 2.7 | 38        |
| 74 | On becoming depressed or anxious in late life: similar vulnerability factors but different effects of stressful life events. <i>British Journal of Psychiatry</i> , 2001, 179, 426-431. | 2.8 | 95        |
| 75 | The placebo response in social phobia. <i>Journal of Psychopharmacology</i> , 2001, 15, 199-203.  | 4.0 | 22        |
| 76 | Cognitive Therapy by Allocation versus Cognitive Therapy by Preference in the Treatment of Panic Disorder. <i>Psychotherapy and Psychosomatics</i> , 2000, 69, 240-243.                 | 8.8 | 31        |
| 77 | Dissociative Style and Directed Forgetting. <i>Cognitive Therapy and Research</i> , 2000, 24, 279-295.  | 1.9 | 52        |
| 78 | Anxiety and Depression in Later Life: Co-Occurrence and Commuality of Risk Factors. <i>American Journal of Psychiatry</i> , 2000, 157, 89-95.   | 7.2 | 505       |
| 79 | Somatoform Dissociation Discriminates Among Diagnostic Categories Over and Above General Psychopathology. <i>Australian and New Zealand Journal of Psychiatry</i> , 1999, 33, 511-520.  | 2.3 | 65        |
| 80 | Hypnotic Recall: A Positive Criterion in the Differential Diagnosis Between Epileptic and Pseudoepileptic Seizures. <i>Epilepsia</i> , 1999, 40, 485-491.                               | 5.1 | 73        |
| 81 | Hoe effectief zijn "evidence based" behandelingen voor paniekstoornis met agorafobie? "Evidence based" behandelingen voor paniekstoornis met agorafobie. <i>Dth</i> , 1999, 19, 19-28.  | 0.2 | 2         |
| 82 | Construction and psychometric characteristics of two self-report questionnaires for the assessment of fear of flying. <i>Psychological Assessment</i> , 1999, 11, 146-158.              | 1.5 | 65        |
| 83 | Dissociation in Temporal Lobe Epilepsy and Pseudo-Epileptic Seizure Patients. <i>Journal of Nervous and Mental Disease</i> , 1999, 187, 713-720.  | 1.0 | 69        |
| 84 | Paroxetine, Clomipramine, and Cognitive Therapy in the Treatment of Panic Disorder. <i>Journal of Clinical Psychiatry</i> , 1999, 60, 831-838.  | 2.2 | 95        |
| 85 | Degree of somatoform and psychological dissociation in dissociative disorder is correlated with reported trauma. <i>Journal of Traumatic Stress</i> , 1998, 11, 711-730.                | 1.8 | 184       |
| 86 | Three controversies about dissociative identity disorder. <i>Journal of Nervous and Mental Disease</i> , 1998, 5, 13-23.  |     | 24        |
| 87 | Anxiety disorders in later life: a report from the longitudinal aging study Amsterdam. <i>Journal of Nervous and Mental Disease</i> , 1998, 13, 717-726.                                |     | 432       |
| 88 | Somatoforme dissociatie: Bijna vergeten, maar meetbaar aanwezig. <i>Dth</i> , 1998, 18, 43-58.  | 0.2 | 2         |
| 89 | Herinneringen aan traumatische ervaringen. <i>Dth</i> , 1998, 18, 59-76.  | 0.2 | 4         |
| 90 | Predictability, controllability, and fear of symptoms of anxiety in epinephrine-induced panic. <i>Biological Psychiatry</i> , 1998, 44, 1017-1026.                                      | 1.3 | 21        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | Somatoform dissociative symptoms as related to animal defensive reactions to predatory imminence and injury.. Journal of Abnormal Psychology, 1998, 107, 63-73.   | 1.9 | 89        |
| 92  | Psychometric Characteristics of the Somatoform Dissociation Questionnaire: A Replication Study<footref rid="foot01"><sup>1</sup></footref>. Psychotherapy and Psychosomatics, 1998, 67, 17-23.                      | 8.8 | 93        |
| 93  | Follow-Up on the Treatment of Panic Disorder With or Without Agoraphobia. Journal of Nervous and Mental Disease, 1998, 186, 414-419.  | 1.0 | 74        |
| 94  | Cognitive and Behavioral Therapies Alone versus in Combination with Fluvoxamine in the Treatment of Obsessive Compulsive Disorder. Journal of Nervous and Mental Disease, 1998, 186, 492-499.                       | 1.0 | 175       |
| 95  | Does Preference for lype of Treatment Matter?. Behavior Modification, 1997, 21, 172-186.  | 1.6 | 32        |
| 96  | Cardiovascular, respiratory, and panic reactions to epinephrine in panic disorder patients. Biological Psychiatry, 1997, 41, 249-251.   | 1.3 | 4         |
| 97  | People who seek help for fear of flying: Typology of flying phobics. Behavior Therapy, 1997, 28, 237-251.   | 2.4 | 54        |
| 98  | On distinguishing types of panic. Journal of Anxiety Disorders, 1996, 10, 173-184.  | 3.2 | 6         |
| 99  | Psychiatric Consultation for Somatizing Patients in the Family Practice Setting: A Feasibility Study. International Journal of Psychiatry in Medicine, 1996, 26, 223-239.   | 1.8 | 17        |
| 100 | The Case for a Dissociative Interpretation of Pseudoepileptic Seizures. Journal of Nervous and Mental Disease, 1996, 184, 468-474.  | 1.0 | 53        |
| 101 | The Development and Psychometric Characteristics of the Somatoform Dissociation Questionnaire (SDQ-20). Journal of Nervous and Mental Disease, 1996, 184, 688-694.  | 1.0 | 460       |
| 102 | Epinephrine and fear of bodily sensations in panic disorder and social phobia. Journal of Psychopharmacology, 1996, 10, 259-265.  | 4.0 | 13        |
| 103 | Dissociative and Hypnotic Experiences in Eating Disorder Patients: An Exploratory Study. American Journal of Clinical Hypnosis, 1995, 38, 97-108.   | 0.6 | 32        |
| 104 | Respiratory Training Prior to Exposure <i>in vivo</i> in the Treatment of Panic Disorder with Agoraphobia: Efficacy and Predictors of Outcome. Australian and New Zealand Journal of Psychiatry, 1995, 29, 104-113. | 2.3 | 21        |
| 105 | The sensitivity to change of measures for obsessive-compulsive disorder. Journal of Anxiety Disorders, 1995, 9, 241-248.  | 3.2 | 44        |
| 106 | Cognitive therapy and exposure in vivo in the treatment of obsessive compulsive disorder. Behaviour Research and Therapy, 1995, 33, 379-390.  | 3.1 | 313       |
| 107 | Fear of fear, trait anxiety and aerobic fitness in relation to state anxiety during adrenalin provocation. Anxiety, Stress and Coping, 1994, 7, 279-289.  | 2.9 | 1         |
| 108 | Assessing the clinical significance of outcome in agoraphobia research: A comparison of two approaches. Behavior Therapy, 1994, 25, 147-158.  | 2.4 | 12        |

| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 109 | Autogenic training and cognitive self-hypnosis for the treatment of recurrent headaches in three different subject groups. <i>Pain</i> , 1994, 58, 331-340.                                | 4.2  | 65        |
| 110 | Dissociative experiences and trauma in eating disorders. <i>International Journal of Eating Disorders</i> , 1993, 13, 187-193.   | 4.0  | 151       |
| 111 | Goal attainment scaling: An idiosyncratic method to assess treatment effectiveness in agoraphobia. <i>Journal of Psychopathology and Behavioral Assessment</i> , 1993, 15, 357-373.        | 1.2  | 11        |
| 112 | Dissociatieve ervaringen bij Vlamingen en Nederlanders; een studie met de Dissociation Questionnaire (DISQ). <i>Dth</i> , 1993, 13, 27-36.   | 0.2  | 5         |
| 113 | The dissociation questionnaire (DISQ): Development and characteristics of a new self-report questionnaire. <i>Clinical Psychology and Psychotherapy</i> , 1993, 1, 21-27.                  | 2.7  | 179       |
| 114 | Adrenaline and the relationship between neurosomaticism, aerobic fitness and mental task performance. <i>Biological Psychology</i> , 1993, 36, 157-181.                                    | 2.2  | 11        |
| 115 | Dissociation and Traumatic Experiences in the General Population of the Netherlands. <i>Psychiatric Services</i> , 1993, 44, 786-788.  | 2.0  | 11        |
| 116 | Autogenic training and self-hypnosis in the control of tension headache. <i>General Hospital Psychiatry</i> , 1992, 14, 408-415.   | 2.4  | 43        |
| 117 | Breathing retraining: A rational placebo?. <i>Clinical Psychology Review</i> , 1992, 12, 141-153.  | 11.4 | 65        |
| 118 | Self-monitoring of panic attacks and retrospective estimates of panic: discordant findings. <i>Behaviour Research and Therapy</i> , 1992, 30, 411-413.                                     | 3.1  | 54        |
| 119 | Hypnosis and autogenic training in the treatment of tension headaches: A two-phase constructive design study with follow-up. <i>Journal of Psychosomatic Research</i> , 1992, 36, 219-228. | 2.6  | 48        |
| 120 | Hyperventilatieprovocatie en ademhalingsoefeningen voorafgaande aan exposure in vivo bij de behandeling van patiënten met paniekstoornis met agorafobie. <i>Dth</i> , 1992, 12, 113-122.   | 0.2  | 0         |
| 121 | Over het registreren van paniek door agorafobici. <i>Dth</i> , 1992, 12, 123-130.  | 0.2  | 0         |
| 122 | Behavioral assessment of avoidance in agoraphobia. <i>Journal of Psychopathology and Behavioral Assessment</i> , 1991, 13, 285-300.  | 1.2  | 21        |
| 123 | Hypnosis and Conversion Disorders. <i>American Journal of Psychotherapy</i> , 1989, 43, 480-493.   | 1.2  | 11        |