

Robin F Krimm

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12124/publications.pdf>

Version: 2024-02-01

36
papers

980
citations

394421

19
h-index

454955

30
g-index

38
all docs

38
docs citations

38
times ranked

553
citing authors

#	ARTICLE	IF	CITATIONS
1	Whole-Mount Staining, Visualization, and Analysis of Fungiform, Circumvallate, and Palate Taste Buds. <i>Journal of Visualized Experiments</i> , 2021, , .	0.3	3
2	Variable Branching Characteristics of Peripheral Taste Neurons Indicates Differential Convergence. <i>Journal of Neuroscience</i> , 2021, 41, 4850-4866.	3.6	15
3	Variation in taste ganglion neuron morphology: insights into taste function and plasticity. <i>Current Opinion in Physiology</i> , 2021, 20, 134-139.	1.8	2
4	TrkB expression and dependence divides gustatory neurons into three subpopulations. <i>Neural Development</i> , 2019, 14, 3.	2.4	7
5	Maintenance of Mouse Gustatory Terminal Field Organization Is Dependent on BDNF at Adulthood. <i>Journal of Neuroscience</i> , 2018, 38, 6873-6887.	3.6	6
6	BDNF is required for taste axon regeneration following unilateral chorda tympani nerve section. <i>Experimental Neurology</i> , 2017, 293, 27-42.	4.1	13
7	The transcription factor <i>Phox2b</i> distinguishes between oral and non-oral sensory neurons in the geniculate ganglion. <i>Journal of Comparative Neurology</i> , 2017, 525, 3935-3950.	1.6	26
8	Taste bud-derived BDNF maintains innervation of a subset of TrkB-expressing gustatory nerve fibers. <i>Molecular and Cellular Neurosciences</i> , 2017, 82, 195-203.	2.2	14
9	Insulin-Like Growth Factors Are Expressed in the Taste System, but Do Not Maintain Adult Taste Buds. <i>PLoS ONE</i> , 2016, 11, e0148315.	2.5	11
10	Postnatal reduction of BDNF regulates the developmental remodeling of taste bud innervation. <i>Developmental Biology</i> , 2015, 405, 225-236.	2.0	17
11	Taste Bud-Derived BDNF Is Required to Maintain Normal Amounts of Innervation to Adult Taste Buds. <i>ENeuro</i> , 2015, 2, ENEURO.0097-15.2015.	1.9	29
12	The neurotrophin receptor p75 regulates gustatory axon branching and promotes innervation of the tongue during development. <i>Neural Development</i> , 2014, 9, 15.	2.4	13
13	BDNF and NT4 play interchangeable roles in gustatory development. <i>Developmental Biology</i> , 2014, 386, 308-320.	2.0	15
14	Taste Neurons Consist of Both a Large TrkB-Receptor-Dependent and a Small TrkB-Receptor-Independent Subpopulation. <i>PLoS ONE</i> , 2013, 8, e83460.	2.5	13
15	Neurotrophin-4 regulates the survival of gustatory neurons earlier in development using a different mechanism than brain-derived neurotrophic factor. <i>Developmental Biology</i> , 2012, 365, 50-60.	2.0	22
16	Lingual and palatal gustatory afferents each depend on both BDNF and NT4, but the dependence is greater for lingual than palatal afferents. <i>Journal of Comparative Neurology</i> , 2010, 518, 3290-3301.	1.6	25
17	Developmental expression of <i>Bdnf</i> , <i>Ntf4/5</i> , and <i>TrkB</i> in the mouse peripheral taste system. <i>Developmental Dynamics</i> , 2010, 239, 2637-2646.	1.8	34
18	BDNF is required for the survival of differentiated geniculate ganglion neurons. <i>Developmental Biology</i> , 2010, 340, 419-429.	2.0	38

#	ARTICLE	IF	CITATIONS
19	Epithelial-Derived Brain-Derived Neurotrophic Factor Is Required for Gustatory Neuron Targeting during a Critical Developmental Period. <i>Journal of Neuroscience</i> , 2009, 29, 3354-3364.	3.6	42
20	Fate mapping of mammalian embryonic taste bud progenitors. <i>Development (Cambridge)</i> , 2009, 136, 1519-1528.	2.5	83
21	Exuberant Neuronal Convergence onto Reduced Taste Bud Targets with Preservation of Neural Specificity in Mice Overexpressing Neurotrophin in the Tongue Epithelium. <i>Journal of Neuroscience</i> , 2007, 27, 13875-13881.	3.6	9
22	Factors that regulate embryonic gustatory development. <i>BMC Neuroscience</i> , 2007, 8, S4.	1.9	41
23	Epithelial overexpression of BDNF and NT4 produces distinct gustatory axon morphologies that disrupt initial targeting. <i>Developmental Biology</i> , 2006, 292, 457-468.	2.0	39
24	Mice lacking the p75 receptor fail to acquire a normal complement of taste buds and geniculate ganglion neurons by adulthood. <i>The Anatomical Record Part A: Discoveries in Molecular, Cellular, and Evolutionary Biology</i> , 2006, 288A, 1294-1302.	2.0	8
25	Refinement of innervation accuracy following initial targeting of peripheral gustatory fibers. <i>Journal of Neurobiology</i> , 2006, 66, 1033-1043.	3.6	27
26	Overexpression of neurotrophin 4 in skin enhances myelinated sensory endings but does not influence sensory neuron number. <i>Journal of Comparative Neurology</i> , 2006, 498, 455-465.	1.6	29
27	Neuron/target plasticity in the peripheral gustatory system. <i>Journal of Comparative Neurology</i> , 2004, 472, 183-192.	1.6	29
28	NT3 expressed in skin causes enhancement of SA1 sensory neurons that leads to postnatal enhancement of Merkel cells. <i>Journal of Comparative Neurology</i> , 2004, 471, 352-360.	1.6	27
29	Epithelial Overexpression of BDNF or NT4 Disrupts Targeting of Taste Neurons That Innervate the Anterior Tongue. <i>Developmental Biology</i> , 2001, 232, 508-521.	2.0	71
30	Cutaneous overexpression of neurotrophin-3 (NT3) selectively restores sensory innervation in NT3 gene knockout mice. , 2000, 43, 40-49.		20
31	Neuron/target matching between chorda tympani neurons and taste buds during postnatal rat development. <i>Journal of Neurobiology</i> , 2000, 43, 98-106.	3.6	24
32	Overexpression of Brain-Derived Neurotrophic Factor Enhances Sensory Innervation and Selectively Increases Neuron Number. <i>Journal of Neuroscience</i> , 1999, 19, 5919-5931.	3.6	100
33	Early dietary sodium restriction disrupts the peripheral anatomical development of the gustatory system. , 1999, 39, 218-226.		17
34	Quantitative Relationships between Taste Bud Development and Gustatory Ganglion Cells. <i>Annals of the New York Academy of Sciences</i> , 1998, 855, 70-75.	3.8	10
35	Innervation of single fungiform taste buds during development in rat. <i>Journal of Comparative Neurology</i> , 1998, 398, 13-24.	1.6	56
36	Early prenatal critical period for chorda tympani nerve terminal field development. <i>Journal of Comparative Neurology</i> , 1997, 378, 254-264.	1.6	35