

# Verena Labi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1215306/publications.pdf>

Version: 2024-02-01

50  
papers

2,759  
citations

236925

25  
h-index

243625

44  
g-index

52  
all docs

52  
docs citations

52  
times ranked

4636  
citing authors

#	ARTICLE	IF	CITATIONS
1	BH3-only proteins Puma and Bim are rate-limiting for $\hat{I}^3$ -radiation-induced apoptosis of lymphoid cells in vivo. <i>Blood</i> , 2005, 106, 4131-4138.	1.4	259
2	Genome-wide association analysis in primary sclerosing cholangitis identifies two non-HLA susceptibility loci. <i>Nature Genetics</i> , 2011, 43, 17-19.	21.4	221
3	Puma cooperates with Bim, the rate-limiting BH3-only protein in cell death during lymphocyte development, in apoptosis induction. <i>Journal of Experimental Medicine</i> , 2006, 203, 2939-2951.	8.5	209
4	How cell death shapes cancer. <i>Cell Death and Disease</i> , 2015, 6, e1675-e1675.	6.3	205
5	BH3-only proteins in cell death initiation, malignant disease and anticancer therapy. <i>Cell Death and Differentiation</i> , 2006, 13, 1325-1338.	11.2	159
6	Targeting the Bcl-2-regulated apoptosis pathway by BH3 mimetics: a breakthrough in anticancer therapy?. <i>Cell Death and Differentiation</i> , 2008, 15, 977-987.	11.2	147
7	Caspase-2 activation in the absence of PIDDosome formation. <i>Journal of Cell Biology</i> , 2009, 185, 291-303.	5.2	144
8	Loss of the BH3-only protein Bmf impairs B cell homeostasis and accelerates $\hat{I}^3$ irradiation-induced thymic lymphoma development. <i>Journal of Experimental Medicine</i> , 2008, 205, 641-655.	8.5	116
9	Bim and Bmf in tissue homeostasis and malignant disease. <i>Oncogene</i> , 2008, 27, S41-S52.	5.9	109
10	AICAR induces apoptosis independently of AMPK and p53 through up-regulation of the BH3-only proteins BIM and NOXA in chronic lymphocytic leukemia cells. <i>Blood</i> , 2010, 116, 3023-3032.	1.4	95
11	Apoptosis of leukocytes triggered by acute DNA damage promotes lymphoma formation. <i>Genes and Development</i> , 2010, 24, 1602-1607.	5.9	95
12	The Nuclear Orphan Receptor NR2F6 Suppresses Lymphocyte Activation and T Helper 17-Dependent Autoimmunity. <i>Immunity</i> , 2008, 29, 205-216.	14.3	93
13	Tissue-specific DNA demethylation is required for proper B-cell differentiation and function. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 5018-5023.	7.1	83
14	The NF- $\hat{A}$ B regulator Bcl-3 and the BH3-only proteins Bim and Puma control the death of activated T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 10979-10984.	7.1	80
15	PIDDosome-independent tumor suppression by Caspase-2. <i>Cell Death and Differentiation</i> , 2012, 19, 1722-1732.	11.2	60
16	Suppression of B-cell lymphomagenesis by the BH3-only proteins Bmf and Bad. <i>Blood</i> , 2010, 115, 995-1005.	1.4	53
17	The Bcl-2 protein family and its role in the development of neoplastic disease. <i>Experimental Gerontology</i> , 2004, 39, 1125-1135.	2.8	49
18	Chemokine-mediated redirection of T cells constitutes a critical mechanism of glucocorticoid therapy in autoimmune CNS responses. <i>Acta Neuropathologica</i> , 2014, 127, 713-729.	7.7	46

#	ARTICLE	IF	CITATIONS
19	A c-Myc/miR17-92/Pten Axis Controls PI3K-Mediated Positive and Negative Selection in B Cell Development and Reconstitutes CD19 Deficiency. <i>Cell Reports</i> , 2016, 16, 419-431.	6.4	45
20	SAFB2 Enables the Processing of Suboptimal Stem-Loop Structures in Clustered Primary miRNA Transcripts. <i>Molecular Cell</i> , 2020, 78, 876-889.e6.	9.7	43
21	Deregulated cell death and lymphocyte homeostasis cause premature lethality in mice lacking the BH3-only proteins Bim and Bmf. <i>Blood</i> , 2014, 123, 2652-2662.	1.4	40
22	<scp>TET</scp> enzymes control antibody production and shape the mutational landscape in germinal centre B cells. <i>FEBS Journal</i> , 2019, 286, 3566-3581.	4.7	37
23	BH3-only protein Bim more critical than Puma in tyrosine kinase inhibitorâ€‘induced apoptosis of human leukemic cells and transduced hematopoietic progenitors carrying oncogenic FLT3. <i>Blood</i> , 2009, 113, 2302-2311.	1.4	31
24	Can the analysis of BH3-only protein knockout mice clarify the issue of â€‘direct versus indirectâ€™™ activation of Bax and Bak?. <i>Cell Death and Differentiation</i> , 2011, 18, 1545-1546.	11.2	30
25	Checkpoint kinase 1 is essential for normal B cell development and lymphomagenesis. <i>Nature Communications</i> , 2017, 8, 1697.	12.8	28
26	Pro-apoptotic Bax is the major and Bak an auxiliary effector in cytokine deprivation-induced mast cell apoptosis. <i>Cell Death and Disease</i> , 2010, 1, e43-e43.	6.3	26
27	Haematopoietic stem cell survival and transplantation efficacy is limited by the BH3â€‘only proteins Bim and Bmf. <i>EMBO Molecular Medicine</i> , 2013, 5, 122-136.	6.9	25
28	The c-Myc/miR17-92/PTEN Axis Tunes PI3K Activity to Control Expression of Recombination Activating Genes in Early B Cell Development. <i>Frontiers in Immunology</i> , 2018, 9, 2715.	4.8	24
29	Transient apoptosis inhibition in donor stem cells improves hematopoietic stem cell transplantation. <i>Journal of Experimental Medicine</i> , 2017, 214, 2967-2983.	8.5	21
30	Loss of the pro-apoptotic BH3-only Bcl-2 family member Bim sustains B lymphopoiesis in the absence of IL-7. <i>International Immunology</i> , 2009, 21, 715-725.	4.0	20
31	Canonical NF-Î²B signaling is uniquely required for the long-term persistence of functional mature B cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 5065-5070.	7.1	20
32	Orphan Nuclear Receptor NR2F6 Suppresses T Follicular Helper Cell Accumulation through Regulation of IL-21. <i>Cell Reports</i> , 2019, 28, 2878-2891.e5.	6.4	20
33	Combined loss of the BH3-only proteins Bim and Bmf restores B-cell development and function in TACI-Ig transgenic mice. <i>Cell Death and Differentiation</i> , 2015, 22, 1477-1488.	11.2	18
34	PUMA-mediated tumor suppression: A tale of two stories. <i>Cell Cycle</i> , 2010, 9, 4269-4275.	2.6	16
35	Minor cell-death defects but reduced tumor latency in mice lacking the BH3-only proteins Bad and Bmf. <i>Oncogene</i> , 2013, 32, 621-630.	5.9	15
36	CHK1 dosage in germinal center B cells controls humoral immunity. <i>Cell Death and Differentiation</i> , 2019, 26, 2551-2567.	11.2	14

#	ARTICLE	IF	CITATIONS
37	Context-specific regulation of cell survival by a miRNA-controlled BIM rheostat. <i>Genes and Development</i> , 2019, 33, 1673-1687.	5.9	13
38	BH3-only protein Noxa contributes to apoptotic control of stress-erythropoiesis. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2013, 18, 1306-1318.	4.9	10
39	Loss of the orphan nuclear receptor NR2F6 enhances CD8+ T-cell memory via IFN- $\gamma$ . <i>Cell Death and Disease</i> , 2021, 12, 187.	6.3	10
40	miR-17-1/492 in lymphocyte development and lymphomagenesis. <i>Cancer Letters</i> , 2019, 446, 73-80.	7.2	8
41	Cell signaling and the aging of B cells. <i>Experimental Gerontology</i> , 2020, 138, 110985.	2.8	6
42	Differential effects of Vav $\beta$ -promoter $\beta$ -driven overexpression of BCLX and BFL1 on lymphocyte survival and B cell lymphomagenesis. <i>FEBS Journal</i> , 2018, 285, 1403-1418.	4.7	5
43	Deletion of Puma and p21Waf1 In Mice Deactivates p53-Induced Cell Death and Cell Cycle Arrest, but Protects Mice From Irradiation-Induced Lymphomagenesis by a Mechanism Involving Hemopoietic Stem Cell Quiescence. <i>Blood</i> , 2010, 116, 90-90.	1.4	5
44	MCL-1 and BCL-XL: blood brothers. <i>Blood</i> , 2021, 137, 1850-1851.	1.4	2
45	Role for BH3-Only Protein NOXA In Growth-Factor Deprivation and Early Erythropoiesis. <i>Blood</i> , 2010, 116, 4235-4235.	1.4	0
46	Lack of the BH3-Only Proteins Bim, Bmf and Puma In Haematopoietic Stem and Progenitor Cells Facilitates Early Reconstitution and Long Term Haematopoiesis.. <i>Blood</i> , 2010, 116, 1542-1542.	1.4	0
47	Deletion of the p53 Target Gene PUMA Prevents Bone Marrow Failure in a Dyskeratosis Congenita Mouse Model. <i>Blood</i> , 2018, 132, 648-648.	1.4	0
48	Orphan Nuclear Receptor NR2F6 Suppresses T Follicular Helper Cell Accumulation Through Direct Regulation of IL-21. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
49	Addiction of B Cell Lymphomas to microRNA-17-92 Is Mediated By a Single Target Gene. <i>Blood</i> , 2019, 134, 1258-1258.	1.4	0
50	Bacterial Infection with <i>Listeria monocytogenes</i> in Mice and Subsequent Analysis of AntigenSpecific CD8 T Cell Responses. <i>Bio-protocol</i> , 2021, 11, e4247.	0.4	0