

Andres Saag

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12170644/publications.pdf>

Version: 2024-02-01

19

papers

403

citations

933447

10

h-index

752698

20

g-index

20

all docs

20

docs citations

20

times ranked

579

citing authors

#	ARTICLE	IF	CITATIONS
1	Evolution of complex symbiotic relationships in a morphologically derived family of lichen-forming fungi. <i>New Phytologist</i> , 2015, 208, 1217-1226.	7.3	105
2	World survey of the genus <i>Lepraria</i> (<i>Stereocaulaceae</i> , lichenized Ascomycota). <i>Lichenologist</i> , 2009, 41, 25-60.	0.8	65
3	Phylogeny of the cetrarioid core (Parmeliaceae) based on five genetic markers. <i>Lichenologist</i> , 2009, 41, 489-511.	0.8	43
4	Evaluation of traditionally circumscribed species in the lichen-forming genus <i>Usnea</i> , section <i>Usnea</i> (Parmeliaceae, Ascomycota) using a six-locus dataset. <i>Organisms Diversity and Evolution</i> , 2016, 16, 497-524.	1.6	32
5	Species delimitation in the lichenized fungal genus <i>Vulpicida</i> (Parmeliaceae, Ascomycota) using gene concatenation and coalescent-based species tree approaches. <i>American Journal of Botany</i> , 2014, 101, 2169-2182.	1.7	19
6	Diversity of lichens and bryophytes in hybrid aspen plantations in Estonia depends on landscape structure. <i>Canadian Journal of Forest Research</i> , 2017, 47, 1202-1214.	1.7	19
7	Phylogenetic relations of European shrubby taxa of the genus <i>Usnea</i> . <i>Lichenologist</i> , 2011, 43, 427-444.	0.8	18
8	A Second Updated World List of Cetrarioid Lichens. <i>Bryologist</i> , 1997, 100, 109.	0.6	17
9	Forest biomass, soil and biodiversity relationships originate from biogeographic affinity and direct ecological effects. <i>Oikos</i> , 2019, 128, 1653-1665.	2.7	16
10	A new circumscription of the lichen genus <i>Nephromopsis</i> (Parmeliaceae, lichenized Ascomycetes). <i>Mycological Progress</i> , 2005, 4, 303-316.	1.4	14
11	Testing morphology-based delimitation of <i>Vulpicida juniperinus</i> and <i>V. tubulosus</i> (Parmeliaceae) using three molecular markers. <i>Lichenologist</i> , 2012, 44, 757-772.	0.8	10
12	Impact of alkaline dust pollution on genetic variation of <i>Usnea subfloridana</i> populations. <i>Fungal Biology</i> , 2016, 120, 1165-1174.	2.5	10
13	Lichen chemistry is concordant with multilocus gene genealogy in the genus <i>Cetrelia</i> (Parmeliaceae). <i>Tj ETQql 1 0.784314 rgBT /Overlo</i>	2.5	10
14	The effect of stand age on biodiversity in a 130-year chronosequence of <i>Populus tremula</i> stands. <i>Forest Ecology and Management</i> , 2022, 504, 119833.	3.2	7
15	A Revision of the North American Lichen Genus <i>Ahtiana</i> (Parmeliaceae). <i>Bryologist</i> , 1995, 98, 596.	0.6	5
16	Unconstrained gene flow between populations of a widespread epiphytic lichen <i>Usnea subfloridana</i> (Parmeliaceae, Ascomycota) in Estonia. <i>Fungal Biology</i> , 2018, 122, 731-737.	2.5	5
17	Third world list of cetrarioid lichens: A databased tool for documentation of nomenclatural dataâ€”lessons learned. <i>Taxon</i> , 2013, 62, 591-603.	0.7	4
18	Microsatellite based genetic diversity of the widespread epiphytic lichen <i>Usnea subfloridana</i> (Parmeliaceae, Ascomycota) in Estonia: comparison of populations from the mainland and an island. <i>MycoKeys</i> , 2019, 58, 27-45.	1.9	2

#	ARTICLE	IF	CITATIONS
19	Seventy-year history of management using low-intensity harvesting methods: weak impact on biodiversity of hemiboreal Scots pine forests. Canadian Journal of Forest Research, 2020, 50, 1268-1280.	1.7	1