

# Andres Saag

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12170644/publications.pdf>

Version: 2024-02-01

19

papers

403

citations

933447

10

h-index

752698

20

g-index

20

all docs

20

docs citations

20

times ranked

579

citing authors

#	ARTICLE	IF	CITATIONS
1	The effect of stand age on biodiversity in a 130-year chronosequence of <i>Populus tremula</i> stands. <i>Forest Ecology and Management</i> , 2022, 504, 119833.	3.2	7
2	Seventy-year history of management using low-intensity harvesting methods: weak impact on biodiversity of hemiboreal Scots pine forests. <i>Canadian Journal of Forest Research</i> , 2020, 50, 1268-1280.	1.7	1
3	Forest biomass, soil and biodiversity relationships originate from biogeographic affinity and direct ecological effects. <i>Oikos</i> , 2019, 128, 1653-1665.	2.7	16
4	Lichen chemistry is concordant with multilocus gene genealogy in the genus <i>Cetrelia</i> (Parmeliaceae). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 2.5</i>	0.6	10
5	Microsatellite based genetic diversity of the widespread epiphytic lichen <i>Usnea subfloridana</i> (Parmeliaceae, Ascomycota) in Estonia: comparison of populations from the mainland and an island. <i>MycoKeys</i> , 2019, 58, 27-45.	1.9	2
6	Unconstrained gene flow between populations of a widespread epiphytic lichen <i>Usnea subfloridana</i> (Parmeliaceae, Ascomycota) in Estonia. <i>Fungal Biology</i> , 2018, 122, 731-737.	2.5	5
7	Diversity of lichens and bryophytes in hybrid aspen plantations in Estonia depends on landscape structure. <i>Canadian Journal of Forest Research</i> , 2017, 47, 1202-1214.	1.7	19
8	Impact of alkaline dust pollution on genetic variation of <i>Usnea subfloridana</i> populations. <i>Fungal Biology</i> , 2016, 120, 1165-1174.	2.5	10
9	Evaluation of traditionally circumscribed species in the lichen-forming genus <i>Usnea</i> , section <i>Usnea</i> (Parmeliaceae, Ascomycota) using a six-locus dataset. <i>Organisms Diversity and Evolution</i> , 2016, 16, 497-524.	1.6	32
10	Evolution of complex symbiotic relationships in a morphologically derived family of lichen-forming fungi. <i>New Phytologist</i> , 2015, 208, 1217-1226.	7.3	105
11	Species delimitation in the lichenized fungal genus <i>Vulpicida</i> (Parmeliaceae, Ascomycota) using gene concatenation and coalescent-based species tree approaches. <i>American Journal of Botany</i> , 2014, 101, 2169-2182.	1.7	19
12	Third world list of cetrarioid lichens: A databased tool for documentation of nomenclatural data—lessons learned. <i>Taxon</i> , 2013, 62, 591-603.	0.7	4
13	Testing morphology-based delimitation of <i>Vulpicida juniperinus</i> and <i>V. tubulosus</i> (Parmeliaceae) using three molecular markers. <i>Lichenologist</i> , 2012, 44, 757-772.	0.8	10
14	Phylogenetic relations of European shrubby taxa of the genus <i>Usnea</i> . <i>Lichenologist</i> , 2011, 43, 427-444.	0.8	18
15	World survey of the genus <i>Lepraria</i> ( <i>Stereocaulaceae</i> , lichenized Ascomycota). <i>Lichenologist</i> , 2009, 41, 25-60.	0.8	65
16	Phylogeny of the cetrarioid core (Parmeliaceae) based on five genetic markers. <i>Lichenologist</i> , 2009, 41, 489-511.	0.8	43
17	A new circumscription of the lichen genus <i>Nephromopsis</i> (Parmeliaceae, lichenized Ascomycetes). <i>Mycological Progress</i> , 2005, 4, 303-316.	1.4	14
18	A Second Updated World List of Cetrarioid Lichens. <i>Bryologist</i> , 1997, 100, 109.	0.6	17

#	ARTICLE	IF	CITATIONS
19	A Revision of the North American Lichen Genus Ahtiana (Parmeliaceae). <i>Bryologist</i> , 1995, 98, 596.	0.6	5