## Anne E M Speckens

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/127392/publications.pdf

Version: 2024-02-01

50276 37204 10,481 175 46 96 citations h-index g-index papers 189 189 189 10876 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	A validation study of the Hospital Anxiety and Depression Scale (HADS) in different groups of Dutch subjects. Psychological Medicine, 1997, 27, 363-370.	4.5	1,723
2	Distinct Neural Correlates of Washing, Checking, and Hoarding SymptomDimensions in Obsessive-compulsive Disorder. Archives of General Psychiatry, 2004, 61, 564.	12.3	811
3	Efficacy of Mindfulness-Based Cognitive Therapy in Prevention of Depressive Relapse. JAMA Psychiatry, 2016, 73, 565.	11.0	558
4	Characteristics and content of intrusive memories in PTSD and their changes with treatment. Journal of Traumatic Stress, 2004, 17, 231-240.	1.8	316
5	Cognitive behavioural therapy for medically unexplained physical symptoms: a randomised controlled trial. BMJ: British Medical Journal, 1995, 311, 1328-1332.	2.3	283
6	A validation study of the Whitely Index, the Illness Attitude Scales, and the Somatosensory Amplification Scale in general medical and general practice patients. Journal of Psychosomatic Research, 1996, 40, 95-104.	2.6	266
7	Decision making and set shifting impairments are associated with distinct symptom dimensions in obsessive-compulsive disorder Neuropsychology, 2006, 20, 409-419.	1.3	242
8	Medically unexplained symptoms, somatisation disorder and hypochondriasis: Course and prognosis. A systematic review. Journal of Psychosomatic Research, 2009, 66, 363-377.	2.6	214
9	The efficacy of mindfulness-based cognitive therapy in recurrent depressed patients with and without a current depressive episode: a randomized controlled trial. Psychological Medicine, 2012, 42, 989-1001.	4.5	211
10	Greater efficiency in attentional processing related to mindfulness meditation. Quarterly Journal of Experimental Psychology, 2010, 63, 1168-1180.	1.1	205
11	Mindfulnessâ€based interventions for psychological and physical health outcomes in cancer patients and survivors: A systematic review and metaâ€analysis of randomized controlled trials. Psycho-Oncology, 2019, 28, 2257-2269.	2.3	178
12	Detecting Somatoform Disorders in Primary Care With the PHQ-15. Annals of Family Medicine, 2009, 7, 232-238.	1.9	165
13	Face-to-Face and Internet-Based Mindfulness-Based Cognitive Therapy Compared With Treatment as Usual in Reducing Psychological Distress in Patients With Cancer: A Multicenter Randomized Controlled Trial. Journal of Clinical Oncology, 2018, 36, 2413-2421.	1.6	151
14	Efficacy of Blended Cognitive Behavior Therapy for High Fear of Recurrence in Breast, Prostate, and Colorectal Cancer Survivors: The SWORD Study, a Randomized Controlled Trial. Journal of Clinical Oncology, 2017, 35, 2173-2183.	1.6	142
15	To discard or not to discard: the neural basis of hoarding symptoms in obsessive-compulsive disorder. Molecular Psychiatry, 2009, 14, 318-331.	7.9	137
16	Neural correlates of anxiety associated with obsessive-compulsive symptom dimensions in normal volunteers. Biological Psychiatry, 2003, 53, 482-493.	1.3	136
17	Effects of mindfulness-based cognitive therapy on neurophysiological correlates of performance monitoring in adult attention-deficit/hyperactivity disorder. Clinical Neurophysiology, 2014, 125, 1407-1416.	1.5	126
18	Social Problem Solving in Adolescents with Suicidal Behavior: A Systematic Review. Suicide and Life-Threatening Behavior, 2005, 35, 365-387.	1.9	124

#	Article	IF	CITATIONS
19	Cognitive–behavioural therapy for depression in people with a somatic disease: meta-analysis of randomised controlled trials. British Journal of Psychiatry, 2010, 197, 11-19.	2.8	120
20	Intrusive memories and rumination in patients with post-traumatic stress disorder: A phenomenological comparison. Memory, 2007, 15, 249-257.	1.7	115
21	Imagery special issue: Intrusive images and memories of earlier adverse events in patients with obsessive compulsive disorder. Journal of Behavior Therapy and Experimental Psychiatry, 2007, 38, 411-422.	1.2	104
22	Mindfulnessâ€based stress reduction added to care as usual for lung cancer patients and/or their partners: A multicentre randomized controlled trial. Psycho-Oncology, 2017, 26, 2118-2126.	2.3	94
23	On the Relationship Between the Practice of Mindfulness Meditation and Personality—an Exploratory Analysis of the Mediating Role of Mindfulness Skills. Mindfulness, 2011, 2, 194-200.	2.8	92
24	Unexplained physical symptoms: outcome, utilization of medical care and associated factors. Psychological Medicine, 1996, 26, 745-752.	4.5	87
25	A Workplace Mindfulness Intervention May Be Associated With Improved Psychological Well-Being and Productivity. A Preliminary Field Study in a Company Setting. Frontiers in Psychology, 2018, 9, 195.	2.1	86
26	Individual differences in disgust sensitivity modulate neural responses to aversive/disgusting stimuli. European Journal of Neuroscience, 2008, 27, 3050-3058.	2.6	85
27	What is the prevalence of fear of cancer recurrence in cancer survivors and patients? A systematic review and individual participant data metaâ€analysis. Psycho-Oncology, 2022, 31, 879-892.	2.3	85
28	Changes in intrusive memories associated with imaginal reliving in posttraumatic stress disorder. Journal of Anxiety Disorders, 2006, 20, 328-341.	3.2	83
29	The diagnostic and prognostic significance of the Whitely Index, the Illness Attitude Scales and the Somatosensory Amplification Scale. Psychological Medicine, 1996, 26, 1085-1090.	4.5	82
30	Neural Responses to Facial Expressions of Disgust but not Fear are Modulated by Washing Symptoms in OCD. Biological Psychiatry, 2007, 61, 1072-1080.	1.3	82
31	Predictors of suicidality in depressive spectrum disorders in the general population: results of the Netherlands Mental Health Survey and Incidence Study. Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 513-521.	3.1	78
32	Mindfulness-Based Stress Reduction for Residents: A Randomized Controlled Trial. Journal of General Internal Medicine, 2018, 33, 429-436.	2.6	74
33	The reassuring value of diagnostic tests: A systematic review. Patient Education and Counseling, 2012, 86, 3-8.	2.2	73
34	The influence of mood on the processing of syntactic anomalies: Evidence from P600. Neuropsychologia, 2010, 48, 3521-3531.	1.6	68
35	Mindfulness-Based Stress Reduction for lung cancer patients and their partners: Results of a mixed methods pilot study. Palliative Medicine, 2015, 29, 652-660.	3.1	68
36	Mindfulness-based cognitive therapy for patients with chronic, treatment-resistant depression: A pragmatic randomized controlled trial. Depression and Anxiety, 2018, 35, 914-924.	4.1	67

#	Article	IF	CITATIONS
37	Psychosexual functioning of partners of men with presumed non-organic erectile dysfunction: Cause or consequence of the disorder?. Archives of Sexual Behavior, 1995, 24, 157-172.	1.9	61
38	Predicting response to cognitive behavioral therapy in contamination-based obsessive–compulsive disorder from functional magnetic resonance imaging. Psychological Medicine, 2014, 44, 2125-2137.	4.5	59
39	Mindfulness-based stress reduction for GPs: results of a controlled mixed methods pilot study in Dutch primary care. British Journal of General Practice, 2016, 66, e99-e105.	1.4	57
40	Mindfulness-Based Cognitive Therapy for Patients with Medically Unexplained Symptoms: A Randomized Controlled Trial. Psychotherapy and Psychosomatics, 2013, 82, 299-310.	8.8	56
41	Effects of Mindfulness-Based Stress Reduction on the Mental Health of Clinical Clerkship Students: A Cluster-Randomized Controlled Trial. Academic Medicine, 2017, 92, 1012-1021.	1.6	56
42	Are Mindfulness and Self-Compassion Related to Psychological Distress and Communication in Couples Facing Lung Cancer? A Dyadic Approach. Mindfulness, 2017, 8, 325-336.	2.8	55
43	Sex Differences in Neural Responses to Disgusting Visual Stimuli: Implications for Disgust-Related Psychiatric Disorders. Biological Psychiatry, 2007, 62, 464-471.	1.3	53
44	Changes in Attentional Processing of Emotional Information Following Mindfulness-Based Cognitive Therapy in People with a History of Depression: Towards an Open Attention for all Emotional Experiences. Cognitive Therapy and Research, 2012, 36, 612-620.	1.9	52
45	The Efficacy of Adapted MBCT on Core Symptoms and Executive Functioning in Adults With ADHD: A Preliminary Randomized Controlled Trial. Journal of Attention Disorders, 2019, 23, 351-362.	2.6	52
46	Discontinuation of antidepressant medication after mindfulness-based cognitive therapy for recurrent depression: Randomised controlled non-inferiority trial. British Journal of Psychiatry, 2016, 208, 366-373.	2.8	51
47	The Effectiveness of Mindfulness-Based Stress Reduction on Psychological Distress and Cognitive Functioning in Patients with Multiple Sclerosis: a Pilot Study. Mindfulness, 2017, 8, 1251-1258.	2.8	49
48	Prognostic association of cardiac anxiety with new cardiac events and mortality following myocardial infarction. British Journal of Psychiatry, 2016, 209, 400-406.	2.8	46
49	The acceptability of psychological treatment in patients with medically unexplained physical symptoms. Journal of Psychosomatic Research, 1995, 39, 855-863.	2.6	45
50	Preventing relapse in recurrent depression using mindfulness-based cognitive therapy, antidepressant medication or the combination: trial design and protocol of the MOMENT study. BMC Psychiatry, 2012, 12, 125.	2.6	45
51	How mood challenges emotional memory formation: An fMRI investigation. NeuroImage, 2011, 56, 1783-1790.	4.2	44
52	Mindfulness and mood stimulate each other in an upward spiral: a mindful walking intervention using experience sampling. Mindfulness, 2016, 7, 1114-1122.	2.8	44
53	Mindfulness-based cognitive therapy <i>v</i> . treatment as usual in adults with ADHD: a multicentre, single-blind, randomised controlled trial. Psychological Medicine, 2019, 49, 55-65.	4.5	43
54	The Maudsley Obsessive–Compulsive Stimuli Set: Validation of a standardized paradigm for symptom-specific provocation in obsessive–compulsive disorder. Psychiatry Research, 2009, 168, 238-241.	3.3	42

#	Article	IF	CITATIONS
55	Fronto-limbic microstructure and structural connectivity in remission from major depression. Psychiatry Research - Neuroimaging, 2012, 204, 40-48.	1.8	41
56	Mindfulness-based cognitive therapy for patients with medically unexplained symptoms: A cost-effectiveness study. Journal of Psychosomatic Research, 2013, 74, 197-205.	2.6	41
57	The Role of the Teacher in Mindfulness-Based Approaches: A Qualitative Study. Mindfulness, 2014, 5, 170-178.	2.8	41
58	The suitability of the Hospital Anxiety and Depression Scale, Distress Thermometer and other instruments to screen for psychiatric disorders in both lung cancer patients and their partners. Journal of Affective Disorders, 2016, 203, 176-183.	4.1	41
59	The contribution of work characteristics, home characteristics and gender to burnout in medical residents. Advances in Health Sciences Education, 2017, 22, 803-818.	3.3	41
60	The Reassurance Questionnaire (RQ): psychometric properties of a self-report questionnaire to assess reassurability. Psychological Medicine, 2000, 30, 841-847.	4.5	40
61	Improvement of mindfulness skills during Mindfulness-Based Cognitive Therapy predicts long-term reductions of neuroticism in persons with recurrent depression in remission. Journal of Affective Disorders, 2017, 213, 112-117.	4.1	40
62	Approach and Avoidance of Emotional Faces in Happy and Sad Mood. Cognitive Therapy and Research, 2013, 37, 1-6.	1.9	36
63	Adding mindfulness-based cognitive therapy to maintenance antidepressant medication for prevention of relapse/recurrence in major depressive disorder: Randomised controlled trial. Journal of Affective Disorders, 2015, 187, 54-61.	4.1	36
64	Discrimination between psychogenic and organic erectile dysfunction. Journal of Psychosomatic Research, 1993, 37, 135-145.	2.6	35
65	Study protocol of the SWORD-study: a randomised controlled trial comparing combined online and face-to-face cognitive behaviour therapy versus treatment as usual in managing fear of cancer recurrence. BMC Psychology, 2015, 3, 12.	2.1	35
66	Does Mindfulness Training Enhance the Professional Development of Residents? A Qualitative Study. Academic Medicine, 2018, 93, 1335-1340.	1.6	35
67	Consolidation and prediction of long-term treatment effect of group and online mindfulness-based cognitive therapy for distressed cancer patients. Acta Oncológica, 2018, 57, 1293-1302.	1.8	35
68	Stress and mindfulness in Parkinson's disease – a survey in 5000 patients. Npj Parkinson's Disease, 2021, 7, 7.	5.3	35
69	Medically unexplained physical symptoms. Journal of Psychosomatic Research, 2004, 57, 517-520.	2.6	34
70	Shorter gaze duration for happy faces in current but not remitted depression: Evidence from eye movements. Psychiatry Research, 2014, 218, 79-86.	3.3	34
71	Prevalence and correlates of lifetime deliberate self-harm and suicidal ideation in naturalistic outpatients: The Leiden Routine Outcome Monitoring study. Journal of Affective Disorders, 2011, 133, 257-264.	4.1	33
72	Stress and Mindfulness in Parkinson's Disease: Clinical Effects and Potential Underlying Mechanisms. Movement Disorders, 2021, 36, 64-70.	3.9	33

#	Article	IF	CITATIONS
73	A qualitative study on mindfulness-based stress reduction for breast cancer patients: how women experience participating with fellow patients. Supportive Care in Cancer, 2016, 24, 1813-1820.	2.2	32
74	Long-term efficacy and cost-effectiveness of blended cognitive behavior therapy for high fear of recurrence in breast, prostate and colorectal Cancer survivors: follow-up of the SWORD randomized controlled trial. BMC Cancer, 2019, 19, 462.	2.6	32
75	Withdrawal of unnecessary antidepressant medication: a randomised controlled trial in primary care. BJGP Open, 2018, 1, bjgpopen17X101265.	1.8	32
76	A BRIEF COGNITIVE-BEHAVIORAL INTERVENTION FOR TREATING DEPRESSION AND PANIC DISORDER IN PATIENTS WITH NONCARDIAC CHEST PAIN: A 24-WEEK RANDOMIZED CONTROLLED TRIAL. Depression and Anxiety, 2013, 30, 670-678.	4.1	31
77	Rose or black-coloured glasses?. Journal of Affective Disorders, 2011, 131, 214-223.	4.1	29
78	One way or another: The opportunities and pitfalls of selfâ€referral and consecutive sampling as recruitment strategies for psychoâ€oncology intervention trials. Psycho-Oncology, 2018, 27, 2056-2059.	2.3	29
79	Multi-dimensional modulations of $\hat{l}\pm$ and $\hat{l}^3$ cortical dynamics following mindfulness-based cognitive therapy in Major Depressive Disorder. Cognitive Neurodynamics, 2015, 9, 13-29.	4.0	28
80	Teacher Competence in Mindfulness-Based Cognitive Therapy for Depression and Its Relation to Treatment Outcome. Mindfulness, 2017, 8, 960-972.	2.8	28
81	Study protocol of a randomized controlled trial comparing Mindfulness-Based Stress Reduction with treatment as usual in reducing psychological distress in patients with lung cancer and their partners: the MILON study. BMC Cancer, 2014, 14, 3.	2.6	27
82	Patients' attitudes to discontinuing not-indicated long-term antidepressant use: barriers and facilitators. Therapeutic Advances in Psychopharmacology, 2019, 9, 204512531987234.	2.7	27
83	The Cardiac Anxiety Questionnaire: Cross-Validation among Cardiac Inpatients. International Journal of Psychiatry in Medicine, 2012, 43, 349-364.	1.8	26
84	Verbal and facialâ€emotional Stroop tasks reveal specific attentional interferences in sad mood. Brain and Behavior, 2012, 2, 74-83.	2.2	26
85	Mindfulness-based cognitive therapy (MBCT) for patients with medically unexplained symptoms: Process of change. Journal of Psychosomatic Research, 2014, 77, 27-33.	2.6	26
86	Predictors and Effects of Usage of an Online Mindfulness Intervention for Distressed Cancer Patients: Usability Study. Journal of Medical Internet Research, 2020, 22, e17526.	4.3	25
87	Study protocol of a multicenter randomized controlled trial comparing the effectiveness of group and individual internet-based Mindfulness-Based Cognitive Therapy with treatment as usual in reducing psychological distress in cancer patients: the BeMind study. BMC Psychology, 2015, 3, 27.	2.1	24
88	Costâ€utility of individual internetâ€based and faceâ€toâ€face Mindfulnessâ€Based Cognitive Therapy compared with treatment as usual in reducing psychological distress in cancer patients. Psycho-Oncology, 2020, 29, 294-303.	2.3	24
89	A randomised controlled trial (MindChamp) of a mindfulnessâ€based intervention for children with ADHD and their parents. Journal of Child Psychology and Psychiatry and Allied Disciplines, 2022, 63, 165-177.	5.2	24
90	Psychiatric diagnosis by telephone: is it an opportunity?. Social Psychiatry and Psychiatric Epidemiology, 2014, 49, 1677-1689.	3.1	23

#	Article	IF	CITATIONS
91	Medically Unexplained Physical Symptoms in Elderly People: A Pilot Study of Psychiatric Geriatric Characteristics. American Journal of Geriatric Psychiatry, 2009, 17, 1085-1088.	1.2	22
92	An Investigation of the Role of Attention in Mindfulness-Based Cognitive Therapy for Recurrently Depressed Patients. Journal of Experimental Psychopathology, 2012, 3, 103-120.	0.8	22
93	Depressed patients in remission show an interaction between variance in the mineralocorticoid receptor NR3C2 gene and childhood trauma on negative memory bias. Psychiatric Genetics, 2015, 25, 99-105.	1.1	22
94	Mindfulness-based stress reduction for breast cancer patients: a mixed method study on what patients experience as a suitable stage to participate. Supportive Care in Cancer, 2017, 25, 3067-3074.	2.2	22
95	Modulation of induced frontocentral theta (Fm-Î) event-related (de-)synchronisation dynamics following mindfulness-based cognitive therapy in Major Depressive Disorder. Cognitive Neurodynamics, 2014, 8, 373-388.	4.0	21
96	Mindfulness-Based Cognitive Therapy for Cancer Patients Delivered via Internet: Qualitative Study of Patient and Therapist Barriers and Facilitators. Journal of Medical Internet Research, 2017, 19, e407.	4.3	21
97	Association between genes, stressful childhood events and processing bias in depression vulnerable individuals. Genes, Brain and Behavior, 2014, 13, 508-516.	2.2	20
98	Mindfulness based cognitive therapy versus treatment as usual in adults with attention deficit hyperactivity disorder (ADHD). BMC Psychiatry, 2015, 15, 216.	2.6	20
99	Exploring associations between psychiatric disorder, psychological distress, and health care utilization in cancer patients. Psycho-Oncology, 2018, 27, 871-878.	2.3	20
100	Development of the Therapeutic Alliance and its Association With Internet-Based Mindfulness-Based Cognitive Therapy for Distressed Cancer Patients: Secondary Analysis of a Multicenter Randomized Controlled Trial. Journal of Medical Internet Research, 2019, 21, e14065.	4.3	20
101	Response time variability and response inhibition predict affective problems in adolescent girls, not in boys: the TRAILS study. European Child and Adolescent Psychiatry, 2012, 21, 277-287.	4.7	19
102	LONG-TERM OUTCOME OF MINDFULNESS-BASED COGNITIVE THERAPY IN RECURRENTLY DEPRESSED PATIENTS WITH AND WITHOUT A DEPRESSIVE EPISODE AT BASELINE. Depression and Anxiety, 2015, 32, 563-569.	4.1	19
103	Reliability and validity of the Dutch version of the Consultation and Relational Empathy Measure in primary care. Family Practice, 2017, 34, 119-124.	1.9	19
104	Eyewitness or Earwitness: The Role of Mental Imagery in Intrusion Development. International Journal of Cognitive Therapy, 2011, 4, 154-164.	2.2	18
105	Cost-utility analysis of a treatment advice to discontinue inappropriate long-term antidepressant use in primary care. Family Practice, 2014, 31, 578-584.	1.9	18
106	What is the contribution of different cognitive biases and stressful childhood events to the presence and number of previous depressive episodes?. Psychiatry Research, 2014, 217, 134-142.	3.3	18
107	Cognitive rehabilitation and mindfulness in multiple sclerosis (REMIND-MS): a study protocol for a randomised controlled trial. BMC Neurology, 2017, 17, 201.	1.8	18
108	A cross-sectional examination of psychological distress, positive mental health and their predictors in medical students in their clinical clerkships. BMC Medical Education, 2017, 17, 219.	2.4	18

7

#	Article	IF	CITATIONS
109	Therapeutic alliance—not therapist competence or group cohesion—contributes to reduction of psychological distress in groupâ€based mindfulnessâ€based cognitive therapy for cancer patients. Clinical Psychology and Psychotherapy, 2019, 26, 309-318.	2.7	18
110	One-year follow up of cardiac anxiety after a myocardial infarction: A latent class analysis. Journal of Psychosomatic Research, 2012, 73, 362-368.	2.6	17
111	Mindfulness training for medical students in their clinical clerkships: two cross-sectional studies exploring interest and participation. BMC Medical Education, 2015, 15, 24.	2.4	17
112	Implicit and explicit self-esteem in remitted depressed patients. Journal of Behavior Therapy and Experimental Psychiatry, 2017, 54, 301-306.	1.2	17
113	Alcohol abuse among elderly patients in a general hospital as identified by the Munich Alcoholism Test. Acta Psychiatrica Scandinavica, 1991, 83, 460-462.	4.5	15
114	Attention Effects of Mindfulness-Based Cognitive Therapy in Formerly Depressed Patients. Journal of Experimental Psychopathology, 2014, 5, 414-424.	0.8	15
115	Can Memory Bias be Modified? The Effects of an Explicit Cued-Recall Training in Two Independent Samples. Cognitive Therapy and Research, 2014, 38, 217-225.	1.9	15
116	Cognitive reactivity as outcome and working mechanism of mindfulness-based cognitive therapy for recurrently depressed patients in remission. Cognition and Emotion, 2018, 32, 371-378.	2.0	15
117	Effectiveness and feasibility of a mindful leadership course for medical specialists: a pilot study. BMC Medical Education, 2020, 20, 34.	2.4	15
118	Effects of Mindfulness-Based Cognitive Therapy on a Behavioural Measure of Rumination in Patients with Chronic, Treatment-Resistant Depression. Cognitive Therapy and Research, 2019, 43, 666-678.	1.9	14
119	The Feasibility, Effectiveness, and Process of Change of Mindfulness-Based Cognitive Therapy for Adults With ADHD: A Mixed-Method Pilot Study. Journal of Attention Disorders, 2020, 24, 928-942.	2.6	14
120	Recovery from recurrent depression: Randomized controlled trial of the efficacy of mindfulness-based compassionate living compared with treatment-as-usual on depressive symptoms and its consolidation at longer term follow-up. Journal of Affective Disorders, 2020, 273, 265-273.	4.1	14
121	Cognitive Behavioural Therapy for Medically Unexplained Physical Symptoms: A Pilot Study. Behavioural and Cognitive Psychotherapy, 1996, 24, 1-16.	1.2	13
122	The impact of comorbid depression on recovery from personality disorders and improvements in psychosocial functioning: Results from a randomized controlled trial. Behaviour Research and Therapy, 2014, 63, 55-62.	3.1	13
123	A randomized controlled trial of Mindfulness-Based Cognitive Therapy (MBCT) versus treatment-as-usual (TAU) for chronic, treatment-resistant depression: study protocol. BMC Psychiatry, 2015, 15, 275.	2.6	13
124	Enhancing shared decision making about discontinuation of antidepressant medication: a concept-mapping study in primary and secondary mental health care. British Journal of General Practice, 2019, 69, e777-e785.	1.4	13
125	Mechanisms of Change in Mindfulness-Based Cognitive Therapy in Adults With ADHD. Journal of Attention Disorders, 2021, 25, 1331-1342.	2.6	13
126	Mindfulness facets and Big Five personality facets in persons with recurrent depression in remission. Personality and Individual Differences, 2017, 110, 109-114.	2.9	12

#	Article	IF	Citations
127	The Co-creation and Feasibility of a Compassion Training as a Follow-up to Mindfulness-Based Cognitive Therapy in Patients with Recurrent Depression. Mindfulness, 2018, 9, 412-422.	2.8	12
128	Automatic approach-avoidance tendencies as a candidate intermediate phenotype for depression: Associations with childhood trauma and the 5-HTTLPR transporter polymorphism. PLoS ONE, 2018, 13, e0193787.	2.5	11
129	A parallel-group, randomized controlled trial into the effectiveness of Mindfulness-Based Compassionate Living (MBCL) compared to treatment-as-usual in recurrent depression: Trial design and protocol. Contemporary Clinical Trials, 2016, 50, 77-83.	1.8	10
130	Mindfulness for Children With ADHD and Mindful Parenting (MindChamp): A Qualitative Study on Feasibility and Effects. Journal of Attention Disorders, 2021, 25, 1931-1942.	2.6	10
131	Cognitive Behavioural Therapy for Unexplained Physical Symptoms: Process and Prognostic Factors. Behavioural and Cognitive Psychotherapy, 1997, 25, 291-294.	1.2	9
132	Prescribing ANtiDepressants Appropriately (PANDA): a cluster randomized controlled trial in primary care. BMC Family Practice, 2013, 14, 6.	2.9	9
133	Inverse Correlation Between Cardiac Injury and Cardiac Anxiety. Journal of Cardiovascular Nursing, 2014, 29, 448-453.	1.1	9
134	Patients with a preference for medication do equally well in mindfulness-based cognitive therapy for recurrent depression as those preferring mindfulness. Journal of Affective Disorders, 2016, 195, 32-39.	4.1	9
135	The relationship between work and home characteristics and work engagement in medical residents. Perspectives on Medical Education, 2017, 6, 227-236.	3.5	9
136	Silent into Nature: Factors Enabling Improvement in a Mindful Walking Retreat in Nature of People with Psychological Symptoms. Ecopsychology, 2018, 10, 77-86.	1.4	9
137	Discontinuation of antidepressant medication in primary care supported by monitoring plus mindfulness-based cognitive therapy versus monitoring alone: design and protocol of a cluster randomized controlled trial. BMC Family Practice, 2019, 20, 105.	2.9	9
138	Putting mindfulness-based cognitive therapy to the test in routine clinical practice: A transdiagnostic panacea or a disorder specific intervention? Journal of Psychiatric Research, 2021, 142, 144-152.	3.1	9
139	Alcohol abuse among general hospital inpatients according to the Munich Alcoholism Test (MALT). General Hospital Psychiatry, 1994, 16, 125-130.	2.4	8
140	Greater mindful eating practice is associated with better reversal learning. Scientific Reports, 2018, 8, 5702.	3.3	8
141	The effectiveness of mindfulness-based cognitive therapy for major depressive disorder: evidence from routine outcome monitoring data. BJPsych Open, 2020, 6, e144.	0.7	8
142	Discontinuing antidepressant medication after mindfulness-based cognitive therapy: a mixed-methods study exploring predictors and outcomes of different discontinuation trajectories, and its facilitators and barriers. BMJ Open, 2020, 10, e039053.	1.9	8
143	Validation of the Dutch Comprehensive Inventory of Mindfulness Experiences (CHIME) and Development of a Short Form (CHIME-SF). Mindfulness, 2019, 10, 1893-1904.	2.8	7
144	Genetic and environmental aetiologies of associations between dispositional mindfulness and ADHD traits: a population-based twin study. European Child and Adolescent Psychiatry, 2019, 28, 1241-1251.	4.7	7

#	Article	IF	Citations
145	Factors associated with relapse and recurrence of major depressive disorder in patients starting mindfulnessâ€based cognitive therapy. Depression and Anxiety, 2022, 39, 113-122.	4.1	7
146	A qualitative analysis of experiences of patients with metastatic breast cancer participating in a mindfulness-based intervention. Palliative Medicine, 2015, 29, 182-183.	3.1	6
147	Added value of Mindfulness-Based Cognitive Therapy for Depression: A Tree-based Qualitative Interaction Analysis. Behaviour Research and Therapy, 2019, 122, 103467.	3.1	6
148	Coherence Between Attentional and Memory Biases in Sad and Formerly Depressed Individuals. Cognitive Therapy and Research, 2014, 38, 334.	1.9	5
149	How medical specialists experience the effects of a mindful leadership course on their leadership capabilities: a qualitative interview study in the Netherlands. BMJ Open, 2019, 9, e031643.	1.9	5
150	Mindfulness and Affect During Mindfulness-Based Cognitive Therapy for Recurrent Depression: an Autoregressive Latent Trajectory Analysis. Mindfulness, 2020, 11, 2360-2370.	2.8	5
151	Prospective Associations Between Home Practice and Depressive Symptoms in Mindfulness-Based Cognitive Therapy for Recurrent Depression: A 15 Months Follow-Up Study. Cognitive Therapy and Research, 2021, 45, 250-261.	1.9	5
152	Ongedifferentieerde somatoforme stoornis. , 2001, , 17-40.		5
153	The feasibility of mindfulness-based cognitive therapy for people with bipolar disorder: a qualitative study. International Journal of Bipolar Disorders, 2020, 8, 33.	2.2	5
154	Adverse or therapeutic? A mixed-methods study investigating adverse effects of Mindfulness-Based Cognitive Therapy in bipolar disorder. PLoS ONE, 2021, 16, e0259167.	2.5	5
155	Exploring Fear of Cancer Recurrence in a Sample of Heterogeneous Distressed Cancer Patients with and Without a Psychiatric Disorder. Journal of Clinical Psychology in Medical Settings, 2021, 28, 419-426.	1.4	4
156	Experiences of Clinical Clerkship Students With Mindfulness-Based Stress Reduction: A Qualitative Study on Long-Term Effects. Frontiers in Psychology, 2022, 13, 785090.	2.1	4
157	Mindfulness-Based Compassionate Living (MBCL): a Qualitative Study into the Added Value of Compassion in Recurrent Depression. Mindfulness, 2021, 12, 2196-2206.	2.8	3
158	Perspectives of bereaved partners of lung cancer patients on the role of mindfulness in dying and grieving: A qualitative study. Palliative Medicine, 2021, 35, 200-208.	3.1	3
159	Temporal Changes in Mindfulness Skills and Positive and Negative Affect and Their Interrelationships During Mindfulness-Based Cognitive Therapy for Cancer Patients. Mindfulness, 2022, 13, 1745-1756.	2.8	3
160	De behandeling van onverklaarde lichamelijke klachten. Dth, 1996, 16, 170-174.	0.2	2
161	The Prognostic Effect of Physical Health Complaints With New Cardiac Events and Mortality in Patients With a Myocardial Infarction. Psychosomatics, 2017, 58, 121-131.	2.5	2
162	P3.05-008 Mindfulness-Based Stress Reduction Added to Care as Usual for Lung CancerÂPatients and Their Partners: AÂRandomized Controlled Trial. Journal of Thoracic Oncology, 2017, 12, S1416-S1417.	1.1	2

#	Article	IF	CITATIONS
163	Mindfulness-Based Cognitive Therapy Versus Treatment as Usual in Adults with ADHD: a Trial-Based Economic Evaluation. Mindfulness, 2019, 10, 1803-1814.	2.8	2
164	Interplay between self-compassion and affect during Mindfulness-Based Compassionate Living for recurrent depression: An Autoregressive Latent Trajectory analysis. Behaviour Research and Therapy, 2021, 146, 103946.	3.1	2
165	Facilitated Attentional Disengagement from Negative Information in Relation to Self-Reported Depressive Symptoms of Dutch Female Undergraduate Students. Psychological Reports, 2011, 108, 252-262.	1.7	1
166	No Evidence for the Association between a Polymorphism in the PCLO Depression Candidate Gene with Memory Bias in Remitted Depressed Patients and Healthy Individuals. PLoS ONE, 2014, 9, e112153.	2.5	1
167	Comments on â€~ <i>Psychological intervention targeting distress for cancer patients: a metaâ€analytic study investigating uptake and adherence</i> <ir> <li>i&gt;'</li> <li>by Brebach and colleagues (2016). Psycho-Oncology, 2016, 25, 997-997.</li> </ir>	2.3	1
168	The Process of Change of Mindfulness-Based Cognitive Therapy for People with Bipolar Disorder: a Qualitative Study. Mindfulness, 2021, 12, 863-872.	2.8	1
169	A multi-method assessment of attentional processes in chronic, treatment-resistant depression. Journal of Psychiatric Research, 2021, 140, 68-76.	3.1	1
170	Too Much or Too Little Antidepressant Medication: Difficult to Change. Two Rcts. Mental Health in Family Medicine, 2017, 13, .	0.2	1
171	Mindfulness-Based Interventions. , 2020, , 1396-1399.		1
172	Discontinuing antidepressant medication after mindfulness-based cognitive therapy: a mixed-methods study exploring predictors and outcomes of different discontinuation trajectories, and its facilitators and barriers. BMJ Open, 2020, 10, e039053.	1.9	1
173	Mindfulness-based cognitive therapy as an alternative to maintenance antidepressant medication to prevent relapse and recurrence in depression. Evidence-Based Mental Health, 2015, 18, 126-126.	4.5	0
174	Preventive cognitive therapy could be a viable and effective addition to antidepressant medication in preventing relapse or recurrence in major depressive disorder. Evidence-Based Mental Health, 2019, 22, e7-e7.	4.5	0
175	Mindfulness-Based Interventions. , 2018, , 1-4.		0