

# Shaobo Wang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1321297/publications.pdf>

Version: 2024-02-01

21  
papers

1,170  
citations

567281

15  
h-index

713466

21  
g-index

25  
all docs

25  
docs citations

25  
times ranked

933  
citing authors

#	ARTICLE	IF	CITATIONS
1	EWSR1-induced circNEIL3 promotes glioma progression and exosome-mediated macrophage immunosuppressive polarization via stabilizing IGF2BP3. <i>Molecular Cancer</i> , 2022, 21, 16.	19.2	115
2	SPI1-induced downregulation of FTO promotes GBM progression by regulating pri-miR-10a processing in an m6A-dependent manner. <i>Molecular Therapy - Nucleic Acids</i> , 2022, 27, 699-717.	5.1	23
3	Comprehensive Analysis of the Tumor Immune Microenvironment Landscape in Glioblastoma Reveals Tumor Heterogeneity and Implications for Prognosis and Immunotherapy. <i>Frontiers in Immunology</i> , 2022, 13, 820673.	4.8	10
4	A Comprehensive Analysis of METTL1 to Immunity and Stemness in Pan-Cancer. <i>Frontiers in Immunology</i> , 2022, 13, 795240.	4.8	15
5	miR-184-3p enriched in cerebrospinal fluid exosomes contributes to progression of glioma and promotes M2-like macrophage polarization. <i>Cancer Science</i> , 2022, 113, 2668-2680.	3.9	13
6	The dual role of glioma exosomal microRNAs: glioma eliminates tumor suppressor miR-1298-5p via exosomes to promote immunosuppressive effects of MDSCs. <i>Cell Death and Disease</i> , 2022, 13, 426.	6.3	32
7	Exosomes derived from hypoxic glioma deliver miR-1246 and miR-10b-5p to normoxic glioma cells to promote migration and invasion. <i>Laboratory Investigation</i> , 2021, 101, 612-624.	3.7	28
8	MicroRNA-29a-3p delivery via exosomes derived from engineered human mesenchymal stem cells exerts tumour suppressive effects by inhibiting migration and vasculogenic mimicry in glioma. <i>Aging</i> , 2021, 13, 5055-5068.	3.1	37
9	Hypoxic glioma-derived exosomes promote M2-like macrophage polarization by enhancing autophagy induction. <i>Cell Death and Disease</i> , 2021, 12, 373.	6.3	93
10	Qki activates Srebp2-mediated cholesterol biosynthesis for maintenance of eye lens transparency. <i>Nature Communications</i> , 2021, 12, 3005.	12.8	22
11	The N6-Methyladenosine-Modified Pseudogene HSPA7 Correlates With the Tumor Microenvironment and Predicts the Response to Immune Checkpoint Therapy in Glioblastoma. <i>Frontiers in Immunology</i> , 2021, 12, 653711.	4.8	25
12	Exosomal miR-1246 from glioma patient body fluids drives the differentiation and activation of myeloid-derived suppressor cells. <i>Molecular Therapy</i> , 2021, 29, 3449-3464.	8.2	47
13	Cell surface GRP78 regulates BACE2 via lysosome-dependent manner to maintain mesenchymal phenotype of glioma stem cells. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021, 40, 20.	8.6	17
14	The Non-N6-Methyladenosine Epitranscriptome Patterns and Characteristics of Tumor Microenvironment Infiltration and Mesenchymal Transition in Glioblastoma. <i>Frontiers in Immunology</i> , 2021, 12, 809808.	4.8	11
15	Hypoxic glioma-derived exosomes deliver microRNA-1246 to induce M2 macrophage polarization by targeting TERF2IP via the STAT3 and NF- $\kappa$ B pathways. <i>Oncogene</i> , 2020, 39, 428-442.	5.9	223
16	TGF $\beta$ -induced beta-site APP-cleaving enzyme 2 upregulation promotes tumorigenesis through the NF- $\kappa$ B signalling pathway in human gliomas. <i>Molecular Oncology</i> , 2020, 14, 407-425.	4.6	14
17	Transfer of MicroRNA via Macrophage-Derived Extracellular Vesicles Promotes Proneural-to-Mesenchymal Transition in Glioma Stem Cells. <i>Cancer Immunology Research</i> , 2020, 8, 966-981.	3.4	55
18	Hypoxia-induced lncRNA PDIA3P1 promotes mesenchymal transition via sponging of miR-124-3p in glioma. <i>Cell Death and Disease</i> , 2020, 11, 168.	6.3	40

#	ARTICLE	IF	CITATIONS
19	Cullin-7 (CUL7) is overexpressed in glioma cells and promotes tumorigenesis via NF- $\kappa$ B activation. Journal of Experimental and Clinical Cancer Research, 2020, 39, 59.	8.6	41
20	Glioma exosomes mediate the expansion and function of myeloid-derived suppressor cells through microRNA-29a/Hbp1 and microRNA-92a/Prkar1a pathways. International Journal of Cancer, 2019, 144, 3111-3126.	5.1	107
21	Immunosuppressive effects of hypoxia-induced glioma exosomes through myeloid-derived suppressor cells via the miR-10a/Rora and miR-21/Pten Pathways. Oncogene, 2018, 37, 4239-4259.	5.9	202