Tomoko Takahashi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1338571/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Successful use of prednisolone and radiation therapy in a dog with intracranial histiocytic sarcoma. Journal of Veterinary Medical Science, 2021, 83, 1782-1785.	0.9	0
2	Interactions between the non-seed region of siRNA and RNA-binding RLC/RISC proteins, Ago and TRBP, in mammalian cells. Nucleic Acids Research, 2014, 42, 5256-5269.	14.5	27
3	Role of .BETA.1 Integrins in Adhesion of Canine Mastocytoma Cells to Extracellular Matrix Proteins. Journal of Veterinary Medical Science, 2007, 69, 495-499.	0.9	3
4	Malignant Transformation of T-cell Large Granular Lymphocyte Leukemia in a Dog. Journal of Veterinary Medical Science, 2007, 69, 677-681.	0.9	8
5	Retinoids Induce Growth Inhibition and Apoptosis in Mast Cell Tumor Cell Lines. Journal of Veterinary Medical Science, 2006, 68, 797-802.	0.9	16
6	Molecular cloning of the canine telomerase reverse transcriptase gene and its expression in neoplastic and non-neoplastic cells. American Journal of Veterinary Research, 2003, 64, 1395-1400.	0.6	13
7	IgG-Mediated Histamine Release from Canine Mastocytoma-Derived Cells. International Archives of Allergy and Immunology, 2001, 125, 228-235.	2.1	34
8	Effect of Retinoids on Growth Inhibition of Two Canine Melanoma Cell Lines Journal of Veterinary Medical Science, 2001, 63, 83-86.	0.9	29
9	Visceral mast cell tumors in dogs: 10 cases (1982-1997). Journal of the American Veterinary Medical Association, 2000, 216, 222-226.	0.5	74
10	VEGF activates protein kinase C-dependent, but Ras-independent Raf-MEK-MAP kinase pathway for DNA synthesis in primary endothelial cells. Oncogene, 1999, 18, 2221-2230.	5.9	524
11	Virally activated ras cooperates with integrin to induce tubulogenesis in sinusoidal endothelial cell lines. Journal of Cellular Physiology, 1998, 176, 223-234.	4.1	35
12	Inhibitory Effects of Glucocorticoids on Proliferation of Canine Mast Cell Tumor Journal of Veterinary Medical Science, 1997, 59, 995-1001.	0.9	32
13	The 230 kDa mature form of KDR/Flk-1 (VEGF receptor-2) activates the PLC-γ pathway and partially induces mitotic signals in NIH3T3 fibroblasts. Oncogene, 1997, 14, 2079-2089.	5.9	278