

Francesca Rigano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1383277/publications.pdf>

Version: 2024-02-01

173
papers

5,545
citations

61984

43
h-index

128289

60
g-index

176
all docs

176
docs citations

176
times ranked

5011
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Determination of the polyphenolic content of berry juices using focusing-modulated comprehensive two-dimensional liquid chromatography coupled to mass spectrometry detection. <i>Analytical and Bioanalytical Chemistry</i> , 2023, 415, 2371-2382. | 3.7 | 5 |
| 2 | Biodegradation Potential of Oil-degrading Bacteria Related to the Genus <i>Thalassospira</i> Isolated from Polluted Coastal Area in Mediterranean Sea. <i>Soil and Sediment Contamination</i> , 2022, 31, 316-332. | 1.9 | 6 |
| 3 | On-line coupling of supercritical fluid extraction with enantioselective supercritical fluid chromatography-triple quadrupole mass spectrometry for the determination of chiral pesticides in hemp seeds: A proof-of-principle study. <i>Food Chemistry</i> , 2022, 373, 131418. | 8.2 | 6 |
| 4 | Untargeted profiling and differentiation of geographical variants of wine samples using headspace solid-phase microextraction flow-modulated comprehensive two-dimensional gas chromatography with the support of tile-based Fisher ratio analysis. <i>Journal of Chromatography A</i> , 2022, 1662, 462735. | 3.7 | 23 |
| 5 | Non-psychoactive cannabinoids identification by linear retention index approach applied to a hand-portable capillary liquid chromatography platform. <i>Analytical and Bioanalytical Chemistry</i> , 2022, 414, 6341-6353. | 3.7 | 7 |
| 6 | Multidimensional gas chromatography: Hyphenation with mass spectrometry. <i>Comprehensive Analytical Chemistry</i> , 2022, , . | 1.3 | 0 |
| 7 | Phytochemical Characterization of <i>Rhus coriaria</i> L. Extracts by Headspace Solid-Phase Micro Extraction Gas Chromatography, Comprehensive Two-Dimensional Liquid Chromatography, and Antioxidant Activity Evaluation. <i>Molecules</i> , 2022, 27, 1727. | 3.8 | 15 |
| 8 | Elucidation of Analyticalâ€“Compositional Fingerprinting of Three Different Species of Chili Pepper by Using Headspace Solid-Phase Microextraction Coupled with Gas Chromatographyâ€“Mass Spectrometry Analysis, and Sensory Profile Evaluation. <i>Molecules</i> , 2022, 27, 2355. | 3.8 | 13 |
| 9 | <i>Listeria monocytogenes</i> exposed to antimicrobial peptides displays differential regulation of lipids and proteins associated to stress response. <i>Cellular and Molecular Life Sciences</i> , 2022, 79, 263. | 5.4 | 7 |
| 10 | Supercritical fluid chromatography-tandem mass spectrometry of oxygen heterocyclic compounds in Citrus essential oils. <i>Analytical and Bioanalytical Chemistry</i> , 2022, 414, 4821-4836. | 3.7 | 4 |
| 11 | Heart-cutting and comprehensive multidimensional gas chromatography: Basic principles. <i>Comprehensive Analytical Chemistry</i> , 2022, , 69-92. | 1.3 | 2 |
| 12 | Elucidation of the Lipid Composition of Hemp (<i>Cannabis sativa</i> L.) Products by Means of Gas Chromatography and Ultra-High Performance Liquid Chromatography Coupled to Mass Spectrometry Detection. <i>Molecules</i> , 2022, 27, 3358. | 3.8 | 16 |
| 13 | Lipids in Archaeological Pottery: A Review on Their Sampling and Extraction Techniques. <i>Molecules</i> , 2022, 27, 3451. | 3.8 | 7 |
| 14 | Distribution of bioactives in entire mill chain from the drupe to the oil and wastes. <i>Natural Product Research</i> , 2021, 35, 4182-4187. | 1.8 | 12 |
| 15 | Apocarotenoids profiling in different <i>Capsicum</i> species. <i>Food Chemistry</i> , 2021, 334, 127595. | 8.2 | 24 |
| 16 | Multidimensional liquid chromatography approaches for analysis of food contaminants. <i>Journal of Separation Science</i> , 2021, 44, 17-34. | 2.5 | 15 |
| 17 | Differentiation of Italian extra virgin olive oils by rapid evaporative ionization mass spectrometry. <i>LWT - Food Science and Technology</i> , 2021, 138, 110715. | 5.2 | 11 |
| 18 | Comprehensive twoâ€“dimensional liquid chromatographyâ€“based qualitative screening of aqueous phases from pyrolysis bioâ€“oils. <i>Electrophoresis</i> , 2021, 42, 58-67. | 2.4 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Cannabis Sativa L.: a comprehensive review on the analytical methodologies for cannabinoids and terpenes characterization. <i>Journal of Chromatography A</i> , 2021, 1637, 461864. | 3.7 | 49 |
| 20 | Phytochemical Investigation and Antioxidant Activity of <i>Globularia alypum</i> L.. <i>Molecules</i> , 2021, 26, 759. | 3.8 | 26 |
| 21 | Influence of Citrus Flavor Addition in Brewing Process: Characterization of the Volatile and Non-Volatile Profile to Prevent Frauds and Adulterations. <i>Separations</i> , 2021, 8, 18. | 2.4 | 13 |
| 22 | Identification of high-value generating molecules from the wastes of tuna fishery industry by liquid chromatography and gas chromatography hyphenated techniques with automated sample preparation. <i>Journal of Separation Science</i> , 2021, 44, 1571-1580. | 2.5 | 15 |
| 23 | The retention index approach in liquid chromatography: An historical review and recent advances. <i>Journal of Chromatography A</i> , 2021, 1640, 461963. | 3.7 | 18 |
| 24 | Preliminary observations on the use of a novel low duty cycle flow modulator for comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2021, 1643, 462076. | 3.7 | 6 |
| 25 | Reversed phase versus hydrophilic interaction liquid chromatography as first dimension of comprehensive two-dimensional liquid chromatography systems for the elucidation of the polyphenolic content of food and natural products. <i>Journal of Chromatography A</i> , 2021, 1645, 462129. | 3.7 | 28 |
| 26 | Determination of multi-pesticide residues in vegetable products using a "reduced-scale" Quechers method and flow-modulated comprehensive two-dimensional gas chromatography-triple quadrupole mass spectrometry. <i>Journal of Chromatography A</i> , 2021, 1645, 462126. | 3.7 | 15 |
| 27 | Evaluation of the Level of Toxic Contaminants and Essential Molecules in the Context of the Re-Use of Tuna Fishery Industry by-Products. <i>Food Analytical Methods</i> , 2021, 14, 2161-2174. | 2.6 | 5 |
| 28 | Pattern-Type Separation of Triacylglycerols by Silver Thiolate—Non-Aqueous Reversed Phase Comprehensive Liquid Chromatography. <i>Separations</i> , 2021, 8, 88. | 2.4 | 11 |
| 29 | Use of a low-cost, lab-made Y-interface for liquid-gas chromatography coupling for the analysis of mineral oils in food samples. <i>Journal of Chromatography A</i> , 2021, 1648, 462191. | 3.7 | 6 |
| 30 | Dietary Intake of Coumarins and Furocoumarins through Citrus Beverages: A Detailed Estimation by a HPLC-MS/MS Method Combined with the Linear Retention Index System. <i>Foods</i> , 2021, 10, 1533. | 4.3 | 13 |
| 31 | Linear retention index approach applied to liquid chromatography coupled to triple quadrupole mass spectrometry to determine oxygen heterocyclic compounds at trace level in finished cosmetics. <i>Journal of Chromatography A</i> , 2021, 1649, 462183. | 3.7 | 15 |
| 32 | Interlaboratory study of a supercritical fluid chromatography method for the determination of pharmaceutical impurities: Evaluation of multi-systems reproducibility. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021, 203, 114206. | 2.8 | 14 |
| 33 | Coumarins, Psoralens and Polymethoxyflavones in Cold-pressed Citrus Essential Oils: a Review. <i>Journal of Essential Oil Research</i> , 2021, 33, 221-239. | 2.7 | 18 |
| 34 | Comparative study of the phenolic profile, antioxidant and antimicrobial activities of leaf extracts of five <i>Juniperus</i> L. (Cupressaceae) taxa growing in Turkey. <i>Natural Product Research</i> , 2020, 34, 1636-1641. | 1.8 | 25 |
| 35 | Characterization of the polyphenolic fraction of pomegranate samples by comprehensive two-dimensional liquid chromatography coupled to mass spectrometry detection. <i>Natural Product Research</i> , 2020, 34, 39-45. | 1.8 | 34 |
| 36 | Combining linear retention index and electron ionization mass spectrometry for a reliable identification in nano liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1610, 460581. | 3.7 | 17 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 37 | African baobab (<i>Adansonia digitata</i>) fruit as promising source of procyanidins. <i>European Food Research and Technology</i> , 2020, 246, 297-306. | 3.3 | 7 |
| 38 | Characterization of monoacylglycerols and diacylglycerols rich in polyunsaturated fatty acids produced by hydrolysis of <i>Mustelus mustelus</i> liver oil catalyzed by an immobilized bacterial lipase. <i>Journal of Chromatography A</i> , 2020, 1613, 460692. | 3.7 | 9 |
| 39 | Determination of free apocarotenoids and apocarotenoid esters in human colostrum. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 1335-1342. | 3.7 | 15 |
| 40 | Recent developments in the carotenoid and carotenoid derivatives chromatography-mass spectrometry analysis in food matrices. <i>TrAC - Trends in Analytical Chemistry</i> , 2020, 132, 116047. | 11.4 | 15 |
| 41 | Identification of Fatty Acid, Lipid and Polyphenol Compounds from <i>Prunus armeniaca</i> L. Kernel Extracts. <i>Foods</i> , 2020, 9, 896. | 4.3 | 9 |
| 42 | Isolation of Microalgae from Mediterranean Seawater and Production of Lipids in the Cultivated Species. <i>Foods</i> , 2020, 9, 1601. | 4.3 | 10 |
| 43 | Characterization of Phenolic Compounds, Vitamin E and Fatty Acids from Monovarietal Virgin Olive Oils of "Picholine marocaine" Cultivar. <i>Molecules</i> , 2020, 25, 5428. | 3.8 | 15 |
| 44 | Polyphenolic compounds with biological activity in guabiroba fruits (<i>Campomanesia</i>). <i>Journal of Food Science</i> , 2020, 41, 1784-1792. | 2.4 | 19 |
| 45 | Miniaturized LC in Molecular Omics. <i>Analytical Chemistry</i> , 2020, 92, 11485-11497. | 6.5 | 30 |
| 46 | Comprehensive Chemical Characterization of the <i>Pistacia vera</i> Fruits through Original NMR Quantification Methods. <i>Applied Sciences (Switzerland)</i> , 2020, 10, 5523. | 2.5 | 3 |
| 47 | Concentration of Potentially Bioactive Compounds in Italian Extra Virgin Olive Oils from Various Sources by Using LC-MS and Multivariate Data Analysis. <i>Foods</i> , 2020, 9, 1120. | 4.3 | 20 |
| 48 | Botanical and Genetic Identification Followed by Investigation of Chemical Composition and Biological Activities on the <i>Scabiosa atropurpurea</i> L. Stem from Tunisian Flora. <i>Molecules</i> , 2020, 25, 5032. | 3.8 | 15 |
| 49 | Choline-chloride and betaine-based deep eutectic solvents for green extraction of nutraceutical compounds from spent coffee ground. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020, 189, 113421. | 2.8 | 40 |
| 50 | Determination of the Metabolite Content of <i>Brassica juncea</i> Cultivars Using Comprehensive Two-Dimensional Liquid Chromatography Coupled with a Photodiode Array and Mass Spectrometry Detection. <i>Molecules</i> , 2020, 25, 1235. | 3.8 | 29 |
| 51 | Exploration of Rapid Evaporative-Ionization Mass Spectrometry as a Shotgun Approach for the Comprehensive Characterization of <i>Kigelia Africana</i> (Lam) Benth. Fruit. <i>Molecules</i> , 2020, 25, 962. | 3.8 | 14 |
| 52 | Hyphenations of 2D capillary-based LC with mass spectrometry. <i>Journal of Chromatography A</i> , 2020, 1613, 369-412. | | 1 |
| 53 | The opposite nitric oxide modulators do not lead to the opposite changes of metabolites under cadmium excess. <i>Journal of Plant Physiology</i> , 2020, 252, 153228. | 3.5 | 5 |
| 54 | Determination of the Phenol and Tocopherol Content in Italian High-Quality Extra-Virgin Olive Oils by Using LC-MS and Multivariate Data Analysis. <i>Food Analytical Methods</i> , 2020, 13, 1027-1041. | 2.6 | 28 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 55 | Towards the determination of an equivalent standard column set between cryogenic and flow-modulated comprehensive two-dimensional gas chromatography. <i>Analytica Chimica Acta</i> , 2020, 1105, 231-236. | 5.4 | 7 |
| 56 | Rapid and miniaturized qualitative and quantitative gas chromatography profiling of human blood total fatty acids. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 2327-2337. | 3.7 | 23 |
| 57 | Comprehensive two-dimensional liquid chromatography as a powerful tool for the analysis of food and food products. <i>TrAC - Trends in Analytical Chemistry</i> , 2020, 127, 115894. | 11.4 | 52 |
| 58 | Evaluation of Italian extra virgin olive oils based on the phenolic compounds composition using multivariate statistical methods. <i>European Food Research and Technology</i> , 2020, 246, 1241-1249. | 3.3 | 11 |
| 59 | Lipid profile of fish species by liquid chromatography coupled to mass spectrometry and a novel linear retention index database. <i>Journal of Separation Science</i> , 2020, 43, 1773-1780. | 2.5 | 11 |
| 60 | Evaluation of matrix effect in one-dimensional and comprehensive two-dimensional liquid chromatography for the determination of the phenolic fraction in extra virgin olive oils. <i>Journal of Separation Science</i> , 2020, 43, 1781-1789. | 2.5 | 19 |
| 61 | A lab-developed interface for liquid-gas chromatography coupling based on the use of a modified programmed-temperature-vaporizing injector. <i>Journal of Chromatography A</i> , 2020, 1622, 461096. | 3.7 | 8 |
| 62 | <i>Brassica incana</i> Ten. (Brassicaceae): Phenolic Constituents, Antioxidant and Cytotoxic Properties of the Leaf and Flowering Top Extracts. <i>Molecules</i> , 2020, 25, 1461. | 3.8 | 24 |
| 63 | Rapid evaporative ionization mass spectrometry coupled with an electrosurgical knife for the rapid identification of Mediterranean Sea species. <i>Analytical and Bioanalytical Chemistry</i> , 2019, 411, 6603-6614. | 3.7 | 16 |
| 64 | Free carotenoids and carotenoids esters composition in Spanish orange and mandarin juices from diverse varieties. <i>Food Chemistry</i> , 2019, 300, 125139. | 8.2 | 16 |
| 65 | Oxygen heterocyclic compound screening in <i>Citrus</i> essential oils by linear retention index approach applied to liquid chromatography coupled to photodiode array detector. <i>Flavour and Fragrance Journal</i> , 2019, 34, 349-364. | 2.6 | 12 |
| 66 | The Contribution of Carotenoids, Phenolic Compounds, and Flavonoids to the Antioxidative Properties of Marine Microalgae Isolated from Mediterranean Morocco. <i>Molecules</i> , 2019, 24, 4037. | 3.8 | 88 |
| 67 | Evaluation of the availability of delphinidin and cyanidin-3-O-sambubioside from <i>Hibiscus sabdariffa</i> and 6-gingerol from <i>Zingiber officinale</i> in colon using liquid chromatography and mass spectrometry detection. <i>European Food Research and Technology</i> , 2019, 245, 2425-2433. | 3.3 | 9 |
| 68 | High-performance liquid chromatography combined with electron ionization mass spectrometry: A review. <i>TrAC - Trends in Analytical Chemistry</i> , 2019, 118, 112-122. | 11.4 | 54 |
| 69 | Recent advances in the coupling of carbon dioxide-based extraction and separation techniques. <i>TrAC - Trends in Analytical Chemistry</i> , 2019, 116, 158-165. | 11.4 | 33 |
| 70 | The Phenolic Fraction of Italian Extra Virgin Olive Oils: Elucidation Through Combined Liquid Chromatography and NMR Approaches. <i>Food Analytical Methods</i> , 2019, 12, 1759-1770. | 2.6 | 38 |
| 71 | Determination of the polyphenolic fraction of <i>Pistacia vera</i> L. kernel extracts by comprehensive two-dimensional liquid chromatography coupled to mass spectrometry detection. <i>Analytical and Bioanalytical Chemistry</i> , 2019, 411, 4819-4829. | 3.7 | 30 |
| 72 | Green Extraction Approaches for Carotenoids and Esters: Characterization of Native Composition from Orange Peel. <i>Antioxidants</i> , 2019, 8, 613. | 5.1 | 37 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 73 | Nitric oxide affects cadmium-induced changes in the lichen <i>Ramalina farinacea</i> . <i>Nitric Oxide - Biology and Chemistry</i> , 2019, 83, 11-18. | 2.7 | 30 |
| 74 | Characterization of peel and pulp proanthocyanidins and carotenoids during ripening in persimmon 'Kaki Tipo' cv, cultivated in Italy. <i>Food Research International</i> , 2019, 120, 800-809. | 6.2 | 21 |
| 75 | Use of an 'Intelligent Knife' (iknife), Based on the Rapid Evaporative Ionization Mass Spectrometry Technology, for Authenticity Assessment of Pistachio Samples. <i>Food Analytical Methods</i> , 2019, 12, 558-568. | 2.6 | 32 |
| 76 | Comprehensive two-dimensional gas chromatography-mass spectrometry using milder electron ionization conditions: A preliminary evaluation. <i>Journal of Chromatography A</i> , 2019, 1589, 134-140. | 3.7 | 15 |
| 77 | On-line liquid chromatography-comprehensive two dimensional gas chromatography with dual detection for the analysis of mineral oil and synthetic hydrocarbons in cosmetic lip care products. <i>Antonica Chimica Acta</i> , 2019, 1048, 221-226. | 5.4 | 14 |
| 78 | Comprehensive lipid profiling in the Mediterranean mussel (<i>Mytilus galloprovincialis</i>) using hyphenated and multidimensional chromatography techniques coupled to mass spectrometry detection. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 3297-3313. | 3.7 | 35 |
| 79 | Use of an Online Extraction Technique Coupled to Liquid Chromatography for Determination of Caffeine in Coffee, Tea, and Cocoa. <i>Food Analytical Methods</i> , 2018, 11, 2637-2644. | 2.6 | 17 |
| 80 | Proposal of a Linear Retention Index System for Improving Identification Reliability of Triacylglycerol Profiles in Lipid Samples by Liquid Chromatography Methods. <i>Analytical Chemistry</i> , 2018, 90, 3313-3320. | 6.5 | 31 |
| 81 | Untargeted profiling of <i>Glycyrrhiza glabra</i> extract with comprehensive two-dimensional liquid chromatography-mass spectrometry using multi-segmented shift gradients in the second dimension: Expanding the metabolic coverage. <i>Electrophoresis</i> , 2018, 39, 1993-2000. | 2.4 | 27 |
| 82 | Analysis of phenolic compounds in different parts of pomegranate (<i>Punica granatum</i>) fruit by HPLC-PDA-ESI/MS and evaluation of their antioxidant activity: application to different Italian varieties. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 3507-3520. | 3.7 | 111 |
| 83 | Accumulation and toxicity of organochlorines in green microalgae. <i>Journal of Hazardous Materials</i> , 2018, 347, 168-175. | 12.4 | 28 |
| 84 | Partial characterization of the pigments produced by the marine-derived fungus <i>Talaromyces albobiverticillius</i> 30548. Towards a new fungal red colorant for the food industry. <i>Journal of Food Composition and Analysis</i> , 2018, 67, 38-47. | 3.9 | 53 |
| 85 | Recent Analytical Techniques Advances in the Carotenoids and Their Derivatives Determination in Various Matrixes. <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 3302-3307. | 5.2 | 33 |
| 86 | Comparison of different analytical techniques for the analysis of carotenoids in tamarillo (<i>Solanum</i>) | 3.0 | 42 |
| 87 | Multilevel characterization of marine microbial biodegradation potentiality by means of flow-modulated comprehensive two-dimensional gas chromatography combined with a triple quadrupole mass spectrometer. <i>Journal of Chromatography A</i> , 2018, 1547, 99-106. | 3.7 | 9 |
| 88 | Authentication of citrus volatiles based on carbon isotope ratios. <i>Journal of Essential Oil Research</i> , 2018, 30, 1-15. | 2.7 | 21 |
| 89 | Novel comprehensive multidimensional liquid chromatography approach for elucidation of the microbiosphere of shikimate-producing <i>Escherichia coli</i> SP1.1/pKD15.071 strain. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 3473-3482. | 3.7 | 8 |
| 90 | Monoacylglycerol and diacylglycerol production by hydrolysis of refined vegetable oil by-products using an immobilized lipase from <i>Serratia</i> sp. W3. <i>Journal of Separation Science</i> , 2018, 41, 4323-4330. | 2.5 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 91 | Metabolic responses of <i>Ulva compressa</i> to single and combined heavy metals. <i>Chemosphere</i> , 2018, 213, 384-394. | 8.2 | 18 |
| 92 | Carotenoids and apocarotenoids determination in intact human blood samples by online supercritical fluid extraction-supercritical fluid chromatography-tandem mass spectrometry. <i>Analytica Chimica Acta</i> , 2018, 1032, 40-47. | 5.4 | 39 |
| 93 | Comprehensive Two-Dimensional Liquid Chromatography Coupled to Mass Spectrometry. <i>Comprehensive Analytical Chemistry</i> , 2018, 79, 81-123. | 1.3 | 3 |
| 94 | Supercritical Fluid Chromatography— Ultra-High Pressure Liquid Chromatography for Red Chili Pepper Fingerprinting by Photodiode Array, Quadrupole-Time-of-Flight and Ion Mobility Mass Spectrometry (SFC— RP-UHPLC-PDA-Q-ToF MS-IMS). <i>Food Analytical Methods</i> , 2018, 11, 3331-3341. | 2.6 | 20 |
| 95 | Apocarotenoids determination in <i>Capsicum chinense</i> Jacq. cv. Habanero, by supercritical fluid chromatography-triple-quadrupole/mass spectrometry. <i>Food Chemistry</i> , 2017, 231, 316-323. | 8.2 | 48 |
| 96 | Highly informative multiclass profiling of lipids by ultra-high performance liquid chromatography— Low resolution (quadrupole) mass spectrometry by using electrospray ionization and atmospheric pressure chemical ionization interfaces. <i>Journal of Chromatography A</i> , 2017, 1509, 69-82. | 3.7 | 18 |
| 97 | Flow-modulated comprehensive two-dimensional gas chromatography combined with a vacuum ultraviolet detector for the analysis of complex mixtures. <i>Journal of Chromatography A</i> , 2017, 1497, 135-143. | 3.7 | 42 |
| 98 | Quali-quantitative characterization of the volatile constituents in <i>Cordia verbenacea</i> D.C. essential oil exploiting advanced chromatographic approaches and nuclear magnetic resonance analysis. <i>Journal of Chromatography A</i> , 2017, 1524, 246-253. | 3.7 | 18 |
| 99 | Determination of amines and phenolic acids in wine with benzoyl chloride derivatization and liquid chromatography—mass spectrometry. <i>Journal of Chromatography A</i> , 2017, 1523, 248-256. | 3.7 | 24 |
| 100 | Direct online extraction and determination by supercritical fluid extraction with chromatography and mass spectrometry of targeted carotenoids from red Habanero peppers (<i>Capsicum chinense</i>) | 0.0 | 0 |
| 101 | Comprehensive two-dimensional liquid chromatography. , 2017, , 403-415. | | 2 |
| 102 | Multidimensional liquid chromatography in food analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2017, 96, 116-123. | 11.4 | 59 |
| 103 | Comprehensive Liquid Chromatography and Other Liquid-Based Comprehensive Techniques Coupled to Mass Spectrometry in Food Analysis. <i>Analytical Chemistry</i> , 2017, 89, 414-429. | 6.5 | 46 |
| 104 | Supercritical fluid chromatography for lipid analysis in foodstuffs. <i>Journal of Separation Science</i> , 2017, 40, 361-382. | 2.5 | 32 |
| 105 | Comprehensive two-dimensional liquid chromatography for polyphenol analysis in foodstuffs. <i>Journal of Separation Science</i> , 2017, 40, 7-24. | 2.5 | 48 |
| 106 | Recent Advances in Comprehensive Two-Dimensional Liquid Chromatography for the Analysis of Natural Products. , 2017, , 287-307. | | 1 |
| 107 | Analysis of lipid profile in lipid storage myopathy. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016, 1029-1030, 157-168. | 2.3 | 6 |
| 108 | Rapid isolation, reliable characterization, and water solubility improvement of polymethoxyflavones from cold-pressed mandarin essential oil. <i>Journal of Separation Science</i> , 2016, 39, 2018-2027. | 2.5 | 20 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Free fatty acid profiling of marine sentinels by nanoLC-EI-MS for the assessment of environmental pollution effects. <i>Science of the Total Environment</i> , 2016, 571, 955-962. | 8.0 | 45 |
| 110 | Characterization of the pigment fraction in sweet bell peppers (<i>Capsicum annuum</i> L.) harvested at green and overripe yellow and red stages by offline multidimensional convergence chromatography/liquid chromatography–mass spectrometry. <i>Journal of Separation Science</i> , 2016, 39, 3281-3291. | 2.5 | 30 |
| 111 | Comprehensive two-dimensional liquid chromatography–tandem mass spectrometry for the simultaneous determination of wine polyphenols and target contaminants. <i>Journal of Chromatography A</i> , 2016, 1458, 54-62. | 3.7 | 69 |
| 112 | Nano Liquid Chromatography Directly Coupled to Electron Ionization Mass Spectrometry for Free Fatty Acid Elucidation in Mussel. <i>Analytical Chemistry</i> , 2016, 88, 4021-4028. | 6.5 | 60 |
| 113 | Chemical characterisation of old cabbage (<i>Brassica oleracea</i> L. var. <i>acephala</i>) seed oil by liquid chromatography and different spectroscopic detection systems. <i>Natural Product Research</i> , 2016, 30, 1646-1654. | 1.8 | 22 |
| 114 | Application of Comprehensive Two-Dimensional Liquid Chromatography for Carotenoid Analysis in Red Mamey (<i>Pouteria sapote</i>) Fruit. <i>Food Analytical Methods</i> , 2016, 9, 2335-2341. | 2.6 | 33 |
| 115 | Bergamot (<i>Citrus bergamia</i> Risso) as a source of nutraceuticals: Limonoids and flavonoids. <i>Journal of Functional Foods</i> , 2016, 20, 10-19. | 3.4 | 62 |
| 116 | Role of the flavonoid-rich fraction in the antioxidant and cytotoxic activities of <i>Bauhinia forficata</i> Link. (Fabaceae) leaves extract. <i>Natural Product Research</i> , 2016, 30, 1229-1239. | 1.8 | 40 |
| 117 | Multidimensional preparative liquid chromatography to isolate flavonoids from bergamot juice and evaluation of their anti-inflammatory potential. <i>Journal of Separation Science</i> , 2015, 38, 4196-4203. | 2.5 | 9 |
| 118 | Lipidomics. <i>Comprehensive Analytical Chemistry</i> , 2015, 68, 395-439. | 1.3 | 4 |
| 119 | Determination of the triacylglycerol fraction in fish oil by comprehensive liquid chromatography techniques with the support of gas chromatography and mass spectrometry data. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 5211-5225. | 3.7 | 36 |
| 120 | Reduced time HPLC analyses for fast quality control of citrus essential oils. <i>Journal of Essential Oil Research</i> , 2015, 27, 307-315. | 2.7 | 32 |
| 121 | Sample preparation techniques coupled to advanced chromatographic methods for marine organisms investigation. <i>Analytica Chimica Acta</i> , 2015, 875, 41-53. | 5.4 | 25 |
| 122 | Underestimated sources of flavonoids, limonoids and dietary fiber: Availability in orange's by-products. <i>Journal of Functional Foods</i> , 2015, 12, 150-157. | 3.4 | 53 |
| 123 | Analysis of human plasma lipids by using comprehensive two-dimensional gas chromatography with dual detection and with the support of high-resolution time-of-flight mass spectrometry for structural elucidation. <i>Journal of Separation Science</i> , 2015, 38, 267-275. | 2.5 | 18 |
| 124 | Complementary Analytical Liquid Chromatography Methods for the Characterization of Aqueous Phase from Pyrolysis of Lignocellulosic Biomasses. <i>Analytical Chemistry</i> , 2014, 86, 11255-11262. | 6.5 | 51 |
| 125 | Flow-modulation low-pressure comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2014, 1372, 236-244. | 3.7 | 44 |
| 126 | Thorough investigation of the oxygen heterocyclic fraction of lime (<i>Citrus aurantifolia</i>) | 2.5 | 13 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 127 | Underestimated sources of flavonoids, limonoids and dietary fibre: Availability in lemon's by-products. <i>Journal of Functional Foods</i> , 2014, 9, 18-26. | 3.4 | 71 |
| 128 | Characterisation of lipid fraction of marine macroalgae by means of chromatography techniques coupled to mass spectrometry. <i>Food Chemistry</i> , 2014, 145, 932-940. | 8.2 | 55 |
| 129 | NMR characterisation and dynamic behaviour of [Pt(bipy)(R-Thiourea) ₂]Cl ₂ and [Pt(phen)(R-Thiourea) ₂]Cl ₂ complexes. <i>Inorganica Chimica Acta</i> , 2014, 410, 1-10. | 2.4 | 11 |
| 130 | High performance characterization of triacylglycerols in milk and milk-related samples by liquid chromatography and mass spectrometry. <i>Journal of Chromatography A</i> , 2014, 1360, 172-187. | 3.7 | 54 |
| 131 | Use of greatly-reduced gas flows in flow-modulated comprehensive two-dimensional gas chromatography-mass spectrometry. <i>Journal of Chromatography A</i> , 2014, 1359, 271-276. | 3.7 | 48 |
| 132 | Continuous vs. segmented second-dimension system gradients for comprehensive two-dimensional liquid chromatography of sugarcane (<i>Saccharum</i> spp.). <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 4315-4324. | 3.7 | 33 |
| 133 | Potential of comprehensive chromatography in food analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2013, 52, 186-205. | 11.4 | 91 |
| 134 | Gas velocity at the point of re-injection: An additional parameter in comprehensive two-dimensional gas chromatography optimization. <i>Journal of Chromatography A</i> , 2013, 1314, 216-223. | 3.7 | 17 |
| 135 | Stop-flow comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection for phospholipid analysis. <i>Journal of Chromatography A</i> , 2013, 1278, 46-53. | 3.7 | 69 |
| 136 | Native carotenoids composition of some tropical fruits. <i>Food Chemistry</i> , 2013, 140, 825-836. | 8.2 | 85 |
| 137 | A new HPLC method developed for the analysis of oxygen heterocyclic compounds in <i>Citrus</i> essential oils. <i>Journal of Essential Oil Research</i> , 2012, 24, 119-129. | 2.7 | 31 |
| 138 | Mass spectrometric elucidation of triacylglycerol content of <i>Brevoortia tyrannus</i> (menhaden) oil using non-aqueous reversed-phase liquid chromatography under ultra high pressure conditions. <i>Journal of Chromatography A</i> , 2012, 1259, 227-236. | 3.7 | 34 |
| 139 | A flow-modulated comprehensive gas chromatography-mass spectrometry method for the analysis of fatty acid profiles in marine and biological samples. <i>Journal of Chromatography A</i> , 2012, 1255, 171-176. | 3.7 | 31 |
| 140 | Ultra high pressure in the second dimension of a comprehensive two-dimensional liquid chromatographic system for carotenoid separation in red chili peppers. <i>Journal of Chromatography A</i> , 2012, 1255, 244-251. | 3.7 | 63 |
| 141 | Use of ionic liquids as stationary phases in hyphenated gas chromatography techniques. <i>Journal of Chromatography A</i> , 2012, 1255, 130-144. | 3.7 | 94 |
| 142 | Mass spectrometry detection in comprehensive liquid chromatography: Basic concepts, instrumental aspects, applications and trends. <i>Mass Spectrometry Reviews</i> , 2012, 31, 523-559. | 5.4 | 86 |
| 143 | Evaluation of a Medium-Polarity Ionic Liquid Stationary Phase in the Analysis of Flavor and Fragrance Compounds. <i>Analytical Chemistry</i> , 2011, 83, 7947-7954. | 6.5 | 77 |
| 144 | Online Comprehensive RPLC-MS—RPLC with Mass Spectrometry Detection for the Analysis of Proteome Samples. <i>Analytical Chemistry</i> , 2011, 83, 2485-2491. | 6.5 | 60 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Chemical Characterization of Sacha Inchi (<i>Plukenetia volubilis</i> L.) Oil. Journal of Agricultural and Food Chemistry, 2011, 59, 13043-13049. | 5.2 | 111 |
| 146 | Determination of phospholipids in milk samples by means of hydrophilic interaction liquid chromatography coupled to evaporative light scattering and mass spectrometry detection. Journal of Chromatography A, 2011, 1218, 6476-6482. | 3.7 | 110 |
| 147 | Comprehensive two-dimensional liquid chromatography with evaporative light scattering detection for the analysis of triacylglycerols in <i>Borago officinalis</i> . Journal of Separation Science, 2011, 34, 688-692. | 2.5 | 24 |
| 148 | Determination of flavanones in <i>Citrus</i> juices by means of one- and two-dimensional liquid chromatography. Journal of Separation Science, 2011, 34, 681-687. | 2.5 | 46 |
| 149 | Analytical characterization of mandarin (<i>Citrus deliciosa</i> Ten.) essential oil. Flavour and Fragrance Journal, 2011, 26, 34-46. | 2.6 | 28 |
| 150 | Employing ultra high pressure liquid chromatography as the second dimension in a comprehensive two-dimensional system for analysis of <i>Stevia rebaudiana</i> extracts. Journal of Chromatography A, 2011, 1218, 2012-2018. | 3.7 | 90 |
| 151 | A flexible loop-type flow modulator for comprehensive two-dimensional gas chromatography. Journal of Chromatography A, 2011, 1218, 3140-3145. | 3.7 | 35 |
| 152 | Study on the chemical composition variability of some processed bergamot (<i>Citrus bergamia</i>) essential oils. Flavour and Fragrance Journal, 2010, 25, 4-12. | 2.6 | 53 |
| 153 | RP-LC-AP-MS analysis of a tryptic digest using a combination of totally porous and partially porous stationary phases. Journal of Separation Science, 2010, 33, 1454-1461. | 2.5 | 38 |
| 154 | High peak capacity separation of peptides through the serial connection of LC shell-packed columns. Journal of Separation Science, 2009, 32, 1129-1136. | 2.5 | 34 |
| 155 | Comprehensive two-dimensional liquid chromatography to quantify polyphenols in red wines. Journal of Chromatography A, 2009, 1216, 7483-7487. | 3.7 | 74 |
| 156 | High efficiency liquid chromatography techniques coupled to mass spectrometry for the characterization of mate extracts. Journal of Chromatography A, 2009, 1216, 7213-7221. | 3.7 | 89 |
| 157 | Quantitative Characterization of Solid Epoxy Resins Using Comprehensive Two Dimensional Liquid Chromatography Coupled with Electrospray Ionization-Time of Flight Mass Spectrometry. Analytical Chemistry, 2009, 81, 4271-4279. | 6.5 | 36 |
| 158 | Determination of Oxygen Heterocyclic Components in Citrus Products by HPLC with UV Detection. Journal of Agricultural and Food Chemistry, 2009, 57, 6543-6551. | 5.2 | 57 |
| 159 | Evaluation of Use of a Dicationic Liquid Stationary Phase in the Fast and Conventional Gas Chromatographic Analysis of Health-Hazardous C ₁₈ Cis/Trans Fatty Acids. Analytical Chemistry, 2009, 81, 5561-5568. | 6.5 | 67 |
| 160 | Elucidation of fatty acid profiles in vegetable oils exploiting group-type patterning and enhanced sensitivity of comprehensive two-dimensional gas chromatography. Journal of Separation Science, 2008, 31, 1797-1802. | 2.5 | 32 |
| 161 | Use of partially porous column as second dimension in comprehensive two-dimensional system for analysis of polyphenolic antioxidants. Journal of Separation Science, 2008, 31, 3297-3308. | 2.5 | 72 |
| 162 | Linear retention indices in gas chromatographic analysis: a review. Flavour and Fragrance Journal, 2008, 23, 297-314. | 2.6 | 192 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 163 | Reliable identification of pesticides using linear retention indices as an active tool in gas chromatographic mass spectrometric analysis. <i>Journal of Chromatography A</i> , 2008, 1186, 430-433. | 3.7 | 15 |
| 164 | Serial coupled columns reversed-phase separations in high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2008, 1188, 208-215. | 3.7 | 45 |
| 165 | Quantification in Comprehensive Two-Dimensional Liquid Chromatography. <i>Analytical Chemistry</i> , 2008, 80, 5418-5424. | 6.5 | 53 |
| 166 | Comprehensive chromatographic methods for the analysis of lipids. <i>TrAC - Trends in Analytical Chemistry</i> , 2007, 26, 191-205. | 11.4 | 73 |
| 167 | Comprehensive gas chromatography coupled to mass spectrometry for the separation of pesticides in a very complex matrix. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 389, 1755-1763. | 3.7 | 39 |
| 168 | Reliable characterization of coffee bean aroma profiles by automated headspace solid phase microextraction-gas chromatography-mass spectrometry with the support of a dual-filter mass spectra library. <i>Journal of Separation Science</i> , 2005, 28, 1101-1109. | 2.5 | 80 |
| 169 | On the genuineness of citrus essential oils. Part LVII. The composition of distilled lime oil. <i>Flavour and Fragrance Journal</i> , 1998, 13, 93-97. | 2.6 | 14 |
| 170 | Characterization of Cold-Pressed Key and Persian Lime Oils by Gas Chromatography, Gas Chromatography/Mass Spectroscopy, High-Performance Liquid Chromatography, and Physicochemical Indices. <i>Journal of Agricultural and Food Chemistry</i> , 1997, 45, 3608-3616. | 5.2 | 39 |
| 171 | On the genuineness of citrus essential oils. Part XLVI. Polymethoxylated flavones of the non-volatile residue of Italian sweet orange and mandarin essential oils. <i>Flavour and Fragrance Journal</i> , 1994, 9, 105-111. | 2.6 | 18 |
| 172 | Analytical evaluation of carotenoids, apocarotenoids, capsaicinoids, and phenolics to assess the effect of a protective treatment on chili peppers dried at different temperatures. <i>European Food Research and Technology</i> , 0, , . | 3.3 | 2 |
| 173 | Sustainability in (bio-)analytical chemistry. <i>Analytical and Bioanalytical Chemistry</i> , 0, , . | 3.7 | 0 |