
Manuel L Lemos

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1419963/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Isolation of Vibrio alginolyticus and Vibrio splendidus from Aquacultured Carpet Shell Clam () Tj ETQq1 1 0.78431 Microbiology, 2005, 71, 98-104.	4 rgBT / 3.1	Overlock 10 244
2	Antibiotic activity of epiphytic bacteria isolated from intertidal seaweeds. Microbial Ecology, 1985, 11, 149-163.	2.8	194
3	Photobacterium damselae subsp. damselae, a bacterium pathogenic for marine animals and humans. Frontiers in Microbiology, 2013, 4, 283.	3.5	169
4	Inhibitory activity of antibioticâ€producing marine bacteria against fish pathogens. Journal of Applied Bacteriology, 1988, 65, 97-101.	1.1	146
5	Chromosome-mediated iron uptake system in pathogenic strains of Vibrio anguillarum. Journal of Bacteriology, 1988, 170, 1920-1925.	2.2	120
6	Modified Medium for the Oxidation-Fermentation Test in the Identification of Marine Bacteria. Applied and Environmental Microbiology, 1985, 49, 1541-1543.	3.1	117
7	Iron uptake by Pasteurella piscicida and its role in pathogenicity for fish. Applied and Environmental Microbiology, 1994, 60, 2990-2998.	3.1	84
8	Gene Expression Profiles of the Spleen, Liver, and Head Kidney in Turbot (Scophthalmus maximus) Along the Infection Process with Aeromonas salmonicida Using an Immune-Enriched Oligo-microarray. Marine Biotechnology, 2011, 13, 1099-1114.	2.4	79
9	Homology of Vibrio anguillarum strains causing epizootics in turbot, salmon and trout reared on the Atlantic coast of Spain. Aquaculture, 1987, 67, 41-52.	3.5	77
10	The Photobacterium damselae subsp. damselae Hemolysins Damselysin and HlyA Are Encoded within a New Virulence Plasmid. Infection and Immunity, 2011, 79, 4617-4627.	2.2	73
11	Characterization of Heme Uptake Cluster Genes in the Fish Pathogen Vibrio anguillarum. Journal of Bacteriology, 2004, 186, 6159-6167.	2.2	71
12	Integrating Conjugative Elements as Vectors of Antibiotic, Mercury, and Quaternary Ammonium Compound Resistance in Marine Aquaculture Environments. Antimicrobial Agents and Chemotherapy, 2012, 56, 2619-2626.	3.2	69
13	Two tonB Systems Function in Iron Transport in Vibrio anguillarum , but Only One Is Essential for Virulence. Infection and Immunity, 2004, 72, 7326-7329.	2.2	62
14	Expressed sequence tags (ESTs) from immune tissues of turbot (Scophthalmus maximus) challenged with pathogens. BMC Veterinary Research, 2008, 4, 37.	1.9	61
15	Structural characterization of vanchrobactin, a new catechol siderophore produced by the fish pathogen Vibrio anguillarum serotype O2. Tetrahedron Letters, 2006, 47, 7113-7116.	1.4	60
16	Synergistic and Additive Effects of Chromosomal and Plasmid-Encoded Hemolysins Contribute to Hemolysis and Virulence in Photobacterium damselae subsp. damselae. Infection and Immunity, 2013, 81, 3287-3299.	2.2	60
17	Genomic and Functional Analysis of ICE <i>Pda</i> Spa1, a Fish-Pathogen-Derived SXT-Related Integrating Conjugative Element That Can Mobilize a Virulence Plasmid. Journal of Bacteriology, 2008, 190, 3353-3361.	2.2	58
18	Population dynamics of heterotrophic bacterial communities associated withFucus vesiculosus andUlva rigida in an estuary. Microbial Ecology, 1988, 15, 345-357.	2.8	55

#	Article	IF	CITATIONS
19	The Expression of Virulence Factors in Vibrio anguillarum Is Dually Regulated by Iron Levels and Temperature. Frontiers in Microbiology, 2019, 10, 2335.	3.5	54
20	Plasmid coding for transferable drug resistance in bacteria isolated from cultured rainbow trout. Applied and Environmental Microbiology, 1984, 48, 872-877.	3.1	53
21	Structure and Biosynthetic Assembly of Piscibactin, a Siderophore from <i>Photobacterium damselae</i> subsp. <i>piscicida</i> , Predicted from Genome Analysis. European Journal of Organic Chemistry, 2012, 2012, 5693-5700.	2.4	49
22	A Transmissible Plasmid-Borne Pathogenicity Island Confers Piscibactin Biosynthesis in the Fish Pathogen Photobacterium damselae subsp. piscicida. Applied and Environmental Microbiology, 2015, 81, 5867-5879.	3.1	48
23	Revisiting the genus Photobacterium: taxonomy, ecology and pathogenesis. International Microbiology, 2017, 20, 1-10.	2.4	47
24	A siderophore biosynthesis gene cluster from the fish pathogen Photobacterium damselae subsp. piscicida is structurally and functionally related to the Yersinia high-pathogenicity island. Microbiology (United Kingdom), 2006, 152, 3327-3341.	1.8	46
25	A gene cluster involved in the biosynthesis of vanchrobactin, a chromosome-encoded siderophore produced by Vibrio anguillarum. Microbiology (United Kingdom), 2006, 152, 3517-3528.	1.8	45
26	Chromosome-Encoded Hemolysin, Phospholipase, and Collagenase in Plasmidless Isolates of Photobacterium damselae subsp. damselae Contribute to Virulence for Fish. Applied and Environmental Microbiology, 2017, 83, .	3.1	44
27	Distribution of plasmid- and chromosome-mediated iron uptake systems in Vibrio anguillarum strains of different origins. Applied and Environmental Microbiology, 1991, 57, 2956-2962.	3.1	44
28	Photobacterium damselae subsp. damselae Major Virulence Factors Dly, Plasmid-Encoded HlyA, and Chromosome-Encoded HlyA Are Secreted via the Type II Secretion System. Infection and Immunity, 2015, 83, 1246-1256.	2.2	42
29	Anguibactin―versus vanchrobactinâ€mediated iron uptake in <i>Vibrio anguillarum</i> : evolution and ecology of a fish pathogen. Environmental Microbiology Reports, 2010, 2, 19-26.	2.4	41
30	Iron-binding proteins and heme compounds as iron sources forVibrio anguillarum. Current Microbiology, 1991, 23, 221-226.	2.2	40
31	Phobalysin, a Small β-Pore-Forming Toxin of Photobacterium damselae subsp. damselae. Infection and Immunity, 2015, 83, 4335-4348.	2.2	40
32	The Siderophore Piscibactin Is a Relevant Virulence Factor for Vibrio anguillarum Favored at Low Temperatures. Frontiers in Microbiology, 2018, 9, 1766.	3.5	40
33	Integrating conjugative elements of the SXT/R391 family from fish-isolated <i>Vibrios</i> encode restriction-modification systems that confer resistance to bacteriophages. FEMS Microbiology Ecology, 2013, 83, 457-467.	2.7	39
34	Two Catechol Siderophores, Acinetobactin and Amonabactin, Are Simultaneously Produced by <i>Aeromonas salmonicida</i> subsp. <i>salmonicida</i> Sharing Part of the Biosynthetic Pathway. ACS Chemical Biology, 2015, 10, 2850-2860.	3.4	38
35	Synthesis and antibacterial activity of conjugates between norfloxacin and analogues of the siderophore vanchrobactin. Bioorganic and Medicinal Chemistry, 2013, 21, 295-302.	3.0	36
36	Identification of heme uptake genes in the fish pathogen Aeromonas salmonicida subsp. salmonicida. Archives of Microbiology, 2008, 190, 439-449.	2.2	35

#	Article	IF	CITATIONS
37	Competitive dominance of antibioticâ€producing marine bacteria in mixed cultures. Journal of Applied Bacteriology, 1991, 71, 228-232.	1.1	34
38	Iron Uptake Mechanisms in the Fish Pathogen Tenacibaculum maritimum. Applied and Environmental Microbiology, 2005, 71, 6947-6953.	3.1	34
39	Identification of Siderophore Biosynthesis Genes Essential for Growth of <i>Aeromonas salmonicida</i> under Iron Limitation Conditions. Applied and Environmental Microbiology, 2008, 74, 2341-2348.	3.1	33
40	Utilization of hemin and hemoglobin by Vibrio vulnificus biotype 2. Applied and Environmental Microbiology, 1996, 62, 2806-2810.	3.1	31
41	Purification and characterization of an antibacterial substance produced by a marine Alteromonas species. Antimicrobial Agents and Chemotherapy, 1989, 33, 1674-1679.	3.2	30
42	Biosynthetic and regulatory elements involved in the production of the siderophore vanchrobactin in Vibrio anguillarum. Microbiology (United Kingdom), 2008, 154, 1400-1413.	1.8	30
43	Iron-binding compounds and related outer membrane proteins in Vibrio cholerae non-O1 strains from aquatic environments. Applied and Environmental Microbiology, 1990, 56, 2410-2416.	3.1	30
44	Heme, an iron supply for vibrios pathogenic for fish. BioMetals, 2007, 20, 615-626.	4.1	29
45	Serological Relationships amongVibrio anguillarumStrains. Journal of Aquatic Animal Health, 1990, 2, 21-29.	1.4	27
46	Identification of heme-binding proteins in the cell membranes ofVibrio anguillarum. FEMS Microbiology Letters, 1996, 135, 265-270.	1.8	27
47	Isolation of mutants of Vibrio anguillarum defective in haeme utilisation and cloning of huvA, a gene coding for an outer membrane protein involved in the use of haeme as iron source. Archives of Microbiology, 2003, 179, 329-338.	2.2	26
48	Subtractive hybridization reveals a high genetic diversity in the fish pathogen Photobacterium damselae subsp. piscicida: evidence of a SXT-like element. Microbiology (United Kingdom), 2005, 151, 2659-2669.	1.8	26
49	FvtA Is the Receptor for the Siderophore Vanchrobactin in <i>Vibrio anguillarum</i> : Utility as a Route of Entry for Vanchrobactin Analogues. Applied and Environmental Microbiology, 2009, 75, 2775-2783.	3.1	26
50	Evidence for horizontal gene transfer, gene duplication and genetic variation as driving forces of the diversity of haemolytic phenotypes inPhotobacterium damselaesubsp.damselae. FEMS Microbiology Letters, 2014, 355, 152-162.	1.8	26
51	A proteomic analysis of the iron response of Photobacterium damselae subsp. damselae reveals metabolic adaptations to iron levels changes and novel potential virulence factors. Veterinary Microbiology, 2017, 201, 257-264.	1.9	26
52	Iron uptake mechanisms as key virulence factors in bacterial fish pathogens. Journal of Applied Microbiology, 2020, 129, 104-115.	3.1	24
53	Heme uptake genes in human and fish isolates of Photobacterium damselae: existence of hutA pseudogenes. Archives of Microbiology, 2005, 183, 347-358.	2.2	23
54	Binding of haemin by the fish pathogen Photobacterium damselae subsp. piscicida. Diseases of Aquatic Organisms, 2002, 48, 109-115.	1.0	23

#	Article	IF	CITATIONS
55	Identification of iron regulated genes in the fish pathogen Aeromonas salmonicida subsp. salmonicida: Genetic diversity and evidence of conserved iron uptake systems. Veterinary Microbiology, 2009, 133, 377-382.	1.9	22
56	Secreted Citrate Serves as Iron Carrier for the Marine Pathogen Photobacterium damselae subsp damselae. Frontiers in Cellular and Infection Microbiology, 2017, 7, 361.	3.9	22
57	Transcriptional organization and regulation of the Vibrio anguillarum heme uptake gene cluster. Gene, 2006, 374, 68-76.	2.2	19
58	Siderophore production by environmental strains ofSalmonellaspecies. FEMS Microbiology Letters, 1989, 57, 7-12.	1.8	18
59	Host-Nonspecific Iron Acquisition Systems and Virulence in the Zoonotic Serovar of Vibrio vulnificus. Infection and Immunity, 2014, 82, 731-744.	2.2	17
60	Genetic Variability of the Heme Uptake System among Different Strains of the Fish Pathogen Vibrio anguillarum : Identification of a New Heme Receptor. Applied and Environmental Microbiology, 2005, 71, 8434-8441.	3.1	15
61	Genetic characterization of pAsa6, a new plasmid from Aeromonas salmonicida subsp. salmonicida that encodes a type III effector protein AopH homolog. Plasmid, 2009, 61, 176-181.	1.4	15
62	Genomic analysis of the marine fish pathogen Photobacterium damselae subsp. piscicida: Insertion sequences proliferation is associated with chromosomal reorganisations and rampant gene decay. Infection, Genetics and Evolution, 2017, 54, 221-229.	2.3	15
63	The Fish Pathogen Vibrio ordalii Under Iron Deprivation Produces the Siderophore Piscibactin. Microorganisms, 2019, 7, 313.	3.6	15
64	Identification of Fur regulated genes in the bacterial fish pathogen Photobacterium damselae ssp. piscicida using the Fur titration assay. BioMetals, 2004, 17, 725-733.	4.1	14
65	Unveiling the pan-genome of the SXT/R391 family of ICEs: molecular characterisation of new variable regions of SXT/R391-like ICEs detected in Pseudoalteromonas sp. and Vibrio scophthalmi. Antonie Van Leeuwenhoek, 2016, 109, 1141-1152.	1.7	14
66	Outer membrane protein FrpA, the siderophore piscibactin receptor of Photobacterium damselae subsp. piscicida, as a subunit vaccine against photobacteriosis in sole (Solea senegalensis). Fish and Shellfish Immunology, 2019, 94, 723-729.	3.6	13
67	Isolation of heme-binding proteins fromVibrio anguillarumusing affinity chromatography. FEMS Microbiology Letters, 1996, 141, 19-23.	1.8	12
68	Distribution of small plasmids in <i>Aeromonas salmonicida</i> subsp. <i>salmonicida</i> strains isolated from NW Spain and Portugal: evidence of clonality in strains isolated from turbot, <i>Psetta maxima</i> (L.). Journal of Fish Diseases, 2008, 31, 469-472.	1.9	12
69	Synthesis and biological activity of analogues of vanchrobactin, a siderophore from Vibrio anguillarum serotype O2. Organic and Biomolecular Chemistry, 2008, 6, 1278.	2.8	12
70	ldentification of the Ferric-Acinetobactin Outer Membrane Receptor in <i>Aeromonas salmonicida</i> subsp. <i>salmonicida</i> and Structure–Activity Relationships of Synthetic Acinetobactin Analogues. ACS Chemical Biology, 2017, 12, 479-493.	3.4	12
71	Virulence properties of three new <i>Photobacterium</i> species affecting cultured fish. Journal of Applied Microbiology, 2020, 129, 37-50.	3.1	12
72	Ferric-reductase activities in whole cells and cell fractions of Vibrio (Listonella) anguillarum. Microbiology (United Kingdom), 1996, 142, 3187-3193.	1.8	12

#	Article	IF	CITATIONS
73	Ferric-reductase activities inVibrio vulnificusbiotypes 1 and 2. FEMS Microbiology Letters, 1999, 172, 205-211.	1.8	10
74	Presence of high-affinity iron uptake systems in fish-isolated and environmental strains of Vibrio anguillarum serotype O3. FEMS Microbiology Letters, 2001, 202, 79-83.	1.8	10
75	Presence of high-affinity iron uptake systems in fish-isolated and environmental strains ofVibrio anguillarumserotype O3. FEMS Microbiology Letters, 2001, 202, 79-83.	1.8	10
76	Isolation of a hemin and hemoglobin binding outer membrane protein of Vibrio vulnificus biotype 2 (serogroup E). FEMS Microbiology Letters, 2006, 156, 187-191.	1.8	10
77	Iron uptake from ferric citrate by Vibrio anguillarum. FEMS Microbiology Letters, 2006, 154, 145-150.	1.8	9
78	Genetic characterization of pPHDP60, a novel conjugative plasmid from the marine fish pathogen Photobacterium damselae subsp. piscicida. Plasmid, 2013, 70, 154-159.	1.4	9
79	Iron assimilation and siderophore production by Vibrio ordalii strains isolated from diseased Atlantic salmon Salmo salar in Chile. Diseases of Aquatic Organisms, 2016, 118, 217-226.	1.0	9
80	Preparation of functionalized magnetic nanoparticles conjugated with feroxamine and their evaluation for pathogen detection. RSC Advances, 2019, 9, 13533-13542.	3.6	9
81	Siderophore production by environmental strains of Salmonella species. FEMS Microbiology Letters, 1989, 57, 7-12.	1.8	9
82	The ABCâ€ŧransporter <i>hutCD</i> genes of <i>Photobacterium damselae</i> subsp. <i>piscicida</i> are essential for haem utilization as iron source and are expressed during infection in fish. Journal of Fish Diseases, 2010, 33, 649-655.	1.9	8
83	The Outer Membrane Protein FstC of Aeromonas salmonicida subsp. salmonicida Acts as Receptor for Amonabactin Siderophores and Displays a Wide Ligand Plasticity. Structure–Activity Relationships of Synthetic Amonabactin Analogues. ACS Infectious Diseases, 2019, 5, 1936-1951.	3.8	8
84	ADHESIVE PROPERTIES AND OTHER VIRULENCE FACTORS IN YERSINIA RUCKERI. , 1990, , 123-139.		8
85	Vibrio neptunius Produces Piscibactin and Amphibactin and Both Siderophores Contribute Significantly to Virulence for Clams. Frontiers in Cellular and Infection Microbiology, 2021, 11, 750567.	3.9	8
86	The marine bivalve molluscs pathogen Vibrio neptunius produces the siderophore amphibactin, which is widespread in molluscs microbiota. Environmental Microbiology, 2020, 22, 5467-5482.	3.8	7
87	dentification and characterisation of the fur genes in <i>Photobacterium damselae</i> ssp. <i>piscicida</i> and ssp. <i>damselae</i> . Diseases of Aquatic Organisms, 2004, 58, 151-156.	1.0	7
88	Application of suppressive subtractive hybridization to the identification of genetic differences between two Lactococcus garvieae strains showing distinct differences in virulence for rainbow trout and mouse. Microbiology (United Kingdom), 2011, 157, 2106-2119.	1.8	6
89	FrpA is the outer membrane piscibactin transporter in Vibrio anguillarum: structural elements in synthetic piscibactin analogues required for transport. Journal of Biological Inorganic Chemistry, 2022, 27, 133-142.	2.6	5
90	Highly preferred site of insertion of Tn7 into the chromosome of Vibrio anguillarum. Plasmid, 1992, 27, 161-163.	1.4	4

#	Article	IF	CITATIONS
91	The Vibriolysin-Like Protease VnpA and the Collagenase ColA Are Required for Full Virulence of the Bivalve Mollusks Pathogen Vibrio neptunius. Antibiotics, 2021, 10, 391.	3.7	4
92	The Temperature-Dependent Expression of the High-Pathogenicity Island Encoding Piscibactin in Vibrionaceae Results From the Combined Effect of the AraC-Like Transcriptional Activator PbtA and Regulatory Factors From the Recipient Genome. Frontiers in Microbiology, 2021, 12, 748147.	3.5	3
93	Selective detection of Aeromonas spp. by a fluorescent probe based on the siderophore amonabactin. Journal of Inorganic Biochemistry, 2022, 230, 111743.	3.5	3
94	Growth of Vibrio anguillarum in different fish sera. Aquaculture, 1992, 107, 277-281.	3.5	2
95	Identification of heme-binding proteins in the cell membranes of Vibrio anguillarum. FEMS Microbiology Letters, 1996, 135, 265-270.	1.8	2
96	Salmonella. , 2011, , 1051-1064.		2
97	Draft Genome Sequences of Five Vibrio neptunius Strains Isolated from Hatcheries of Bivalve Mollusks. Microbiology Resource Announcements, 2021, 10, .	0.6	1
98	Isolation of heme-binding proteins from Vibrio anguillarum using affinity chromatography. FEMS Microbiology Letters, 1996, 141, 19-23.	1.8	1
99	Ferric-reductase activities in Vibrio vulnificus biotypes 1 and 2. FEMS Microbiology Letters, 1999, 172, 205-211.	1.8	1
100	Iron uptake from ferric citrate by Vibrio anguillarum. FEMS Microbiology Letters, 1997, 154, 145-150.	1.8	1
101	BIOMETALS 2008 (Santiago de Compostela). BioMetals, 2009, 22, 1-2.	4.1	0
102	Synthesis of Functionalized Magnetic Nanoparticles, Their Conjugation with the Siderophore Feroxamine and its Evaluation for Bacteria Detection. Journal of Visualized Experiments, 2020, , .	0.3	0
103	Susceptibility to Bismuth(III) of Aquaculture Bacterial Pathogens: Effectiveness of Bismuth–Deferiprone Therapy against Vibrio anguillarum Infection in Fish. Microorganisms, 2021, 9, 2399.	3.6	0
104	Isolation of a hemin and hemoglobin binding outer membrane protein of Vibrio vulnificus biotype 2 (serogroup E). FEMS Microbiology Letters, 1997, 156, 187-191.	1.8	0