
Jan Mulder

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1429408/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Survival of occlusal ART restorations using high-viscosity glass-ionomer with and without chlorhexidine: A 2-year split-mouth quadruple-blind randomized controlled clinical trial. Journal of Advanced Research, 2019, 17, 117-123.	9.5	12
2	ls preventing micro-cavities in dentine from progressing with a sealant successful?. British Dental Journal, 2019, 226, 590-594.	0.6	5
3	High-viscosity glass-ionomer vs. composite resin restorations in persons with disability: Five-year follow-up of clinical trial. Brazilian Oral Research, 2019, 33, e099.	1.4	13
4	Three-year survival of ART high-viscosity glass-ionomer and resin composite restorations in people with disability. Clinical Oral Investigations, 2018, 22, 461-467.	3.0	25
5	Frequency of remnants of sealants left behind in pits and fissures of occlusal surfaces after 2 and 3Âyears. Clinical Oral Investigations, 2017, 21, 143-149.	3.0	4
6	Manual therapy compared with physical therapy in patients with non-specific neck pain: a randomized controlled trial. Chiropractic & Manual Therapies, 2017, 25, 12.	1.5	17
7	3-year survival rates of retained composite resin and ART sealants using two assessment criteria. Brazilian Oral Research, 2017, 31, e35.	1.4	9
8	A study on the survival of primary molars with intact and with defective restorations. International Journal of Paediatric Dentistry, 2016, 26, 383-390.	1.8	17
9	Dutch influenza vaccination rate drops for fifth consecutive year. Vaccine, 2015, 33, 4886-4891.	3.8	20
10	Caries experience in the primary dentition and presence of plaque in 7-year-old Chinese children: A 4-year time-lag study. Journal of International Society of Preventive and Community Dentistry, 2015, 5, 205.	1.0	3
11	The 3.5-year survival rates of primary molars treated according to three treatment protocols: a controlled clinical trial. Clinical Oral Investigations, 2014, 18, 1061-1069.	3.0	52
12	Oral Health Training Programs for Community and Professional Health Care Workers in Nairobi East District Increases Identification of HIV-Infected Patients. PLoS ONE, 2014, 9, e90927.	2.5	8
13	ls high-viscosity glass-ionomer-cement a successor to amalgam for treating primary molars?. Dental Materials, 2014, 30, 1172-1178.	3.5	39
14	Do light cured ART conventional high-viscosity glass-ionomer sealants perform better than resin-composite sealants: A 4-year randomized clinical trial. Dental Materials, 2014, 30, 487-492.	3.5	32
15	Amalgam and ART restorations in children: a controlled clinical trial. Clinical Oral Investigations, 2014, 18, 117-124.	3.0	35
16	What happens to cavitated primary teeth over time? A 3.5-year prospective cohort study in China. International Dental Journal, 2013, 63, 183-188.	2.6	13
17	A Community-Based Oral Health Promotion Model for HIV Patients in Nairobi, East District in Kenya: A Study Protocol. Journal of Public Health Research, 2013, 2, jphr.2013.e5.	1.2	5
18	Caries experience in a child population in a deprived area of Brazil, using ICDAS II. Clinical Oral Investigations, 2012, 16, 513-520.	3.0	78