

# MarÃ-a-Guadalupe Ramirez-DueÃ±as

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1469235/publications.pdf>

Version: 2024-02-01

19  
papers

527  
citations

840776

11  
h-index

794594

19  
g-index

19  
all docs

19  
docs citations

19  
times ranked

1213  
citing authors

#	ARTICLE	IF	CITATIONS
1	Low NKp30, NKp46 and NKG2D expression and reduced cytotoxic activity on NK cells in cervical cancer and precursor lesions. <i>BMC Cancer</i> , 2009, 9, 186.	2.6	164
2	Effect of d-limonene on immune response in BALB/c mice with lymphoma. <i>International Immunopharmacology</i> , 2005, 5, 829-838.	3.8	78
3	KIR/HLA Gene Profile Implication in Systemic Sclerosis Patients from Mexico. <i>Journal of Immunology Research</i> , 2019, 2019, 1-11.	2.2	68
4	Serum levels of macrophage migration inhibitory factor are associated with rheumatoid arthritis course. <i>Rheumatology International</i> , 2012, 32, 2307-2311.	3.0	33
5	The $\gamma$ 319C/+49G/CT60G Haplotype of CTLA-4 Gene Confers Susceptibility to Rheumatoid Arthritis in Mexican Population. <i>Cell Biochemistry and Biophysics</i> , 2013, 67, 1217-1228.	1.8	29
6	Protease-activated receptor-2 (PAR-2) in cervical cancer proliferation. <i>Gynecologic Oncology</i> , 2008, 108, 19-26.	1.4	26
7	Cytokines (IL-15, IL-21, and IFN- $\gamma$ ) in rheumatoid arthritis: association with positivity to autoantibodies (RF, anti-CCP, anti-MCV, and anti-PAD14) and clinical activity. <i>Clinical Rheumatology</i> , 2019, 38, 3061-3071.	2.2	22
8	Macrophage migration inhibitory factor polymorphisms are a potential susceptibility marker in systemic sclerosis from southern Mexican population: association with MIF mRNA expression and cytokine profile. <i>Clinical Rheumatology</i> , 2019, 38, 1643-1654.	2.2	17
9	The functional class evaluated in rheumatoid arthritis is associated with soluble TGF- $\beta$ 1 serum levels but not with G915C (Arg25Pro) TGF- $\beta$ 1 polymorphism. <i>Rheumatology International</i> , 2012, 32, 367-372.	3.0	15
10	<i>KIR2DL2</i> and <i>KIR2DS2</i> as genetic markers to the methotrexate response in rheumatoid arthritis patients. <i>Immunopharmacology and Immunotoxicology</i> , 2016, 38, 303-309.	2.4	15
11	Immunosuppressive activity of proteases in cervical carcinoma. <i>Gynecologic Oncology</i> , 2005, 98, 111-117.	1.4	13
12	MIF and TNF- $\alpha$ serum levels in rheumatoid arthritis patients treated with disease-modifying antirheumatic drugs: a cross-sectional study. <i>Immunopharmacology and Immunotoxicology</i> , 2015, 37, 207-213.	2.4	9
13	Distribution of KIR genes and KIR2DS4 gene variants in two Mexican Mestizo populations. <i>Human Immunology</i> , 2017, 78, 614-620.	2.4	8
14	TNFA -308G>A and -238G>A polymorphisms and risk to systemic sclerosis: impact on TNF- $\alpha$ serum levels, TNFA mRNA expression, and autoantibodies. <i>Clinical and Experimental Medicine</i> , 2019, 19, 439-447.	3.6	8
15	The 3'UTR 1188A/C polymorphism of IL-12p40 is not associated with susceptibility for developing plaque psoriasis in Mestizo population from western Mexico. <i>Immunology Letters</i> , 2015, 163, 221-226.	2.5	7
16	Transforming growth factor beta isoforms and TGF- $\beta$ 1 and TGF- $\beta$ 2 expression in systemic sclerosis patients. <i>Clinical and Experimental Medicine</i> , 2023, 23, 471-481.	3.6	6
17	<i>PTPN22</i> 1858C>T polymorphism is associated with increased CD154 expression and higher CD4+ T cells percentage in rheumatoid arthritis patients. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22710.	2.1	4
18	University students' knowledge and attitudes towards leprosy. <i>Journal of Infection in Developing Countries</i> , 2013, 7, 658-664.	1.2	3

#	ARTICLE	IF	CITATIONS
19	Diversity of KIR/HLA Genotypes and Their Association with Psoriasis Vulgaris in the Western Mexican Population. <i>Genes</i> , 2020, 11, 338.	2.4	2