Andrew C Harris

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1473070/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Cigarette Smoke Extract, but Not Electronic Cigarette Aerosol Extract, Inhibits Monoamine Oxidase in vitro and Produces Greater Acute Aversive/Anhedonic Effects Than Nicotine Alone on Intracranial Self-Stimulation in Rats. Frontiers in Neuroscience, 2022, 16, .	2.8	3
2	Behavioral predictors of individual differences in opioid addiction vulnerability as measured using i.v. self-administration in rats. Drug and Alcohol Dependence, 2021, 221, 108561.	3.2	9
3	Repeated morphine exposure activates synaptogenesis and other neuroplasticity-related gene networks in the dorsomedial prefrontal cortex of male and female rats. Drug and Alcohol Dependence, 2021, 221, 108598.	3.2	17
4	Magnitude of open-field thigmotaxis during mecamylamine-precipitated nicotine withdrawal in rats is influenced by mecamylamine dose, duration of nicotine infusion, number of withdrawal episodes, and age. Pharmacology Biochemistry and Behavior, 2021, 205, 173185.	2.9	3
5	Individual Differences in Different Measures of Opioid Self-Administration in Rats Are Accounted for by a Single Latent Variable. Frontiers in Psychiatry, 2021, 12, 712163.	2.6	1
6	Further pharmacological characterization of a preclinical model of the early development of nicotine withdrawal. Drug and Alcohol Dependence, 2021, 226, 108870.	3.2	2
7	β-Carbolines found in cigarette smoke elevate intracranial self-stimulation thresholds in rats. Pharmacology Biochemistry and Behavior, 2020, 198, 173041.	2.9	6
8	Higher anhedonia during withdrawal from initial opioid exposure is protective against subsequent opioid self-administration in rats. Psychopharmacology, 2020, 237, 2279-2291.	3.1	15
9	Comparison of the Relative Abuse Liability of Electronic Cigarette Aerosol Extracts and Nicotine Alone in Adolescent Rats: A Behavioral Economic Analysis. International Journal of Environmental Research and Public Health, 2020, 17, 860.	2.6	1
10	Non-nicotine constituents in e-cigarette aerosol extract attenuate nicotine's aversive effects in adolescent rats. Drug and Alcohol Dependence, 2019, 203, 51-60.	3.2	9
11	Locomotor activity does not predict individual differences in morphine self-administration in rats. Pharmacology Biochemistry and Behavior, 2018, 166, 48-56.	2.9	15
12	Consortium on Methods Evaluating Tobacco: Research Tools to Inform US Food and Drug Administration Regulation of Snus. Nicotine and Tobacco Research, 2018, 20, 1292-1300.	2.6	9
13	Effects of nicotine-containing and "nicotine-free―e-cigarette refill liquids on intracranial self-stimulation in rats. Drug and Alcohol Dependence, 2018, 185, 1-9.	3.2	16
14	Substitutability of nicotine alone and an electronic cigarette liquid using a concurrent choice assay in rats: A behavioral economic analysis. Drug and Alcohol Dependence, 2018, 185, 58-66.	3.2	13
15	Propylene glycol, a major electronic cigarette constituent, attenuates the adverse effects of high-dose nicotine as measured by intracranial self-stimulation in rats. Drug and Alcohol Dependence, 2018, 193, 162-168.	3.2	12
16	Status and future directions of preclinical behavioral pharmacology in tobacco regulatory science Behavior Analysis (Washington, D C), 2018, 18, 252-274.	0.5	11
17	Similar precipitated withdrawal effects on intracranial self-stimulation during chronic infusion of an e-cigarette liquid or nicotine alone. Pharmacology Biochemistry and Behavior, 2017, 161, 1-5.	2.9	12
18	Self-administration of Smokeless Tobacco Products in Rats. Tobacco Regulatory Science (discontinued), 2016, 2, 329-342.	0.2	12

ANDREW C HARRIS

#	Article	IF	CITATIONS
19	The Anti-(+)-Methamphetamine Monoclonal Antibody mAb7F9 Attenuates Acute (+)-Methamphetamine Effects on Intracranial Self-Stimulation in Rats. PLoS ONE, 2015, 10, e0118787.	2.5	11
20	A Two-Day Continuous Nicotine Infusion Is Sufficient to Demonstrate Nicotine Withdrawal in Rats as Measured Using Intracranial Self-Stimulation. PLoS ONE, 2015, 10, e0144553.	2.5	7
21	Blockade of cholinergic transmission elicits somatic signs in nicotine-naÃ⁻ve adolescent rats. Frontiers in Pharmacology, 2015, 6, 239.	3.5	3
22	Intracranial self-stimulation reward thresholds during morphine withdrawal in rats bred for high (HiS) and low (LoS) saccharin intake. Brain Research, 2015, 1602, 119-126.	2.2	21
23	Animal models to assess the abuse liability of tobacco products: Effects of smokeless tobacco extracts on intracranial self-stimulation. Drug and Alcohol Dependence, 2015, 147, 60-67.	3.2	27
24	Effects of nicotine and minor tobacco alkaloids on intracranial-self-stimulation in rats. Drug and Alcohol Dependence, 2015, 153, 330-334.	3.2	44
25	Effects of oxytocin on nicotine withdrawal in rats. Pharmacology Biochemistry and Behavior, 2014, 116, 84-89.	2.9	38
26	Restraint stress attenuates nicotine's locomotor stimulant but not discriminative stimulus effects in rats. Pharmacology Biochemistry and Behavior, 2014, 124, 92-100.	2.9	8
27	Mecamylamine elicits withdrawal-like signs in rats following a single dose of nicotine. Psychopharmacology, 2013, 225, 291-302.	3.1	15
28	Affective and neuroendocrine effects of withdrawal from chronic, long-acting opiate administration. Brain Research, 2013, 1538, 73-82.	2.2	9
29	Delivery of nicotine in an extract of a smokeless tobacco product reduces its reinforcement-attenuating and discriminative stimulus effects in rats. Psychopharmacology, 2012, 220, 565-576.	3.1	25
30	Combined active and passive immunization against nicotine: Minimizing monoclonal antibody requirements using a target antibody concentration strategy. International Immunopharmacology, 2011, 11, 1809-1815.	3.8	24
31	A lack of association between severity of nicotine withdrawal and individual differences in compensatory nicotine self-administration in rats. Psychopharmacology, 2011, 217, 153-166.	3.1	41
32	Comparison of the behavioral effects of cigarette smoke and pure nicotine in rats. Pharmacology Biochemistry and Behavior, 2010, 96, 217-227.	2.9	44
33	Passive immunization with a nicotine-specific monoclonal antibody decreases brain nicotine levels but does not precipitate withdrawal in nicotine-dependent rats. Pharmacology Biochemistry and Behavior, 2009, 93, 105-111.	2.9	34
34	Correlates of individual differences in compensatory nicotine self-administration in rats following a decrease in nicotine unit dose. Psychopharmacology, 2009, 205, 599-611.	3.1	15
35	Effects of the NMDA receptor antagonist memantine on the expression and development of acute opiate dependence as assessed by withdrawal-potentiated startle and hyperalgesia. Psychopharmacology, 2008, 196, 649-660.	3.1	29
36	Combined Active and Passive Immunization Enhances the Efficacy of Immunotherapy against Nicotine in Rats. Journal of Pharmacology and Experimental Therapeutics, 2008, 325, 985-993.	2.5	35

ANDREW C HARRIS

#	Article	IF	CITATIONS
37	Compensatory nicotine self-administration in rats during reduced access to nicotine: An animal model of smoking reduction Experimental and Clinical Psychopharmacology, 2008, 16, 86-97.	1.8	22
38	Prevalence, magnitude, and correlates of an extinction burst in drug-seeking behavior in rats trained to self-administer nicotine during unlimited access (23Âh/day) sessions. Psychopharmacology, 2007, 194, 395-402.	3.1	28
39	Acute opioid dependence: characterizing the early adaptations underlying drug withdrawal. Psychopharmacology, 2005, 178, 353-366.	3.1	72
40	Elevated startle during withdrawal from acute morphine: a model of opiate withdrawal and anxiety. Psychopharmacology, 2004, 171, 140-147.	3.1	65
41	Potentiated startle and hyperalgesia during withdrawal from acute morphine: effects of multiple opiate exposures. Psychopharmacology, 2004, 176, 266-273.	3.1	42
42	Delay Discounting and Performance on the Prisoner's Dilemma Game. Psychological Record, 2002, 52, 429-440.	0.9	90