Jorge Franco

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1476770/publications.pdf

Version: 2024-02-01

218677 206112 2,399 46 26 48 h-index citations g-index papers 49 49 49 2502 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Association Analysis of Historical Bread Wheat Germplasm Using Additive Genetic Covariance of Relatives and Population Structure. Genetics, 2007, 177, 1889-1913.	2.9	426
2	Genomic Prediction of Gene Bank Wheat Landraces. G3: Genes, Genomes, Genetics, 2016, 6, 1819-1834.	1.8	159
3	Genetic Characterization of CIMMYT Inbred Maize Lines and Open Pollinated Populations Using Large Scale Fingerprinting Methods. Crop Science, 2002, 42, 1832-1840.	1.8	141
4	Core Hunter: an algorithm for sampling genetic resources based on multiple genetic measures. BMC Bioinformatics, 2009, 10, 243.	2.6	138
5	Diversity analysis of 80,000 wheat accessions reveals consequences and opportunities of selection footprints. Nature Communications, 2020, 11, 4572.	12.8	129
6	Classifying Genetic Resources by Categorical and Continuous Variables. Crop Science, 1998, 38, 1688-1696.	1.8	115
7	A Sampling Strategy for Conserving Genetic Diversity when Forming Core Subsets. Crop Science, 2005, 45, 1035-1044.	1.8	92
8	Sampling Strategies for Conserving Maize Diversity When Forming Core Subsets Using Genetic Markers. Crop Science, 2006, 46, 854-864.	1.8	80
9	Evaluation of cowpea germplasm lines for protein and mineral concentrations in grains. Plant Genetic Resources: Characterisation and Utilisation, 2011, 9, 515-522.	0.8	77
10	Statistical methods for classifying genotypes. Euphytica, 2004, 137, 19-37.	1.2	75
11	Unlocking the genetic diversity of Creole wheats. Scientific Reports, 2016, 6, 23092.	3.3	75
12	The Development of Quality Control Genotyping Approaches: A Case Study Using Elite Maize Lines. PLoS ONE, 2016, 11, e0157236.	2.5	67
13	Varietal differentiation of Tannat, Cabernet-Sauvignon and Merlot grapes and wines according to their anthocyanic composition. European Food Research and Technology, 2007, 225, 111-117.	3.3	66
14	Classification of Peruvian highland maize races using plant traits. Genetic Resources and Crop Evolution, 2008, 55, 151-162.	1.6	62
15	QuEChERS Adaptability for the Analysis of Pesticide Residues in Beehive Products Seeking the Development of an Agroecosystem Sustainability Monitor. Journal of Agricultural and Food Chemistry, 2015, 63, 4484-4492.	5.2	56
16	Gene flow among different teosinte taxa and into the domesticated maize gene pool. Genetic Resources and Crop Evolution, 2011, 58, 1243-1261.	1.6	51
17	Genetic diversity and population structure of native maize populations in Latin America and the Caribbean. PLoS ONE, 2017, 12, e0173488.	2.5	50
18	A Twoâ€Stage, Threeâ€Way Method for Classifying Genetic Resources in multiple Environments. Crop Science, 1999, 39, 259-267.	1.8	45

#	Article	IF	CITATIONS
19	A Multivariate Method for Classifying Cultivars and Studying Group × Environment × Trait Interaction. Crop Science, 2003, 43, 1249-1258.	1.8	42
20	Genetic and phenotypic diversity in a germplasm working collection of cultivated tropical yams (Dioscorea spp.). Genetic Resources and Crop Evolution, 2012, 59, 1753-1765.	1.6	38
21	The Modified Location Model for Classifying Genetic Resources. Crop Science, 2002, 42, 1719-1726.	1.8	36
22	Genetic Characterization of a Core Set of a Tropical Maize Race Tuxpeño for Further Use in Maize Improvement. PLoS ONE, 2012, 7, e32626.	2.5	36
23	Evaluation of Carribean Maize Accessions to Develop a Core Subset. Crop Science, 1998, 38, 1378-1386.	1.8	33
24	Classifying Mexican Maize Accessions Using Hierarchical and Density Search Methods. Crop Science, 1997, 37, 972-980.	1.8	32
25	Assessment of Reaction Patterns of Hybrids to <i>Striga hermonthica</i> (Del.) Benth. under Artificial Infestation in Kenya and Nigeria. Crop Science, 2012, 52, 2528-2537.	1.8	30
26	Toward a Costâ€Effective Fingerprinting Methodology to Distinguish Maize Openâ€Pollinated Varieties. Crop Science, 2010, 50, 467-477.	1.8	28
27	Genetic diversity of physical, nutritional and functional properties of cowpea grain and relationships among the traits. Plant Genetic Resources: Characterisation and Utilisation, 2016, 14, 67-76.	0.8	20
28	Comparing a Preliminary Racial Classification with a Numerical Classification of the Maize Landraces of Uruguay. Crop Science, 2003, 43, 718.	1.8	19
29	Effects of larval diet on the development and reproduction of Argyrotaenia sphaleropa (Meyrick) (Lepidoptera: Tortricidae). Neotropical Entomology, 2003, 32, 551-557.	1.2	17
30	A Sequential Clustering Strategy for Classifying Gene Bank Accessions. Crop Science, 1997, 37, 1656-1662.	1.8	13
31	Re-defining the yam (<i>Dioscorea</i> spp.) core collection using morphological traits. Plant Genetic Resources: Characterisation and Utilisation, 2018, 16, 193-200.	0.8	13
32	Establishment of a cassava (<i>Manihot esculenta</i> Crantz) core collection based on agro-morphological descriptors. Plant Genetic Resources: Characterisation and Utilisation, 2012, 10, 119-127.	0.8	12
33	The impact of sample selection strategies on genetic diversity and representativeness in germplasm bank collections. BMC Plant Biology, 2019, 19, 520.	3.6	12
34	Effectiveness of essential oils for postharvest control of Phyllosticta citricarpa (citrus black spot) on citrus fruit. Postharvest Biology and Technology, 2016, 121, 1-8.	6.0	11
35	Biology of Bonagota cranaodes (Meyrick) (Lepidoptera: Tortricidae) on seven natural foods. Neotropical Entomology, 2004, 33, 299-306.	1.2	11
36	Hierarchical Multipleâ€Factor Analysis for Classifying Genotypes Based on Phenotypic and Genetic Data. Crop Science, 2010, 50, 105-117.	1.8	10

#	Article	IF	CITATIONS
37	Life history and assessment of grapevine phylloxera leaf galling incidence on Vitis species in Uruguay. SpringerPlus, 2013, 2, 181.	1.2	10
38	Intrafloral phenology of Trifolium polymorphum Poir. (Leguminosae) aerial flowers and reproductive implications. Acta Botanica Brasilica, 2009, 23, 881-888.	0.8	9
39	The Modified Location Model for Classifying Genetic Resources. Crop Science, 2002, 42, 1727-1736.	1.8	8
40	Parasitoid Niches of <i>Encarsia formosa </i> and <i>Encarsia Lycopersici </i> (Hymenoptera: Aphelinidae) Exploiting <i>Trialeurodes vaporariorum </i> (Hemiptera: Aleyrodidae). Florida Entomologist, 2012, 95, 1024-1030.	0.5	8
41	Use of Optimization Modeling to Assess the Effect of Timber and Carbon Pricing on Harvest Scheduling, Carbon Sequestration, and Net Present Value of Eucalyptus Plantations. Forests, 2021, 12, 651.	2.1	8
42	Strategic use of Iranian bread wheat landrace accessions for genetic improvement: Core set formulation and validation. Plant Breeding, 2021, 140, 87-99.	1.9	8
43	Effect of Different Diets on the Development, Mortality, Survival, Food Uptake and Fecundity of <i>Tupiocoris cucurbitaceus < i> (Hemiptera: Miridae). Florida Entomologist, 2014, 97, 1816-1824.</i>	0.5	6
44	Stand Characterization of Eucalyptus spp. Plantations in Uruguay Using Airborne Lidar Scanner Technology. Remote Sensing, 2020, 12, 3947.	4.0	6
45	Relationship between male moths of Cryptoblabes gnidiella (Millière) (Lepidoptera: Pyralidae) caught in sex pheromone traps and cumulative degree-days in vineyards in southern Uruguay. SpringerPlus, 2013, 2, 258.	1.2	4
46	Genetic Structure, Core Collection, and Regeneration Quality in White Dent Corn Landraces. Crop Science, 2018, 58, 1644-1658.	1.8	4