
Kota Mizumoto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1481114/publications.pdf Version: 2024-02-01



Κοτλ Μιζυμοτο

#	Article	lF	CITATIONS
1	Wnt Signals Can Function as Positional Cues in Establishing Cell Polarity. Developmental Cell, 2006, 10, 391-396.	7.0	155
2	Two βs or not two βs: regulation of asymmetric division by β-catenin. Trends in Cell Biology, 2007, 17, 465-473.	7.9	131
3	Cortical β-Catenin and APC Regulate Asymmetric Nuclear β-Catenin Localization during Asymmetric Cell Division in C. elegans. Developmental Cell, 2007, 12, 287-299.	7.0	113
4	Wnt Regulates Spindle Asymmetry to Generate Asymmetric Nuclear β-Catenin in C.Âelegans. Cell, 2011, 146, 942-954.	28.9	92
5	Interaxonal Interaction Defines Tiled Presynaptic Innervation in C.Âelegans. Neuron, 2013, 77, 655-666.	8.1	61
6	Systematic phenomics analysis of autism-associated genes reveals parallel networks underlying reversible impairments in habituation. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 656-667.	7.1	57
7	An intersectional gene regulatory strategy defines subclass diversity of C. elegans motor neurons. ELife, 2017, 6, .	6.0	42
8	CRISPR-Cas9 human gene replacement and phenomic characterization in <i>Caenorhabditis elegans</i> to understand the functional conservation of human genes and decipher variants of uncertain significance. DMM Disease Models and Mechanisms, 2018, 11, .	2.4	38
9	Two Wnts Instruct Topographic Synaptic Innervation in C.Âelegans. Cell Reports, 2013, 5, 389-396.	6.4	34
10	Altered expression of wheat AINTEGUMENTA homolog, WANT-1, in pistil and pistil-like transformed stamen of an alloplasmic line with Aegilops crassa cytoplasm. Development Genes and Evolution, 2009, 219, 175-187.	0.9	24
11	Tumor suppressor APC is an attenuator of spindle-pulling forces during <i>C. elegans</i> asymmetric cell division. Proceedings of the National Academy of Sciences of the United States of America, 2018, 115, E954-E963.	7.1	24
12	Nuclear and chloroplast genome genetic diversity in the wild einkorn wheat, Triticum urartu, revealed by AFLP and SSLP analyses. Hereditas, 2002, 137, 208-214.	1.4	22
13	Rap2 and TNIK control Plexin-dependent tiled synaptic innervation in C. elegans. ELife, 2018, 7, .	6.0	18
14	Characterization of wheat Bell1-type homeobox genes in floral organs of alloplasmic lines with Aegilops crassa cytoplasm. BMC Plant Biology, 2011, 11, 2.	3.6	14
15	Preferential expression of a HLP homolog encoding a mitochondrial L14 ribosomal protein in stamens of common wheat. Gene, 2004, 343, 281-289.	2.2	12
16	Intrinsic and extrinsic mechanisms of synapse formation and specificity in C. elegans. Cellular and Molecular Life Sciences, 2019, 76, 2719-2738.	5.4	12
17	Gradient-independent Wnt signaling instructs asymmetric neurite pruning in C. elegans. ELife, 2019, 8, .	6.0	7
18	Sustained expression of unc-4 homeobox gene and unc-37/Groucho in postmitotic neurons specifies the spatial organization of the cholinergic synapses in C. elegans. ELife, 2021, 10, .	6.0	6

#	Article	IF	CITATIONS
19	GFPnovo2, a brighter GFP variant for in vivo labeling in. MicroPublication Biology, 2018, 2018, .	0.1	4
20	Origin, dispersal and genomic structure of a low-copy-number hypervariable RFLP clone in Triticum and Aegilops species Genes and Genetic Systems, 2003, 78, 291-300.	0.7	3