Anne Bousquet-Melou

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1509307/publications.pdf Version: 2024-02-01

		430874	434195
32	1,072	18	31
papers	citations	h-index	g-index
32	32	32	1326
all docs	docs citations	times ranked	citing authors

43

#	Article	IF	CITATIONS
1	Lavender sensitivity to water stress: Comparison between eleven varieties across two phenological stages. Industrial Crops and Products, 2022, 177, 114531.	5.2	4
2	Chemical interaction between Quercus pubescens and its companion species is not emphasized under drought stress. European Journal of Forest Research, 2021, 140, 333-343.	2.5	1
3	Mediterranean woody plant specialized metabolites affect germination of Linum perenne at its dry and upper thermal limits. Plant and Soil, 2020, 446, 291-305.	3.7	4
4	Vegetation dynamics and regeneration of Pinus pinea forests in Mount Lebanon: Towards the progressive disappearance of pine. Ecological Engineering, 2020, 152, 105866.	3.6	8
5	Soil scarification favors natural regeneration of Pinus pinea in Lebanon forests: Evidences from field and laboratory experiments. Forest Ecology and Management, 2020, 459, 117840.	3.2	5
6	Exogenous Isoprene Confers Physiological Benefits in a Negligible Isoprene Emitter (Acer) Tj ETQq0 0 0 rgBT /Ov	erl <u>gck</u> 10 ⁻	Tf 50 542 Td (
7	Allelopathic effects of volatile organic compounds released from <i>Pinus halepensis</i> needles and roots. Ecology and Evolution, 2019, 9, 8201-8213.	1.9	42
8	Phenolics of the understory shrub Cotinus coggygria influence Mediterranean oak forests diversity and dynamics. Forest Ecology and Management, 2019, 441, 262-270.	3.2	14
9	Resistance of native oak to recurrent drought conditions simulating predicted climatic changes in the <scp>Mediterranean</scp> region. Plant, Cell and Environment, 2018, 41, 2299-2312.	5.7	20
10	Do litter-mediated plant-soil feedbacks influence Mediterranean oak regeneration? A two-year pot experiment. Plant and Soil, 2018, 430, 59-71.	3.7	12
11	Biotic interactions in a Mediterranean oak forest: role of allelopathy along phenological development of woody species. European Journal of Forest Research, 2017, 136, 699-710.	2.5	18
12	The Impact of Competition and Allelopathy on the Trade-Off between Plant Defense and Growth in Two Contrasting Tree Species. Frontiers in Plant Science, 2016, 7, 594.	3.6	78
13	Secondary metabolites of <i><scp>P</scp>inus halepensis</i> alter decomposer organisms and litter decomposition during afforestation of abandoned agricultural zones. Journal of Ecology, 2014, 102, 411-424.	4.0	68
14	Allelochemicals of Pinus halepensis as Drivers of Biodiversity in Mediterranean Open Mosaic Habitats During the Colonization Stage of Secondary Succession. Journal of Chemical Ecology, 2013, 39, 298-311.	1.8	59
15	How nutrient availability influences acclimation to shade of two (pioneer and late-successional) Mediterranean tree species?. European Journal of Forest Research, 2013, 132, 325-333.	2.5	9
16	Effects of different site preparation treatments on species diversity, composition, and plant traits in Pinus halepensis woodlands. Plant Ecology, 2011, 212, 627-638.	1.6	21
17	Fertilization and allelopathy modify Pinus halepensis saplings crown acclimation to shade. Trees - Structure and Function, 2011, 25, 497-507.	1.9	13
18	Variations in Allelochemical Composition of Leachates of Different Organs and Maturity Stages of	1.0	49

18Variations in Allelochemical Composition of Leachates of Different Organs and Maturity Stages of
Pinus halepensis. Journal of Chemical Ecology, 2009, 35, 970-979.1.8

#	Article	IF	CITATIONS
19	Regeneration failure of Pinus halepensis Mill.: The role of autotoxicity and some abiotic environmental parameters. Forest Ecology and Management, 2008, 255, 2928-2936.	3.2	75
20	Water deficit stress induces different monoterpene and sesquiterpene emission changes in Mediterranean species. Relationship between terpene emissions and plant water potential. Chemosphere, 2007, 67, 276-284.	8.2	152
21	Monoterpene and sesquiterpene emissions of three Mediterranean species through calcareous and siliceous soils in natural conditions. Atmospheric Environment, 2007, 41, 629-639.	4.1	58
22	Effect of Intraspecific Competition and Substrate Type on Terpene Emissions from Some Mediterranean Plant Species. Journal of Chemical Ecology, 2007, 33, 277-286.	1.8	23
23	Potential allelopathic effect of Pinus halepensis in the secondary succession: an experimental approach. Chemoecology, 2006, 16, 97-105.	1.1	83
24	Chemical composition of the volatile oil of Laggera aurita Schulz from Burkina-Faso. Biochemical Systematics and Ecology, 2006, 34, 815-818.	1.3	13
25	Contribution of some Mediterranean plants to BVOC in the atmosphere of an open and a closed environment: a preliminary study. WIT Transactions on Ecology and the Environment, 2006, , .	0.0	0
26	Allelopathic potential of Medicago arborea, a Mediterranean invasive shrub. Chemoecology, 2005, 15, 193-198.	1.1	24
27	Comparison of essential oil composition of two varieties of Cistus ladanifer. Biochemical Systematics and Ecology, 2003, 31, 339-343.	1.3	25
28	Phenolic compounds content in Pinus halepensis Mill. needles: a bioindicator of air pollution. Chemosphere, 2003, 52, 239-248.	8.2	69
29	Phenols and Flavonoids in Aleppo Pine Needles as Bioindicators of Air Pollution. Journal of Environmental Quality, 2003, 32, 2265-2271.	2.0	51
30	Evolutionary divergence in the pan-Atlantic mangrove Avicennia germinans. New Phytologist, 2000, 145, 115-125.	7.3	23
31	Inter-specific variation in the concentration of two iridoid glucosides in Avicennia L. (Avicenniaceae) Tj ETQq1 1 C	.784314 ı 1.3	gBT /Overloc
32	Iridoid glucosides from Avicennia germinans. Phytochemistry, 1995, 38, 893-894.	2.9	37