

Noriyuki Matsuda, æ³/₄ç”°æ²ä¹<

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1532973/publications.pdf>

Version: 2024-02-01

50
papers

7,825
citations

136950

32
h-index

182427

51
g-index

52
all docs

52
docs citations

52
times ranked

10469
citing authors

#	ARTICLE	IF	CITATIONS
1	PINK1 stabilized by mitochondrial depolarization recruits Parkin to damaged mitochondria and activates latent Parkin for mitophagy. <i>Journal of Cell Biology</i> , 2010, 189, 211-221.	5.2	1,600
2	Ubiquitin is phosphorylated by PINK1 to activate parkin. <i>Nature</i> , 2014, 510, 162-166.	27.8	1,185
3	Molecular mechanisms and physiological functions of mitophagy. <i>EMBO Journal</i> , 2021, 40, e104705.	7.8	553
4	UV-Induced Ubiquitylation of XPC Protein Mediated by UV-DDB-Ubiquitin Ligase Complex. <i>Cell</i> , 2005, 121, 387-400.	28.9	517
5	PINK1 autophosphorylation upon membrane potential dissipation is essential for Parkin recruitment to damaged mitochondria. <i>Nature Communications</i> , 2012, 3, 1016.	12.8	465
6	MG53 nucleates assembly of cell membrane repair machinery. <i>Nature Cell Biology</i> , 2009, 11, 56-64.	10.3	396
7	p62/SQSTM1 cooperates with Parkin for perinuclear clustering of depolarized mitochondria. <i>Genes To Cells</i> , 2010, 15, 887-900.	1.2	345
8	Phosphorylated ubiquitin chain is the genuine Parkin receptor. <i>Journal of Cell Biology</i> , 2015, 209, 111-128.	5.2	217
9	The ubiquitin signal and autophagy: an orchestrated dance leading to mitochondrial degradation. <i>EMBO Reports</i> , 2016, 17, 300-316.	4.5	197
10	Parkin-catalyzed Ubiquitin-Ester Transfer Is Triggered by PINK1-dependent Phosphorylation. <i>Journal of Biological Chemistry</i> , 2013, 288, 22019-22032.	3.4	173
11	A Dimeric PINK1-containing Complex on Depolarized Mitochondria Stimulates Parkin Recruitment. <i>Journal of Biological Chemistry</i> , 2013, 288, 36372-36384.	3.4	168
12	Diverse Effects of Pathogenic Mutations of Parkin That Catalyze Multiple Monoubiquitylation in Vitro. <i>Journal of Biological Chemistry</i> , 2006, 281, 3204-3209.	3.4	166
13	Proteostasis and neurodegeneration: The roles of proteasomal degradation and autophagy. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2014, 1843, 197-204.	4.1	153
14	Critical role of mitochondrial ubiquitination and the OPTN-ATG9A axis in mitophagy. <i>Journal of Cell Biology</i> , 2020, 219, .	5.2	114
15	Endosomal Rab cycles regulate Parkin-mediated mitophagy. <i>ELife</i> , 2018, 7, .	6.0	113
16	Direct interactions between NEDD8 and ubiquitin E2 conjugating enzymes upregulate cullin-based E3 ligase activity. <i>Nature Structural and Molecular Biology</i> , 2007, 14, 167-168.	8.2	105
17	Unconventional PINK1 localization mechanism to the outer membrane of depolarized mitochondria drives Parkin recruitment. <i>Journal of Cell Science</i> , 2015, 128, 964-78.	2.0	103
18	EL5, a rice N-acetylchitooligosaccharide elicitor-responsive RING-H2 finger protein, is a ubiquitin ligase which functions in vitro in co-operation with an elicitor-responsive ubiquitin-conjugating enzyme, OsUBC5b. <i>Plant Journal</i> , 2002, 30, 447-455.	5.7	98

#	ARTICLE	IF	CITATIONS
37	Discovery and Optimization of Inhibitors of the Parkinson's Disease Associated Protein DJ-1. ACS Chemical Biology, 2018, 13, 2783-2793.	3.4	27
38	A palmitoylated RING finger ubiquitin ligase and its homologue in the brain membranes. Journal of Neurochemistry, 2003, 86, 749-762.	3.9	25
39	Constitutive Activation of PINK1 Protein Leads to Proteasome-mediated and Non-apoptotic Cell Death Independently of Mitochondrial Autophagy. Journal of Biological Chemistry, 2016, 291, 16162-16174.	3.4	23
40	Loss of peptide: <i>N</i> -glycanase causes proteasome dysfunction mediated by a sugar-recognizing ubiquitin ligase. Proceedings of the National Academy of Sciences of the United States of America, 2021, 118, .	7.1	23
41	Modes of interaction between the Arabidopsis Rab protein, Ara4, and its putative regulator molecules revealed by a yeast expression system. Plant Journal, 2000, 21, 341-349.	5.7	21
42	Unexpected mitochondrial matrix localization of Parkinson's disease-related DJ-1 mutants but not wild-type DJ-1. Genes To Cells, 2016, 21, 772-788.	1.2	21
43	Cleaved PGAM5 is released from mitochondria depending on proteasome-mediated rupture of the outer mitochondrial membrane during mitophagy. Journal of Biochemistry, 2019, 165, 19-25.	1.7	19
44	Overexpression of PRA2, a Rab/Yipt-family Small GTPase from Pea Pisum sativum, Aggravates the Growth Defect of Yeast ypt Mutants.. Cell Structure and Function, 2000, 25, 11-20.	1.1	9
45	Two sides of a coin: Physiological significance and molecular mechanisms for damage-induced mitochondrial localization of PINK1 and Parkin. Neuroscience Research, 2020, 159, 16-24.	1.9	8
46	The PARK2/Parkin receptor on damaged mitochondria revisited—uncovering the role of phosphorylated ubiquitin chains. Autophagy, 2015, 11, 1700-1701.	9.1	6
47	Tagged tags engage disposal. Nature, 2015, 524, 294-295.	27.8	6
48	Mammalian BCAS3 and C16orf70 associate with the phagophore assembly site in response to selective and non-selective autophagy. Autophagy, 2021, 17, 2011-2036.	9.1	6
49	Unfolding is the driving force for mitochondrial import and degradation of the Parkinson's disease-related protein DJ-1. Journal of Cell Science, 2021, 134, .	2.0	3
50	Cleaved PGAM5 dephosphorylates nuclear serine/arginine-rich proteins during mitophagy. Biochimica Et Biophysica Acta - Molecular Cell Research, 2021, 1868, 119045.	4.1	2