

Karin Leandersson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1559039/publications.pdf>

Version: 2024-02-01

54
papers

2,945
citations

201674

27
h-index

175258

52
g-index

55
all docs

55
docs citations

55
times ranked

6129
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Branching Copy-Number Evolution and Parallel Immune Profiles across the Regional Tumor Space of Resected Pancreatic Cancer. <i>Molecular Cancer Research</i> , 2022, 20, 749-761. | 3.4 | 3 |
| 2 | High Infiltration of CD68 ⁺ /CD163 ⁺ Macrophages Is an Adverse Prognostic Factor after Neoadjuvant Chemotherapy in Esophageal and Gastric Adenocarcinoma. <i>Journal of Innate Immunity</i> , 2022, 14, 615-628. | 3.8 | 8 |
| 3 | Peripheral Blood Mononuclear Cell Populations Correlate with Outcome in Patients with Metastatic Breast Cancer. <i>Cells</i> , 2022, 11, 1639. | 4.1 | 8 |
| 4 | T cells, B cells, and PD-L1 expression in esophageal and gastric adenocarcinoma before and after neoadjuvant chemotherapy: relationship with histopathological response and survival. <i>Oncolmmunology</i> , 2021, 10, 1921443. | 4.6 | 14 |
| 5 | Infiltration of NK and plasma cells is associated with a distinct immune subset in non-small cell lung cancer. <i>Journal of Pathology</i> , 2021, 255, 243-256. | 4.5 | 17 |
| 6 | The Immune Landscape of Colorectal Cancer. <i>Cancers</i> , 2021, 13, 5545. | 3.7 | 14 |
| 7 | Clinical relevance of systemic monocytic-MDSCs in patients with metastatic breast cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 435-448. | 4.2 | 44 |
| 8 | Prognostic implications of the expression levels of different immunoglobulin heavy chain-encoding RNAs in early breast cancer. <i>Npj Breast Cancer</i> , 2020, 6, 28. | 5.2 | 25 |
| 9 | Deletion of Nemo-like Kinase in T Cells Reduces Single-Positive CD8 ⁺ Thymocyte Population. <i>Journal of Immunology</i> , 2020, 205, 1830-1841. | 0.8 | 4 |
| 10 | Co-localization of CD169 ⁺ macrophages and cancer cells in lymph node metastases of breast cancer patients is linked to improved prognosis and PDL1 expression. <i>Oncolmmunology</i> , 2020, 9, 1848067. | 4.6 | 9 |
| 11 | Topographical Distribution and Spatial Interactions of Innate and Semi-Innate Immune Cells in Pancreatic and Other Periampullary Adenocarcinoma. <i>Frontiers in Immunology</i> , 2020, 11, 558169. | 4.8 | 18 |
| 12 | Tumor-Associated CD68 ⁺ , CD163 ⁺ , and MARCO ⁺ Macrophages as Prognostic Biomarkers in Patients With Treatment-Naïve Gastroesophageal Adenocarcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 534761. | 2.8 | 20 |
| 13 | Inflammatory macrophage derived TNF α downregulates estrogen receptor α via FOXO3a inactivation in human breast cancer cells. <i>Experimental Cell Research</i> , 2020, 390, 111932. | 2.6 | 7 |
| 14 | The Generation and Identity of Human Myeloid-Derived Suppressor Cells. <i>Frontiers in Oncology</i> , 2020, 10, 109. | 2.8 | 77 |
| 15 | Quantitative, qualitative and spatial analysis of lymphocyte infiltration in periampullary and pancreatic adenocarcinoma. <i>International Journal of Cancer</i> , 2020, 146, 3461-3473. | 5.1 | 39 |
| 16 | Chemotherapy, host response and molecular dynamics in periampullary cancer: the CHAMP study. <i>BMC Cancer</i> , 2020, 20, 308. | 2.6 | 9 |
| 17 | Complement inhibitor factor H expressed by breast cancer cells differentiates CD14 ⁺ human monocytes into immunosuppressive macrophages. <i>Oncolmmunology</i> , 2020, 9, 1731135. | 4.6 | 20 |
| 18 | Human G-MDSCs are neutrophils at distinct maturation stages promoting tumor growth in breast cancer. <i>Life Science Alliance</i> , 2020, 3, e202000893. | 2.8 | 14 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | The STAT3 inhibitor galiellalactone inhibits the generation of MDSC-like monocytes by prostate cancer cells and decreases immunosuppressive and tumorigenic factors. <i>Prostate</i> , 2019, 79, 1611-1621. | 2.3 | 47 |
| 20 | Clinical impact of T cells, B cells and the PD-1/PD-L1 pathway in muscle invasive bladder cancer: a comparative study of transurethral resection and cystectomy specimens. <i>Oncolmmunology</i> , 2019, 8, e1644108. | 4.6 | 34 |
| 21 | Impact of systemic therapy on circulating leukocyte populations in patients with metastatic breast cancer. <i>Scientific Reports</i> , 2019, 9, 13451. | 3.3 | 21 |
| 22 | Localization and Regulation of Polymeric Ig Receptor in Healthy and Diseased Human Kidney. <i>American Journal of Pathology</i> , 2019, 189, 1933-1944. | 3.8 | 10 |
| 23 | Wnt5a is a TLR2/4-ligand that induces tolerance in human myeloid cells. <i>Communications Biology</i> , 2019, 2, 176. | 4.4 | 24 |
| 24 | Expression of PD-L1 and PD-1 in Chemoradiotherapy-Naïve Esophageal and Gastric Adenocarcinoma: Relationship With Mismatch Repair Status and Survival. <i>Frontiers in Oncology</i> , 2019, 9, 136. | 2.8 | 36 |
| 25 | Pre-diagnostic anthropometry, sex, and risk of colorectal cancer according to tumor immune cell composition. <i>Oncolmmunology</i> , 2019, 8, e1664275. | 4.6 | 5 |
| 26 | Docetaxel promotes the generation of anti-tumorigenic human macrophages. <i>Experimental Cell Research</i> , 2018, 362, 525-531. | 2.6 | 34 |
| 27 | Infiltration of CD4+ T cells, IL-17+ T cells and FoxP3+ T cells in human breast cancer. <i>Cancer Biomarkers</i> , 2018, 20, 395-409. | 1.7 | 22 |
| 28 | Expression of programmed cell death protein 1 (PD-1) and its ligand PD-L1 in colorectal cancer: Relationship with sidedness and prognosis. <i>Oncolmmunology</i> , 2018, 7, e1465165. | 4.6 | 59 |
| 29 | The clinical impact of tumour-infiltrating lymphocytes in colorectal cancer differs by anatomical subsite: A cohort study. <i>International Journal of Cancer</i> , 2017, 141, 1654-1666. | 5.1 | 65 |
| 30 | Papillary renal cell carcinoma-derived chemerin, IL-8, and CXCL16 promote monocyte recruitment and differentiation into foam-cell macrophages. <i>Laboratory Investigation</i> , 2017, 97, 1296-1305. | 3.7 | 28 |
| 31 | The clinical importance of tumour-infiltrating macrophages and dendritic cells in periampullary adenocarcinoma differs by morphological subtype. <i>Journal of Translational Medicine</i> , 2017, 15, 152. | 4.4 | 33 |
| 32 | On the origin of myeloid-derived suppressor cells. <i>Oncotarget</i> , 2017, 8, 3649-3665. | 1.8 | 156 |
| 33 | The prognostic impact of tumor-infiltrating lymphocytes in colorectal cancer differs by anatomical subsite.. <i>Journal of Clinical Oncology</i> , 2017, 35, 47-47. | 1.6 | 1 |
| 34 | The integrative clinical impact of tumor-infiltrating T lymphocytes and NK cells in relation to B lymphocyte and plasma cell density in esophageal and gastric adenocarcinoma. <i>Oncotarget</i> , 2017, 8, 72108-72126. | 1.8 | 53 |
| 35 | Prognostic significance of professional antigen presenting cells according to morphological subtype of periampullary adenocarcinoma.. <i>Journal of Clinical Oncology</i> , 2017, 35, 121-121. | 1.6 | 0 |
| 36 | Cartilage oligomeric matrix protein contributes to the development and metastasis of breast cancer. <i>Oncogene</i> , 2016, 35, 5585-5596. | 5.9 | 74 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 37 | Cancer-associated fibroblast-secreted CXCL16 attracts monocytes to promote stroma activation in triple-negative breast cancers. <i>Nature Communications</i> , 2016, 7, 13050. | 12.8 | 135 |
| 38 | Dual mechanisms of action of the RNA-binding protein human antigen R explains its regulatory effect on melanoma cell migration. <i>Translational Research</i> , 2016, 172, 45-60. | 5.0 | 19 |
| 39 | The Prognostic Impact of NK/NKT Cell Density in Periampullary Adenocarcinoma Differs by Morphological Type and Adjuvant Treatment. <i>PLoS ONE</i> , 2016, 11, e0156497. | 2.5 | 32 |
| 40 | Systemic Monocytic-MDSCs Are Generated from Monocytes and Correlate with Disease Progression in Breast Cancer Patients. <i>PLoS ONE</i> , 2015, 10, e0127028. | 2.5 | 116 |
| 41 | Prognostic stromal gene signatures in breast cancer. <i>Breast Cancer Research</i> , 2015, 17, 23. | 5.0 | 67 |
| 42 | S100A9 expressed in ER ⁺ /PgR ⁺ breast cancers induces inflammatory cytokines and is associated with an impaired overall survival. <i>British Journal of Cancer</i> , 2015, 113, 1234-1243. | 6.4 | 35 |
| 43 | Expression of functional toll like receptor 4 in estrogen receptor/progesterone receptor-negative breast cancer. <i>Breast Cancer Research</i> , 2015, 17, 130. | 5.0 | 41 |
| 44 | Heterogeneity among septic shock patients in a set of immunoregulatory markers. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2014, 33, 313-324. | 2.9 | 15 |
| 45 | WNT5A induces release of exosomes containing pro-angiogenic and immunosuppressive factors from malignant melanoma cells. <i>Molecular Cancer</i> , 2014, 13, 88. | 19.2 | 213 |
| 46 | Infiltration of CD3 ⁺ and CD68 ⁺ cells in bladder cancer is subtype specific and affects the outcome of patients with muscle-invasive tumors Grant support: The Swedish Cancer Society, the Swedish research council, the Nilsson Cancer foundation, the BioCARE Strategic Cancer Research program, the Lund Medical Faculty, and FoU Landstinget Kronoberg and SÄdra Region <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 791-797. | 1.6 | 106 |
| 47 | A high frequency of MDSCs in sepsis patients, with the granulocytic subtype dominating in gram-positive cases. <i>Journal of Leukocyte Biology</i> , 2014, 96, 685-693. | 3.3 | 128 |
| 48 | Wnt5a Inhibits Human Monocyte-Derived Myeloid Dendritic Cell Generation. <i>Scandinavian Journal of Immunology</i> , 2013, 78, 194-204. | 2.7 | 21 |
| 49 | Wnt5a Induces a Tolerogenic Phenotype of Macrophages in Sepsis and Breast Cancer Patients. <i>Journal of Immunology</i> , 2012, 188, 5448-5458. | 0.8 | 100 |
| 50 | The presence of tumor associated macrophages in tumor stroma as a prognostic marker for breast cancer patients. <i>BMC Cancer</i> , 2012, 12, 306. | 2.6 | 531 |
| 51 | T cells developing in fetal thymus of T-cell receptor α -chain transgenic mice colonize β T-cell-specific epithelial niches but lack long-term reconstituting potential. <i>Immunology</i> , 2006, 119, 134-142. | 4.4 | 6 |
| 52 | Wnt-5a mRNA translation is suppressed by the Elav-like protein HuR in human breast epithelial cells. <i>Nucleic Acids Research</i> , 2006, 34, 3988-3999. | 14.5 | 86 |
| 53 | Wnt-5a/Ca ²⁺ -Induced NFAT Activity Is Counteracted by Wnt-5a/Yes-Cdc42-Casein Kinase 1 α Signaling in Human Mammary Epithelial Cells. <i>Molecular and Cellular Biology</i> , 2006, 26, 6024-6036. | 2.3 | 144 |
| 54 | Expression and signaling activity of Wnt-5a/discoidin domain receptor-1 and Syk plays distinct but decisive roles in breast cancer patient survival. <i>Clinical Cancer Research</i> , 2005, 11, 520-8. | 7.0 | 89 |