

# Henning KelbÄk

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1577826/publications.pdf>

Version: 2024-02-01

34  
papers

1,270  
citations

567281

15  
h-index

377865

34  
g-index

34  
all docs

34  
docs citations

34  
times ranked

2207  
citing authors

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 1  | Mechanisms of Very Late Drug-Eluting Stent Thrombosis Assessed by Optical Coherence Tomography. <i>Circulation</i> , 2016, 133, 650-660.  | 1.6  | 260       |
| 2  | Early Versus Standard Care Invasive Examination and Treatment of Patients With Non-ST-Segment Elevation Acute Coronary Syndrome. <i>Circulation</i> , 2018, 138, 2741-2750.   | 1.6  | 168       |
| 3  | No Major Differences in 30-Day Outcomes in High-Risk Patients Randomized to Off-Pump Versus On-Pump Coronary Bypass Surgery. <i>Circulation</i> , 2010, 121, 498-504.   | 1.6  | 140       |
| 4  | Final infarct size measured by cardiovascular magnetic resonance in patients with ST elevation myocardial infarction predicts long-term clinical outcome: an observational study. <i>European Heart Journal Cardiovascular Imaging</i> , 2013, 14, 387-395.   | 1.2  | 124       |
| 5  | Differential clinical outcomes after 1 year versus 5 years in a randomised comparison of zotarolimus-eluting and sirolimus-eluting coronary stents (the SORT OUT III study): a multicentre, open-label, randomised superiority trial. <i>Lancet</i> , The, 2014, 383, 2047-2056.  | 13.7 | 96        |
| 6  | The Third DANish Study of Optimal Acute Treatment of Patients with ST-segment Elevation Myocardial Infarction: Ischemic postconditioning or deferred stent implantation versus conventional primary angioplasty and complete revascularization versus treatment of culprit lesion only. <i>American Heart Journal</i> , 2015, 169, 613-621. | 2.7  | 61        |
| 7  | Impact of Acute Hyperglycemia on Myocardial Infarct Size, Area at Risk, and Salvage in Patients With STEMI and the Association With Exenatide Treatment: Results From a Randomized Study. <i>Diabetes</i> , 2014, 63, 2474-2485.  | 0.6  | 59        |
| 8  | Biolimus-Eluting Stents With Biodegradable Polymer Versus Bare-Metal Stents in Acute Myocardial Infarction. <i>Circulation: Cardiovascular Interventions</i> , 2014, 7, 355-364.  | 3.9  | 56        |
| 9  | Influence of pre-infarction angina, collateral flow, and pre-procedural TIMI flow on myocardial salvage index by cardiac magnetic resonance in patients with ST-segment elevation myocardial infarction. <i>European Heart Journal Cardiovascular Imaging</i> , 2012, 13, 433-443.  | 1.2  | 48        |
| 10 | Left Ventricular Hypertrophy Is Associated With Increased Infarct Size and Decreased Myocardial Salvage in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. <i>Journal of the American Heart Association</i> , 2017, 6, .  | 3.7  | 39        |
| 11 | Fractional Flow Reserve-Guided Complete Revascularization Improves the Prognosis in Patients With ST-Segment Elevation Myocardial Infarction and Severe Nonculprit Disease. <i>Circulation: Cardiovascular Interventions</i> , 2017, 10, .  | 3.9  | 39        |
| 12 | One-year results of total arterial revascularization vs. conventional coronary surgery: CARRPO trial. <i>European Heart Journal</i> , 2008, 30, 1005-1011.  | 2.2  | 22        |
| 13 | Comparison of Selvester QRS score with magnetic resonance imaging measured infarct size in patients with ST elevation myocardial infarction. <i>Journal of Electrocardiology</i> , 2012, 45, 414-419.   | 0.9  | 20        |
| 14 | Absolute quantitation of left ventricular wall and cavity parameters using ECG-gated PET. <i>Journal of Nuclear Cardiology</i> , 2004, 11, 38-46.   | 2.1  | 17        |
| 15 | Graft patency after off-pump versus on-pump coronary artery surgery in high-risk patients. <i>Scandinavian Cardiovascular Journal</i> , 2010, 44, 161-167.  | 1.2  | 17        |
| 16 | Low whole-body insulin sensitivity in patients with ischaemic heart disease is associated with impaired myocardial glucose uptake predictive of poor outcome after revascularisation. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2002, 29, 991-998.  | 6.4  | 11        |
| 17 | Coronary and Skeletal Muscle Enzyme Changes during a 14 km Run. <i>Acta Medica Scandinavica</i> , 1988, 224, 183-186.   | 0.0  | 10        |
| 18 | Scintigraphic evaluation of routine filterwire distal protection in percutaneous coronary intervention for acute ST-segment elevation myocardial infarction: a randomized controlled trial. <i>Journal of Nuclear Cardiology</i> , 2009, 16, 784-791.   | 2.1  | 10        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Zotarolimus-eluting vs. sirolimus-eluting coronary stents in patients with and without acute coronary syndromes: a SORT OUT III substudy. <i>European Journal of Clinical Investigation</i> , 2012, 42, 1047-1054.   | 3.4 | 10        |
| 20 | Association Between Early Q Waves and Reperfusion Success in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention. <i>Circulation: Cardiovascular Interventions</i> , 2017, 10, .  | 3.9 | 10        |
| 21 | Pre- and afterload reduction in chronic mitral regurgitation: a double-blind randomized placebo-controlled trial of the acute and 2 weeks' effect of nifedipine or isosorbide dinitrate treatment on left ventricular function and the severity of mitral regurg. <i>British Journal of Clinical Pharmacology</i> , 1996, 41, 493-497. | 2.4 | 9         |
| 22 | Stent Thrombosis is the Primary Cause of ST-Segment Elevation Myocardial Infarction following Coronary Stent Implantation: A Five Year Follow-Up of the SORT OUT II Study. <i>PLoS ONE</i> , 2014, 9, e113399.   | 2.5 | 8         |
| 23 | Comparison of methods of fractional area change for detection of regional left ventricular dysfunction. <i>International Journal of Cardiovascular Imaging</i> , 2000, 16, 257-266.  | 0.6 | 6         |
| 24 | Subacute cardiac rubidium-82 positron emission tomography (82Rb-PET) to assess myocardial area at risk, final infarct size, and myocardial salvage after STEMI. <i>Journal of Nuclear Cardiology</i> , 2018, 25, 970-981.  | 2.1 | 6         |
| 25 | Regional myocardial oxygen consumption estimated by carbon-11 acetate and positron emission tomography before and after repetitive ischemia. <i>Journal of Nuclear Cardiology</i> , 2000, 7, 228-234.  | 2.1 | 4         |
| 26 | Variability of insulin-stimulated myocardial glucose uptake in healthy elderly subjects. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2002, 29, 1600-1607.  | 6.4 | 4         |
| 27 | Impact of age on reperfusion success and long-term prognosis in ST-segment elevation myocardial infarction – A cardiac magnetic resonance imaging study. <i>IJC Heart and Vasculature</i> , 2021, 33, 100731.  | 1.1 | 4         |
| 28 | Amlodipine reduces myocardial ischaemia during exercise without compromising left ventricular function in patients with silent ischaemia: a randomised, double-blind, placebo-controlled study. <i>European Journal of Heart Failure</i> , 1999, 1, 395-400.   | 7.1 | 2         |
| 29 | Observer variation is not eliminated by standardised analysis of dobutamine-atropine stress echocardiography. <i>International Journal of Cardiovascular Imaging</i> , 2002, 18, 169-179.  | 0.6 | 2         |
| 30 | Restenosis in coronary bare metal stents. Importance of time to follow-up: A comparison of coronary angiograms 6 months and 4 years after implantation. <i>Scandinavian Cardiovascular Journal</i> , 2009, 43, 87-93.  | 1.2 | 2         |
| 31 | A mismatch index based on the difference between measured left ventricular ejection fraction and that estimated by infarct size at three months following reperfused acute myocardial infarction. <i>Journal of Electrocardiology</i> , 2014, 47, 191-196.   | 0.9 | 2         |
| 32 | The importance of Î²2-agonists in myocardial infarction: Findings from the Eastern Danish Heart Registry. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2016, 5, 551-559.   | 1.0 | 2         |
| 33 | Longitudinal shortening of subepicardial myocytes in severe ischaemic cardiomyopathy: insights from gadolinium contrast cardiac magnetic resonance imaging. <i>ESC Heart Failure</i> , 2017, 4, 670-674.   | 3.1 | 1         |
| 34 | Bleeding Episodes in Patients With Non-ST-Segment Elevation Acute Coronary Syndrome Undergoing Very Early Versus Standard Care Invasive Examination (from the Very Early vs Deferred Invasive) <i>TJ ETQq0 0 0 rgBT /Overlock 10 Tf 50 14</i><br>170, 10-16.   | 1.6 | 1         |