

Henning KelbÄk

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1577826/publications.pdf>

Version: 2024-02-01

34
papers

1,270
citations

567281

15
h-index

377865

34
g-index

34
all docs

34
docs citations

34
times ranked

2207
citing authors

#	ARTICLE	IF	CITATIONS
1	Bleeding Episodes in Patients With Non-ST-Segment Elevation Acute Coronary Syndrome Undergoing Very Early Versus Standard Care Invasive Examination (from the Very Early vs Deferred Invasive) Trial. <i>Circulation</i> , 2017, 135, 10-16.	1.6	10
2	Impact of age on reperfusion success and long-term prognosis in ST-segment elevation myocardial infarction – A cardiac magnetic resonance imaging study. <i>IJC Heart and Vasculature</i> , 2021, 33, 100731.	1.1	4
3	Subacute cardiac rubidium-82 positron emission tomography (82Rb-PET) to assess myocardial area at risk, final infarct size, and myocardial salvage after STEMI. <i>Journal of Nuclear Cardiology</i> , 2018, 25, 970-981.	2.1	6
4	Early Versus Standard Care Invasive Examination and Treatment of Patients With Non-ST-Segment Elevation Acute Coronary Syndrome. <i>Circulation</i> , 2018, 138, 2741-2750.	1.6	168
5	Left Ventricular Hypertrophy Is Associated With Increased Infarct Size and Decreased Myocardial Salvage in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	39
6	Association Between Early Q Waves and Reperfusion Success in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention. <i>Circulation: Cardiovascular Interventions</i> , 2017, 10, .	3.9	10
7	Fractional Flow Reserve-Guided Complete Revascularization Improves the Prognosis in Patients With ST-Segment Elevation Myocardial Infarction and Severe Nonculprit Disease. <i>Circulation: Cardiovascular Interventions</i> , 2017, 10, .	3.9	39
8	Longitudinal shortening of subepicardial myocytes in severe ischaemic cardiomyopathy: insights from gadolinium contrast cardiac magnetic resonance imaging. <i>ESC Heart Failure</i> , 2017, 4, 670-674.	3.1	1
9	Mechanisms of Very Late Drug-Eluting Stent Thrombosis Assessed by Optical Coherence Tomography. <i>Circulation</i> , 2016, 133, 650-660.	1.6	260
10	The importance of Î²2-agonists in myocardial infarction: Findings from the Eastern Danish Heart Registry. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2016, 5, 551-559.	1.0	2
11	The Third DANish Study of Optimal Acute Treatment of Patients with ST-segment Elevation Myocardial Infarction: Ischemic postconditioning or deferred stent implantation versus conventional primary angioplasty and complete revascularization versus treatment of culprit lesion only. <i>American Heart Journal</i> , 2015, 169, 613-621.	2.7	61
12	Stent Thrombosis is the Primary Cause of ST-Segment Elevation Myocardial Infarction following Coronary Stent Implantation: A Five Year Follow-Up of the SORT OUT II Study. <i>PLoS ONE</i> , 2014, 9, e113399.	2.5	8
13	Impact of Acute Hyperglycemia on Myocardial Infarct Size, Area at Risk, and Salvage in Patients With STEMI and the Association With Exenatide Treatment: Results From a Randomized Study. <i>Diabetes</i> , 2014, 63, 2474-2485.	0.6	59
14	Differential clinical outcomes after 1 year versus 5 years in a randomised comparison of zotarolimus-eluting and sirolimus-eluting coronary stents (the SORT OUT III study): a multicentre, open-label, randomised superiority trial. <i>Lancet</i> , 2014, 383, 2047-2056.	13.7	96
15	A mismatch index based on the difference between measured left ventricular ejection fraction and that estimated by infarct size at three months following reperfused acute myocardial infarction. <i>Journal of Electrocardiology</i> , 2014, 47, 191-196.	0.9	2
16	Bioluminescence-Eluting Stents With Biodegradable Polymer Versus Bare-Metal Stents in Acute Myocardial Infarction. <i>Circulation: Cardiovascular Interventions</i> , 2014, 7, 355-364.	3.9	56
17	Final infarct size measured by cardiovascular magnetic resonance in patients with ST elevation myocardial infarction predicts long-term clinical outcome: an observational study. <i>European Heart Journal Cardiovascular Imaging</i> , 2013, 14, 387-395.	1.2	124
18	Influence of pre-infarction angina, collateral flow, and pre-procedural TIMI flow on myocardial salvage index by cardiac magnetic resonance in patients with ST-segment elevation myocardial infarction. <i>European Heart Journal Cardiovascular Imaging</i> , 2012, 13, 433-443.	1.2	48

#	ARTICLE	IF	CITATIONS
19	Comparison of Selvester QRS score with magnetic resonance imaging measured infarct size in patients with ST elevation myocardial infarction. <i>Journal of Electrocardiology</i> , 2012, 45, 414-419.	0.9	20
20	Zotarolimus-eluting vs. sirolimus-eluting coronary stents in patients with and without acute coronary syndromes: a SORT OUT III substudy. <i>European Journal of Clinical Investigation</i> , 2012, 42, 1047-1054.	3.4	10
21	No Major Differences in 30-Day Outcomes in High-Risk Patients Randomized to Off-Pump Versus On-Pump Coronary Bypass Surgery. <i>Circulation</i> , 2010, 121, 498-504.	1.6	140
22	Graft patency after off-pump versus on-pump coronary artery surgery in high-risk patients. <i>Scandinavian Cardiovascular Journal</i> , 2010, 44, 161-167.	1.2	17
23	Restenosis in coronary bare metal stents. Importance of time to follow-up: A comparison of coronary angiograms 6 months and 4 years after implantation. <i>Scandinavian Cardiovascular Journal</i> , 2009, 43, 87-93.	1.2	2
24	Scintigraphic evaluation of routine filterwire distal protection in percutaneous coronary intervention for acute ST-segment elevation myocardial infarction: a randomized controlled trial. <i>Journal of Nuclear Cardiology</i> , 2009, 16, 784-791.	2.1	10
25	One-year results of total arterial revascularization vs. conventional coronary surgery: CARRPO trial. <i>European Heart Journal</i> , 2008, 30, 1005-1011.	2.2	22
26	Absolute quantitation of left ventricular wall and cavity parameters using ECG-gated PET. <i>Journal of Nuclear Cardiology</i> , 2004, 11, 38-46.	2.1	17
27	Low whole-body insulin sensitivity in patients with ischaemic heart disease is associated with impaired myocardial glucose uptake predictive of poor outcome after revascularisation. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2002, 29, 991-998.	6.4	11
28	Variability of insulin-stimulated myocardial glucose uptake in healthy elderly subjects. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2002, 29, 1600-1607.	6.4	4
29	Between observer variation is not eliminated by standardised analysis of dobutamine-atropine stress echocardiography. <i>International Journal of Cardiovascular Imaging</i> , 2002, 18, 169-179.	0.6	2
30	Regional myocardial oxygen consumption estimated by carbon-11 acetate and positron emission tomography before and after repetitive ischemia.... <i>Journal of Nuclear Cardiology</i> , 2000, 7, 228-234.	2.1	4
31	Comparison of methods of fractional area change for detection of regional left ventricular dysfunction. <i>International Journal of Cardiovascular Imaging</i> , 2000, 16, 257-266.	0.6	6
32	Amlodipine reduces myocardial ischaemia during exercise without compromising left ventricular function in patients with silent ischaemia: a randomised, double-blind, placebo-controlled study. <i>European Journal of Heart Failure</i> , 1999, 1, 395-400.	7.1	2
33	Pre- and afterload reduction in chronic mitral regurgitation: a double-blind randomized placebo-controlled trial of the acute and 2 weeks' effect of nifedipine or isosorbide dinitrate treatment on left ventricular function and the severity of mitral regurg. <i>British Journal of Clinical Pharmacology</i> , 1996, 41, 493-497.	2.4	9
34	Coronary and Skeletal Muscle Enzyme Changes during a 14 km Run. <i>Acta Medica Scandinavica</i> , 1988, 224, 183-186.	0.0	10