

# Volker Daniel

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1602578/publications.pdf>

Version: 2024-02-01

136  
papers

2,822  
citations

186265

28  
h-index

233421

45  
g-index

138  
all docs

138  
docs citations

138  
times ranked

3181  
citing authors

#	ARTICLE	IF	CITATIONS
1	Predicting neurological recovery after traumatic spinal cord injury by time-resolved analysis of monocyte subsets. <i>Brain</i> , 2021, 144, 3159-3174.	7.6	9
2	Higher CD19+CD25+ Bregs are independently associated with better graft function in renal transplant recipients. <i>BMC Nephrology</i> , 2021, 22, 180.	1.8	5
3	Relationship of transitional regulatory B and regulatory T cells and immunosuppressive drug doses in stable renal transplant recipients. <i>Immunity, Inflammation and Disease</i> , 2021, 9, 1252-1271.	2.7	3
4	CD4+CD25+CD127-Foxp3+ and CD8+CD28- Tregs in Renal Transplant Recipients: Phenotypic Patterns, Association With Immunosuppressive Drugs, and Interaction With Effector CD8+ T Cells and CD19+IL-10+ Bregs. <i>Frontiers in Immunology</i> , 2021, 12, 716559.	4.8	5
5	Phase 2 Trial of Oncolytic H-1 Parvovirus Therapy Shows Safety and Signs of Immune System Activation in Patients With Metastatic Pancreatic Ductal Adenocarcinoma. <i>Clinical Cancer Research</i> , 2021, 27, 5546-5556.	7.0	22
6	Fatal late-onset CAR T-cell-mediated encephalitis after axicabtagene-ciloleucel in a patient with large B-cell lymphoma. <i>Blood Advances</i> , 2021, 5, 3789-3793.	5.2	10
7	Selenium-Binding Protein 1 (SELENBP1) as Biomarker for Adverse Clinical Outcome After Traumatic Spinal Cord Injury. <i>Frontiers in Neuroscience</i> , 2021, 15, 680240.	2.8	6
8	LIPUS vs. reaming in non-union treatment: Cytokine expression course as a tool for evaluation and differentiation of non-union therapy. <i>Journal of Orthopaedics</i> , 2020, 17, 208-214.	1.3	0
9	Assessment of CAR T Cell Frequencies in Axicabtagene Ciloleucel and Tisagenlecleucel Patients Using Duplex Quantitative PCR. <i>Cancers</i> , 2020, 12, 2820.	3.7	13
10	Severe underquantification of HIV-1 group O isolates by major commercial PCR-based assays. <i>Clinical Microbiology and Infection</i> , 2020, 26, 1688.e1-1688.e7.	6.0	2
11	Improving antibody-based therapies by chemical engineering of antibodies with multimeric cell-penetrating peptides for elevated intracellular delivery. <i>Journal of Controlled Release</i> , 2020, 322, 200-208.	9.9	30
12	Two of a kind? Immunological and clinical risk factors differ between recurrent implantation failure and recurrent miscarriage. <i>Journal of Reproductive Immunology</i> , 2020, 141, 103166.	1.9	16
13	NK cell subsets in idiopathic recurrent miscarriage and renal transplant patients. <i>Journal of Reproductive Immunology</i> , 2020, 138, 103098.	1.9	5
14	Phase I trial of donor-derived modified immune cell infusion in kidney transplantation. <i>Journal of Clinical Investigation</i> , 2020, 130, 2364-2376.	8.2	29
15	Circulating NKG2A <sup>hi</sup> NKG2D <sup>+</sup> CD56dimCD16 <sup>+</sup> Natural Killer (NK) Cells as Mediators of Functional Immunosurveillance in Kidney Transplant Recipients. <i>Annals of Transplantation</i> , 2020, 25, e925162.	0.9	1
16	CXCR5+IFN- $\gamma$ +CD8 <sup>+</sup> T Lymphocytes as a Potential Inhibitor of DSA Formation in Renal Transplant Recipients. <i>Transplantation</i> , 2020, 104, 2264-2265.	1.0	0
17	Patients with idiopathic recurrent miscarriage have abnormally high TGF- $\beta$ blood NK, NKT and T cells in the presence of abnormally low TGF- $\beta$ plasma levels. <i>BMC Immunology</i> , 2019, 20, 10.	2.2	17
18	Association of pre- and early post-transplant serum amino acids and metabolites of amino acids and liver transplant outcome. <i>Transplant Immunology</i> , 2018, 46, 42-48.	1.2	2

#	ARTICLE	IF	CITATIONS
19	Changes of NK cell subsets with time post-transplant in peripheral blood of renal transplant recipients. <i>Transplant Immunology</i> , 2018, 49, 59-71.	1.2	7
20	Chemokine analysis as a novel diagnostic modality in the early prediction of the outcome of non-union therapy: a matched pair analysis. <i>Journal of Orthopaedic Surgery and Research</i> , 2018, 13, 249.	2.3	2
21	A Preliminary Study of Contrast-Enhanced Ultrasound (CEUS) and Cytokine Expression Analysis (CEA) as Early Predictors for the Outcome of Tibial Non-Union Therapy. <i>Diagnostics</i> , 2018, 8, 55.	2.6	8
22	Cytokines in relation to hCG are significantly altered in asymptomatic women with miscarriage – a pilot study. <i>Reproductive Biology and Endocrinology</i> , 2018, 16, 93.	3.3	13
23	SaO011A PHASE-I CLINICAL TRIAL OF DONOR-DERIVED MIC CELL INFUSION FOR THE INDUCTION OF DONOR-SPECIFIC HYPORESPONSIVENESS AFTER LIVING DONOR KIDNEY TRANSPLANTATION (TOL-1 STUDY). <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i320-i320.	0.7	0
24	Evaluation of matrix metalloproteases as early biomarkers for bone regeneration during the applied Masquelet therapy for non-unions. <i>Injury</i> , 2018, 49, 1732-1738.	1.7	17
25	Increased natural killer cell subsets with inhibitory cytokines and inhibitory surface receptors in patients with recurrent miscarriage and decreased or normal subsets in kidney transplant recipients late post-transplant. <i>Clinical and Experimental Immunology</i> , 2018, 193, 241-254.	2.6	17
26	Association of Graft Effluent Parameters with Donor Body Mass Index, Graft Quality, and Post-Transplant Events. <i>Annals of Transplantation</i> , 2018, 23, 136-143.	0.9	1
27	Induction of Donor-Specific Immune Tolerance with Clinical MIC Cell Infusion – a Phase I Study (TOL-1). <i>Blood</i> , 2018, 132, 4539-4539.	1.4	0
28	Exploratory study to suggest the possibility of MMP-8 and MMP-9 serum levels as early markers for remission after traumatic spinal cord injury. <i>Spinal Cord</i> , 2017, 55, 8-15.	1.9	21
29	Association of peripheral NK cell counts with Helios+IFN- $\gamma$ Tregs in patients with good long-term renal allograft function. <i>Clinical and Experimental Immunology</i> , 2017, 188, 467-479.	2.6	16
30	Pre-Pregnancy Levels of Peripheral Natural Killer Cells as Markers for Immunomodulatory Treatment in Patients with Recurrent Miscarriage. <i>Archivum Immunologiae Et Therapiae Experimentalis</i> , 2017, 65, 339-346.	2.3	16
31	CCL-2 as a possible early marker for remission after traumatic spinal cord injury. <i>Spinal Cord</i> , 2017, 55, 1002-1009.	1.9	20
32	Endothelial precursor cell cross-match using Tie2-enriched spleen cells. <i>Clinical Transplantation</i> , 2017, 31, e13118.	1.6	1
33	Oncolytic H-1 Parvovirus Shows Safety and Signs of Immunogenic Activity in a First Phase I/IIa Glioblastoma Trial. <i>Molecular Therapy</i> , 2017, 25, 2620-2634.	8.2	199
34	Time-course of plasma inflammatory mediators in a rat model of brain death. <i>Transplant Immunology</i> , 2017, 43-44, 21-26.	1.2	3
35	Low utility of serum 25-hydroxyvitamin D 3 and 1, 25-dihydroxyvitamin D 3 in predicting peripheral Treg and Th17 cell counts in ESRD and renal transplant patients. <i>Transplant Immunology</i> , 2017, 43-44, 3-10.	1.2	3
36	The “killer cell story” in recurrent miscarriage: Association between activated peripheral lymphocytes and uterine natural killer cells. <i>Journal of Reproductive Immunology</i> , 2017, 119, 9-14.	1.9	57

#	ARTICLE	IF	CITATIONS
37	Decreased NK cell immunity in kidney transplant recipients late post-transplant and increased NK-cell immunity in patients with recurrent miscarriage. PLoS ONE, 2017, 12, e0186349.	2.5	14
38	A non-controlled, single arm, open label, phase II study of intravenous and intratumoral administration of ParvOryx in patients with metastatic, inoperable pancreatic cancer: ParvOryx02 protocol. BMC Cancer, 2017, 17, 576.	2.6	36
39	Helios expression and Foxp3 TSDR methylation of IFN $\gamma$ + and IFN $\gamma$ - Treg from kidney transplant recipients with good long-term graft function. PLoS ONE, 2017, 12, e0173773.	2.5	11
40	The treatment of nonunions with application of BMP-7 increases the expression pattern for angiogenic and inflammable cytokines: a matched pair analysis. Journal of Inflammation Research, 2016, Volume 9, 155-165.	3.5	18
41	Clinical relevance of preformed IgG and IgM antibodies against donor endothelial progenitor cells in recipients of living donor kidney grafts. Clinical Transplantation, 2016, 30, 124-130.	1.6	8
42	Early post-transplant neopterin associated with one year survival and bacteremia in liver transplant recipients. Human Immunology, 2016, 77, 115-120.	2.4	14
43	Early post-operative acute phase response in patients with early graft dysfunction is predictive of 6-month and 12-month mortality in liver transplant recipients. Human Immunology, 2016, 77, 952-960.	2.4	18
44	IFN $\gamma$ + and IFN $\gamma$ <sup>hi</sup> Treg subsets with stable and unstable Foxp3 expression in kidney transplant recipients with good long-term graft function. Transplant Immunology, 2016, 39, 1-9.	1.2	5
45	Low-dose oral cholecalciferol is associated with higher numbers of Helios+ and total Tregs than oral calcitriol in renal allograft recipients: an observational study. BMC Pharmacology & Toxicology, 2016, 17, 24.	2.4	4
46	IL-23 plasma level is strongly associated with CMV status and reactivation of CMV in renal transplant recipients. BMC Immunology, 2016, 17, 35.	2.2	11
47	Immunosuppressive drugs affect induction of IFN $\gamma$ + Treg in vitro. Human Immunology, 2016, 77, 146-152.	2.4	13
48	Serum levels of chemokines CCL4 and CCL5 in cirrhotic patients indicate the presence of hepatocellular carcinoma. British Journal of Cancer, 2015, 113, 756-762.	6.4	49
49	Immune profiling in patients with recurrent miscarriage. Journal of Reproductive Immunology, 2015, 108, 136-141.	1.9	23
50	Posttraumatic Inflammation as a Key to Neuroregeneration after Traumatic Spinal Cord Injury. International Journal of Molecular Sciences, 2015, 16, 7900-7916.	4.1	52
51	Association of low serum TGF- $\beta$ 2 level in hantavirus infected patients with severe disease. BMC Immunology, 2015, 16, 19.	2.2	4
52	Patients with idiopathic recurrent miscarriage show higher levels of DR+ activated T-cells that are less responsive to mitogens. Journal of Reproductive Immunology, 2015, 112, 82-87.	1.9	17
53	IFN $\gamma$ <sup>hi</sup> Treg in-vivo and in-vitro represent both activated nTreg and peripherally induced aTreg and remain phenotypically stable in-vitro after removal of the stimulus. BMC Immunology, 2015, 16, 45.	2.2	21
54	Immunonutrition - the influence of early postoperative glutamine supplementation in enteral/parenteral nutrition on immune response, wound healing and length of hospital stay in multiple trauma patients and patients after extensive surgery. GMS Interdisciplinary Plastic and Reconstructive Surgery DGPW, 2015, 4, Doc15.	0.1	11

#	ARTICLE	IF	CITATIONS
55	Role of Human Corneal Endothelial Cells in T-Cell-Mediated Alloimmune Attack In Vitro. , 2014, 55, 1213.		20
56	Interferon-Gamma Producing Regulatory T Cells as a Diagnostic and Therapeutic Tool in Organ Transplantation. International Reviews of Immunology, 2014, 33, 195-211.	3.3	41
57	Differences in the Induction of Induced Human CD4+ CD25+ FoxP3+ T-Regulatory Cells and CD3+ CD8+ CD28 <sup>hi</sup> T-Suppressor Cells Subset Phenotypes In Vitro: Comparison of Phorbol 12-Myristate 13-Acetate/Ionomycin and Phytohemagglutinin Stimulation. Transplantation Proceedings, 2013, 45, 1822-1831.	0.6	15
58	HIV-Specific CD8+ T Lymphocytes in Blood of Long-Term HIV-Infected Hemophilia Patients. BioResearch Open Access, 2013, 2, 399-411.	2.6	0
59	CD4+CD25+Foxp3+IFN $\gamma$ +CD178+ human induced Treg (iTreg) contribute to suppression of alloresponses by apoptosis of responder cells. Human Immunology, 2013, 74, 151-162.	2.4	16
60	Plasticity and Overlap of In Vitro-Induced Regulatory T-Cell Markers in Healthy Humans. Transplantation Proceedings, 2013, 45, 1816-1821.	0.6	8
61	Increased serum levels of quinolinic acid indicate enhanced severity of hepatic dysfunction in patients with liver cirrhosis. Human Immunology, 2013, 74, 60-66.	2.4	14
62	Strong association of phenylalanine and tryptophan metabolites with activated cytomegalovirus infection in kidney transplant recipients. Human Immunology, 2012, 73, 186-192.	2.4	21
63	CD4+CD25+Foxp3+IFN $\gamma$ + Treg are immunosuppressive in vitro and increase with intensity of the alloresponse in pretransplant MLC. Transplant Immunology, 2012, 27, 114-121.	1.2	18
64	In-vitro inhibition of IFN $\gamma$ + iTreg mediated by monoclonal antibodies against cell surface determinants essential for iTreg function. BMC Immunology, 2012, 13, 47.	2.2	8
65	Whole-Exome Sequencing Links CARD11 Inactivation with SCID. Blood, 2012, 120, 258-258.	1.4	0
66	CD4+CD25+Foxp3+IFN $\gamma$ + human induced T regulatory cells are induced by interferon $\gamma$ and suppress alloresponses nonspecifically. Human Immunology, 2011, 72, 699-707.	2.4	31
67	Cytokine expression during early and late phase of acute Puumala hantavirus infection. BMC Immunology, 2011, 12, 65.	2.2	59
68	Association of high IFN $\gamma$ plasma levels with low B $\alpha$ cell counts in renal transplant recipients with stable long-term graft function. Clinical Transplantation, 2010, 24, 281-289.	1.6	5
69	Increased Peripheral Blood Leukocyte Subsets with Regulatory Phenotype in Clinically Stable Long-Term HIV-Infected Hemophilia Patients on HAART May Be Beneficial and Contribute to a Decrease in Autoimmunity. Viral Immunology, 2010, 23, 87-97.	1.3	1
70	Increased pretransplantation plasma kynurenine levels do not protect from but predict acute kidney allograft rejection. Human Immunology, 2010, 71, 1067-1072.	2.4	15
71	Clinical Relevance of Immune Monitoring in Solid Organ Transplantation. International Reviews of Immunology, 2009, 28, 155-184.	3.3	4
72	Association of IL-12+ DC with High CD3+CD4-DR+ Lymphocyte Counts in Long-term HIV-infected Hemophilia Patients With Clinically Stable Disease. Journal of Clinical Immunology, 2008, 28, 58-72.	3.8	5

#	ARTICLE	IF	CITATIONS
73	Observational support for an immunoregulatory role of CD3 <sup>+</sup> CD4 <sup>+</sup> CD25 <sup>+</sup> IFN- $\gamma$ <sup>+</sup> blood lymphocytes in kidney transplant recipients with good long-term graft outcome. <i>Transplant International</i> , 2008, 21, 646-660.	1.6	56
74	Soluble IL-2 receptor and tumour necrosis factor- $\alpha$ in plasma of haemophilia patients infected with HIV. <i>Clinical and Experimental Immunology</i> , 2008, 87, 287-292.	2.6	29
75	Association of T cell and macrophage dysfunction with surface gp120-immunoglobulin-complement complexes in HIV-infected patients. <i>Clinical and Experimental Immunology</i> , 2008, 93, 152-156.	2.6	27
76	Dendritische Zellen und Immunsuppression nach Organtransplantation / Dendritic cells and immunosuppression after organ transplantation. <i>Laboratoriums Medizin</i> , 2008, 32, 131-139.	0.6	3
77	Dendritic cells and immunosuppression after organ transplantation 1. <i>Laboratoriums Medizin</i> , 2008, 32, -.	0.6	1
78	Normal or Even Increased Dendritic Cell and Peripheral Blood Lymphocyte Subsets With Regulatory Phenotype in Clinically Stable Long-Term HIV-Infected Patients With Hemophilia on Highly Active Antiretroviral Therapy. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2008, 47, 1-15.	2.1	29
79	Dysregulated Cytokine Responses During Cytomegalovirus Infection in Renal Transplant Recipients. <i>Transplantation</i> , 2008, 86, 275-285.	1.0	27
80	<i>Short Communication:</i> Decreasing Soluble CD30 and Increasing IFN- $\gamma$ Plasma Levels Are Indicators of Effective Highly Active Antiretroviral Therapy. <i>AIDS Research and Human Retroviruses</i> , 2007, 23, 886-890.	1.1	22
81	Evidence for IFN- $\gamma$ up- and IL-4 downregulation late post-transplant in patients with good kidney graft outcome. <i>Clinical Transplantation</i> , 2007, 21, 449-459.	1.6	19
82	Post-Transplant sCD30 and Neopterin as Predictors of Chronic Allograft Nephropathy: Impact of Different Immunosuppressive Regimens. <i>American Journal of Transplantation</i> , 2006, 6, 1865-1874.	4.7	74
83	Association of Circulating Interleukin (IL)-12 $\alpha$ and IL-10 $\alpha$ -Producing Dendritic Cells with Time Posttransplant, Dose of Immunosuppression, and Plasma Cytokines in Renal-Transplant Recipients. <i>Transplantation</i> , 2005, 79, 1498-1506.	1.0	27
84	Effectivity of a T-Cell-Adapted Induction Therapy With Anti-Thymocyte Globulin (Sangstat). <i>Journal of Heart and Lung Transplantation</i> , 2005, 24, 708-713.	0.6	26
85	Evidence for autoantibody-induced CD4 depletion mediated by apoptotic and non-apoptotic mechanisms in HIV-positive long-term surviving haemophilia patients. <i>Clinical and Experimental Immunology</i> , 2004, 135, 94-104.	2.6	13
86	Dissociation of CD4 <sup>+</sup> cell counts from viral load and association with immune complexes in HIV+ hemophilia patients. <i>Immunology Letters</i> , 2004, 91, 23-32.	2.5	3
87	Evaluation of posttransplantation soluble CD30 for diagnosis of acute renal allograft rejection1. <i>Transplantation</i> , 2003, 75, 421-423.	1.0	81
88	Associations of Dichlorodiphenyltrichloroethane (DDT) 4.4 and Dichlorodiphenyldichloroethylene (DDE) 4.4 Blood Levels with Plasma IL-4. <i>Archives of Environmental Health</i> , 2002, 57, 541-547.	0.4	59
89	Association of Elevated Blood Levels of Pentachlorophenol (PCP) with Cellular and Humoral Immunodeficiencies. <i>Archives of Environmental Health</i> , 2001, 56, 77-83.	0.4	34
90	Associations of blood levels of PCB, HCHS, and HCB with numbers of lymphocyte subpopulations, in vitro lymphocyte response, plasma cytokine levels, and immunoglobulin autoantibodies.. <i>Environmental Health Perspectives</i> , 2001, 109, 173-178.	6.0	80

#	ARTICLE	IF	CITATIONS
91	Association of immune complexes and plasma viral load with CD4+ cell depletion, CD8+DR+ and CD16+ cell counts in HIV+ hemophilia patients. Implications for the immunopathogenesis of HIV-induced CD4+ lymphocyte depletion. <i>Immunology Letters</i> , 2001, 76, 69-78.	2.5	7
92	Increased Soluble Fas in HIV-Infected Hemophilia Patients with CD4+and CD8+Cell Count Increases and Viral Load and Immune Complex Decreases. <i>AIDS Research and Human Retroviruses</i> , 2001, 17, 329-335.	1.1	6
93	Impact of Apolipoprotein(a) Phenotypes on Long-Term Renal Transplant Survival. <i>Journal of the American Society of Nephrology: JASN</i> , 2001, 12, 1052-1058.	6.1	11
94	Immunomodulatory Effects of Acupuncture in the Treatment of Allergic Asthma: A Randomized Controlled Study. <i>Journal of Alternative and Complementary Medicine</i> , 2000, 6, 519-525.	2.1	114
95	Reduction of viral load and immune complex load on CD4+ lymphocytes as a consequence of highly active antiretroviral treatment (HAART) in HIV-infected hemophilia patients. <i>Immunology Letters</i> , 1999, 69, 283-289.	2.5	18
96	CD4 depletion in HIV-infected haemophilia patients is associated with rapid clearance of immune complex-coated CD4+lymphocytes. <i>Clinical and Experimental Immunology</i> , 1999, 115, 477-484.	2.6	8
97	P-glycoprotein expression is not a useful predictor of acute or chronic kidney graft rejection. <i>Transplant International</i> , 1999, 12, 10-17.	1.6	11
98	Predictive indicators of rejection or infection in renal transplant patients. <i>Transplantation Proceedings</i> , 1999, 31, 1364-1365.	0.6	8
99	P-glycoprotein expression is not a useful predictor of acute or chronic kidney graft rejection. <i>Transplant International</i> , 1999, 12, 10-17.	1.6	7
100	Circulating interleukin-1 receptor antagonist (IL-1RA) serum levels in patients undergoing orthotopic heart transplantation. <i>Transplant International</i> , 1998, 11, 443-448.	1.6	3
101	Superior 3-year kidney graft function in patients with impaired pretransplant Th2 responses. <i>Transplant International</i> , 1998, 11, S350-S356.	1.6	9
102	HIV-induced IL-6/IL-10 dysregulation of CD4 cells is associated with defective B cell help and autoantibody formation against CD4 cells. <i>Clinical and Experimental Immunology</i> , 1998, 111, 20-29.	2.6	16
103	Association of viral load in plasma samples of HIV-infected hemophilia patients with autoantibodies and gp120-containing immune complexes on CD4+ lymphocytes. <i>Immunology Letters</i> , 1998, 60, 179-187.	2.5	7
104	Immunological effects of acupuncture on allergic asthma. <i>Focus on Alternative and Complementary Therapies</i> , 1998, 3, 187-187.	0.1	0
105	T-lymphocyte populations, cytokines and other growth factors in serum and urine of children with idiopathic nephrotic syndrome. <i>Clinical Nephrology</i> , 1997, 47, 289-97.	0.7	69
106	Identification of complement activation sites in human immunodeficiency virus type-1 glycoprotein gp120. <i>Blood</i> , 1996, 87, 2329-2336.	1.4	29
107	CD8+ Lymphocyte Decrease in HIV Disease: Association with Anti-CD4+ but Not with Anti-CD8+ Lymphocyte Autoantibodies. <i>Vox Sanguinis</i> , 1996, 70, 86-91.	1.5	10
108	Association of T cell dysfunction with the presence of IgG autoantibodies on CD4+ lymphocytes in haemophilia patients; results of a 10-year study. <i>Clinical and Experimental Immunology</i> , 1996, 104, 4-10.	2.6	21

#	ARTICLE	IF	CITATIONS
109	In vitro Cytokine Treatment of B Cell Defects in HIV-Infected Hemophilia Patients. <i>Vox Sanguinis</i> , 1995, 69, 27-37.	1.5	3
110	Sequential occurrence of IgM, IgM/IgG, and gp120-IgM/IgG complement complexes on CD4+ lymphocytes in relation to CD4+ blood lymphocyte depletion in HIV+ hemophilia patients: results of a 10-year study. <i>Immunology Letters</i> , 1995, 47, 97-102.	2.5	20
111	Impaired In-Vitro Lymphocyte Responses in Patients with Elevated Pentachlorophenol (PCP) Blood Levels. <i>Archives of Environmental Health</i> , 1995, 50, 287-292.	0.4	57
112	Cytokine monitoring of infection and rejection in renal transplant recipients. <i>Transplantation Proceedings</i> , 1995, 27, 884-6.	0.6	18
113	Isotypes and IgG Subclasses of Anti-Fab Antibodies in Human Immunodeficiency Virus-Infected Hemophilia Patients. <i>Vox Sanguinis</i> , 1994, 66, 37-45.	1.5	17
114	Complement activation by recombinant HIV-1 glycoprotein gp120. <i>Journal of Immunology</i> , 1994, 152, 6028-34.	0.8	42
115	Autoantibodies against CD4+ lymphocytes in HIV-infected hemophilia patients. <i>Acta Microbiologica Et Immunologica Hungarica</i> , 1994, 41 Suppl, 11-5.	0.8	0
116	CD4+ Lymphocyte Depletion in HIV-Infected Patients is Associated with gp120-Immunoglobulin-Complement Attachment to CD4+ Cells. <i>Vox Sanguinis</i> , 1993, 64, 31-36.	1.5	21
117	Molecular Mimicry between HIV-1 and Antigen Receptor Molecules: A Clue to the Pathogenesis of AIDS. <i>Vox Sanguinis</i> , 1993, 65, 10-17.	1.5	54
118	CD4+ Lymphocyte Depletion in HIV-Infected Patients is Associated with gp120-Immunoglobulin-Complement Attachment to CD4+ Cells. <i>Vox Sanguinis</i> , 1993, 64, 31-36.	1.5	22
119	Plasma GM-CSF, IL-6, and IL-3 monitoring allows differentiation between infection and rejection in some renal transplant recipients: preliminary results of a retrospective study. <i>Transplantation Proceedings</i> , 1993, 25, 893-6.	0.6	2
120	Striking inverse association of IgG-anti-Fab gamma antibodies and CD4 cell counts in patients with acquired immunodeficiency syndrome (AIDS)/AIDS-related complex. <i>Blood</i> , 1992, 79, 954-957.	1.4	33
121	Autoantibodies in HIV-Infected Hemophilia Patients against Different Epitopes on CD4+Lymphocytes and Recombinant CD4. <i>Vox Sanguinis</i> , 1992, 62, 39-44.	1.5	10
122	Anti-IgG Autoantibodies in HIV-Infected Hemophilia Patients. <i>Vox Sanguinis</i> , 1992, 62, 224-229.	1.5	31
123	Striking inverse association of IgG-anti-Fab gamma antibodies and CD4 cell counts in patients with acquired immunodeficiency syndrome (AIDS)/AIDS-related complex. <i>Blood</i> , 1992, 79, 954-957.	1.4	36
124	Preliminary evidence that monitoring of plasma granulocyte-macrophage colony-stimulating factor may be helpful to differentiate between infection and rejection in renal transplant patients. <i>Transplantation Proceedings</i> , 1992, 24, 2770-2.	0.6	2
125	Autoantibodies against CD4 cells are associated with CD4 helper defects in human immunodeficiency virus-infected patients. <i>Blood</i> , 1991, 77, 133-140.	1.4	62
126	Autoantibodies against CD4 cells are associated with CD4 helper defects in human immunodeficiency virus-infected patients. <i>Blood</i> , 1991, 77, 133-140.	1.4	0



#	ARTICLE	IF	CITATIONS
127	Autoantibodies against CD4 cells are associated with CD4 helper defects in human immunodeficiency virus-infected patients. <i>Blood</i> , 1991, 77, 133-40.	1.4	18
128	Autoantibodies against CD4 and CD8 Positive T Lymphocytes in HIV-Infected Hemophilia Patients. <i>Vox Sanguinis</i> , 1989, 57, 172-176.	1.5	38
129	Lymphocyte autoantibodies and alloantibodies in HIV-positive haemophilia patients. <i>Clinical and Experimental Immunology</i> , 1989, 75, 178-83.	2.6	48
130	Non-complement-fixing antibodies as indicators for impending renal allograft rejection. <i>Transplantation Proceedings</i> , 1989, 21, 702-3.	0.6	6
131	Correlation of in vitro Immune Defects with Impaired Gamma Interferon Response in Human-Immunodeficiency-Virus-Infected Individuals. <i>Vox Sanguinis</i> , 1988, 54, 92-95.	1.5	13
132	Correlation of in vitro Immune Defects with Impaired Gamma Interferon Response in Human-Immunodeficiency-Virus-Infected Individuals. <i>Vox Sanguinis</i> , 1988, 54, 92-95.	1.5	10
133	Suppression of donor-reactive T cells by pretransplant sera of renal transplant recipients. <i>Transplantation Proceedings</i> , 1988, 20, 291-2.	0.6	10
134	Suppression and stimulation of T cell clones of renal transplant recipients by autologous post-transplant sera: a role for regulatory antibodies. <i>Transplantation Proceedings</i> , 1987, 19, 4210-3.	0.6	1
135	ASSESSMENT OF PLASMA NEOPTERIN IN CLINICAL KIDNEY TRANSPLANTATION. <i>Transplantation</i> , 1986, 41, 454-458.	1.0	70
136	Correlation of Immune Defects in Hemophilia with HTLV-III Antibody Titers. <i>Vox Sanguinis</i> , 1986, 51, 35-39.	1.5	28