

Tim J Croudace

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1603358/publications.pdf>

Version: 2024-02-01

130
papers

8,686
citations

38742

50
h-index

48315

88
g-index

131
all docs

131
docs citations

131
times ranked

10474
citing authors

#	ARTICLE	IF	CITATIONS
1	On Dimensionality, Measurement Invariance, and Suitability of Sum Scores for the PHQ-9 and the GAD-7. <i>Assessment</i> , 2022, 29, 355-366.	3.1	38
2	Determinants of patient-reported outcome trajectories and symptomatic recovery in Improving Access to Psychological Therapies (IAPT) services. <i>Psychological Medicine</i> , 2022, 52, 3231-3240.	4.5	4
3	Neurological Signs at the First Psychotic Episode as Correlates of Long-Term Outcome: Results From the AESOP-10 Study. <i>Schizophrenia Bulletin</i> , 2021, 47, 118-127.	4.3	20
4	Diagnostic stability and outcome after first episode psychosis. <i>Journal of Mental Health</i> , 2021, 30, 104-112.	1.9	9
5	An item response theory analysis of an item pool for the recovering quality of life (ReQoL) measure. <i>Quality of Life Research</i> , 2021, 30, 267-276.	3.1	6
6	Symptom remission at 12-weeks strongly predicts long-term recovery from the first episode of psychosis. <i>Psychological Medicine</i> , 2020, 50, 1452-1462.	4.5	16
7	Prevalence of and recovery from common mental disorder including psychotic experiences in the UK Primary Care Improving Access to Psychological Therapies (IAPT) Programme. <i>Journal of Affective Disorders</i> , 2020, 272, 84-90.	4.1	27
8	Exploring the item sets of the Recovering Quality of Life (ReQoL) measures using factor analysis. <i>Quality of Life Research</i> , 2019, 28, 1005-1015.	3.1	11
9	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. <i>Programme Grants for Applied Research</i> , 2019, 7, 1-108.	1.0	6
10	Recovering Quality of Life (ReQoL): a new generic self-reported outcome measure for use with people experiencing mental health difficulties. <i>British Journal of Psychiatry</i> , 2018, 212, 42-49.	2.8	141
11	Reduction in adolescent depression after contact with mental health services: a longitudinal cohort study in the UK. <i>Lancet Psychiatry</i> , 2017, 4, 120-127.	7.4	86
12	Ethnicity and long-term course and outcome of psychotic disorders in a UK sample: The AESOP-10 study. <i>British Journal of Psychiatry</i> , 2017, 211, 88-94.	2.8	71
13	Monoamine Oxidase A (MAOA) Gene and Personality Traits from Late Adolescence through Early Adulthood: A Latent Variable Investigation. <i>Frontiers in Psychology</i> , 2017, 8, 1736.	2.1	12
14	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 496.	1.6	15
15	Clinical services for adults with an intellectual disability and epilepsy: A comparison of management alternatives. <i>PLoS ONE</i> , 2017, 12, e0180266.	2.5	13
16	Friendships and Family Support Reduce Subsequent Depressive Symptoms in At-Risk Adolescents. <i>PLoS ONE</i> , 2016, 11, e0153715.	2.5	151
17	Effects of ignoring clustered data structure in confirmatory factor analysis of ordered polytomous items: a simulation study based on PANSS. <i>International Journal of Methods in Psychiatric Research</i> , 2016, 25, 205-219.	2.1	6
18	Protocol for the Mindful Student Study: a randomised controlled trial of the provision of a mindfulness intervention to support university students' well-being and resilience to stress. <i>BMJ Open</i> , 2016, 6, e012300.	1.9	27

#	ARTICLE	IF	CITATIONS
19	An evaluation of computerized adaptive testing for general psychological distress: combining GHQ-12 and Affectometer-2 in an item bank for public mental health research. BMC Medical Research Methodology, 2016, 16, 58.	3.1	17
20	Ten-year outcomes in first episode psychotic major depression patients compared with schizophrenia and bipolar patients. Schizophrenia Research, 2016, 176, 417-422.	2.0	23
21	Evaluation of ethnic disparities in detection of depression and anxiety in primary care during the maternal period: Combined analysis of routine and cohort data. British Journal of Psychiatry, 2016, 208, 453-461.	2.8	46
22	Calibrating well-being, quality of life and common mental disorder items: Psychometric epidemiology in public mental health research. British Journal of Psychiatry, 2016, 209, 162-168.	2.8	53
23	Outcomes following first-episode psychosis – Why we should intervene early in all ages, not only in youth. Australian and New Zealand Journal of Psychiatry, 2016, 50, 1055-1063.	2.3	25
24	Computerized adaptive testing of population psychological distress: simulation-based evaluation of GHQ-30. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 895-906.	3.1	8
25	Biological and psychosocial risk factors for psychotic major depression. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 233-245.	3.1	19
26	Maternal psychological distress in primary care and association with child behavioural outcomes at age three. European Child and Adolescent Psychiatry, 2016, 25, 601-613.	4.7	13
27	Understanding causes of and developing effective interventions for schizophrenia and other psychoses. Programme Grants for Applied Research, 2016, 4, 1-184.	1.0	3
28	Childhood adversity subtypes and depressive symptoms in early and late adolescence. Development and Psychopathology, 2015, 27, 885-899.	2.3	54
29	Diagnostic change 10 years after a first episode of psychosis. Psychological Medicine, 2015, 45, 2757-2769.	4.5	74
30	Ten-Year Outcomes of First-Episode Psychoses in the MRC A†SOP-10 Study. Journal of Nervous and Mental Disease, 2015, 203, 379-386.	1.0	90
31	Clinical effectiveness and cost-effectiveness of tailored intensive liaison between primary and secondary care to identify individuals at risk of a first psychotic illness (the LEGs study): a cluster-randomised controlled trial. Lancet Psychiatry, the, 2015, 2, 984-993.	7.4	47
32	Schizophrenia and other psychotic disorders in Caribbean-born migrants and their descendants in England: systematic review and meta-analysis of incidence rates, 1950–2013. Social Psychiatry and Psychiatric Epidemiology, 2015, 50, 1039-1055.	3.1	54
33	Mood, anxiety and psychotic phenomena measure a common psychopathological factor. Psychological Medicine, 2015, 45, 1483-1493.	4.5	128
34	Factors of psychological distress: clinical value, measurement substance, and methodological artefacts. Social Psychiatry and Psychiatric Epidemiology, 2015, 50, 515-524.	3.1	26
35	Validity of single item responses to short message service texts to monitor depression: an mHealth sub-study of the UK ACUDep trial. BMC Medical Research Methodology, 2015, 15, 56.	3.1	14
36	Mortality in Schizophrenia and Other Psychoses: A 10-Year Follow-up of the Ó”SOP First-Episode Cohort. Schizophrenia Bulletin, 2015, 41, 664-673.	4.3	128

#	ARTICLE	IF	CITATIONS
37	Public mental health: evidenced-based priorities. <i>Lancet, The</i> , 2015, 385, 1472-1475.	13.7	31
38	A comparison of approaches for assessing covariate effects in latent class analysis. <i>Longitudinal and Life Course Studies</i> , 2015, 6, .	0.6	33
39	Heterogeneity in Patient-Reported Outcomes following Low-Intensity Mental Health Interventions: A Multilevel Analysis. <i>PLoS ONE</i> , 2014, 9, e99658.	2.5	16
40	Elevated morning cortisol is a stratified population-level biomarker for major depression in boys only with high depressive symptoms. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 3638-3643.	7.1	97
41	Adolescent Self-control Predicts Midlife Hallucinatory Experiences: 40-Year Follow-up of a National Birth Cohort. <i>Schizophrenia Bulletin</i> , 2014, 40, 1543-1551.	4.3	10
42	Multilevel ordinal factor analysis of the Positive and Negative Syndrome Scale (PANSS). <i>International Journal of Methods in Psychiatric Research</i> , 2014, 23, 25-35.	2.1	12
43	Reappraising the long-term course and outcome of psychotic disorders: the AESOP-10 study. <i>Psychological Medicine</i> , 2014, 44, 2713-2726.	4.5	192
44	Early development, stress and depression across the life course: pathways to depression in a national British birth cohort. <i>Psychological Medicine</i> , 2014, 44, 2845-2854.	4.5	38
45	Postwar environment and long-term mental health problems in former child soldiers in Northern Uganda: the WAYS study. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 425-430.	3.7	28
46	Factor analysis of treatment outcomes from a UK specialist addiction service: Relationship between the Leeds Dependence Questionnaire, Social Satisfaction Questionnaire and 10-item Clinical Outcomes in Routine Evaluation Drug and Alcohol Review, 2014, 33, 643-650.	2.1	7
47	Item bank measurement of depression: will one dimension work?. <i>Journal of Clinical Epidemiology</i> , 2014, 67, 4-6.	5.0	9
48	The influence of different types of war experiences on depression and anxiety in a Ugandan cohort of war-affected youth: the WAYS study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014, 49, 1783-1792.	3.1	19
49	Measuring mental health and wellbeing outcomes for children and adolescents to inform practice and policy: a review of child self-report measures. <i>Child and Adolescent Psychiatry and Mental Health</i> , 2014, 8, 14.	2.5	111
50	Lifetime affect and midlife cognitive function: prospective birth cohort study. <i>British Journal of Psychiatry</i> , 2014, 204, 194-199.	2.8	24
51	Comparison of high and low intensity contact between secondary and primary care to detect people at ultra-high risk for psychosis: study protocol for a theory-based, cluster randomized controlled trial. <i>Trials</i> , 2013, 14, 222.	1.6	10
52	Reliability Generalization for Childhood Autism Rating Scale. <i>Journal of Autism and Developmental Disorders</i> , 2013, 43, 2855-2865.	2.7	29
53	Cohort profile: mental health following extreme trauma in a northern Ugandan cohort of War-Affected Youth Study (The WAYS Study). <i>SpringerPlus</i> , 2013, 2, 300.	1.2	35
54	Rumination, anxiety, depressive symptoms and subsequent depression in adolescents at risk for psychopathology: a longitudinal cohort study. <i>BMC Psychiatry</i> , 2013, 13, 250.	2.6	84

#	ARTICLE	IF	CITATIONS
55	Predictors of human rotation. <i>Laterality</i> , 2013, 18, 265-281.	1.0	10
56	The catechol-O-methyltransferase gene (COMT) and cognitive function from childhood through adolescence. <i>Biological Psychology</i> , 2013, 92, 359-364.	2.2	16
57	The validity of patient- and clinician-rated measures of needs and the therapeutic relationship in psychosis: A pooled analysis. <i>Psychiatry Research</i> , 2013, 209, 711-720.	3.3	7
58	Adolescent self-organization predicts midlife memory in a prospective birth cohort study.. <i>Psychology and Aging</i> , 2013, 28, 958-968.	1.6	18
59	Meta-analytic approaches to determine gender differences in the age-incidence characteristics of schizophrenia and related psychoses. <i>International Journal of Methods in Psychiatric Research</i> , 2013, 22, 36-45.	2.1	19
60	Comparing growth trajectories of risk behaviors from late adolescence through young adulthood: An accelerated design.. <i>Developmental Psychology</i> , 2013, 49, 1732-1738.	1.6	50
61	Psychological Distress during Pregnancy in a Multi-Ethnic Community: Findings from the Born in Bradford Cohort Study. <i>PLoS ONE</i> , 2013, 8, e60693.	2.5	34
62	Who Do You Think Is in Control in Addiction? A Pilot Study on Drug-related Locus of Control Beliefs. <i>Addictive Disorders and Their Treatment</i> , 2012, 11, 195-205.	0.5	16
63	Association between pubertal development and depressive symptoms in girls from a UK cohort. <i>Psychological Medicine</i> , 2012, 42, 2579-2589.	4.5	57
64	Childhood cognitive function and adult psychopathology: associations with psychotic and non-psychotic symptoms in the general population. <i>British Journal of Psychiatry</i> , 2012, 201, 124-130.	2.8	33
65	Factors influencing the costs of epilepsy in adults with an intellectual disability. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2012, 21, 205-210.	2.0	14
66	Mokken scale analysis of mental health and well-being questionnaire item responses: a non-parametric IRT method in empirical research for applied health researchers. <i>BMC Medical Research Methodology</i> , 2012, 12, 74.	3.1	175
67	Incidence of Schizophrenia and Other Psychoses in England, 1950-2009: A Systematic Review and Meta-Analyses. <i>PLoS ONE</i> , 2012, 7, e31660.	2.5	385
68	Comparative Effectiveness of Cognitive Therapies Delivered Face-to-Face or over the Telephone: An Observational Study Using Propensity Methods. <i>PLoS ONE</i> , 2012, 7, e42916.	2.5	60
69	5-HTTLPR and Early Childhood Adversities Moderate Cognitive and Emotional Processing in Adolescence. <i>PLoS ONE</i> , 2012, 7, e48482.	2.5	39
70	Depressive symptoms during adolescence: comparison between epidemiological and high risk sampling. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 1333-1341.	3.1	11
71	Changes of hand preference in Parkinson's disease. <i>Journal of Neural Transmission</i> , 2012, 119, 693-696.	2.8	6
72	Use of the theory of planned behaviour to assess factors influencing the identification of individuals at ultra-high risk for psychosis in primary care. <i>Microbial Biotechnology</i> , 2012, 6, 265-275.	1.7	7

#	ARTICLE	IF	CITATIONS
73	Cortisol levels in response to starting school in children at increased risk for social phobia. <i>Psychoneuroendocrinology</i> , 2012, 37, 462-474.	2.7	33
74	The validity of subjective quality of life measures in psychotic patients with severe psychopathology and cognitive deficits: an item response model analysis. <i>Quality of Life Research</i> , 2012, 21, 237-246.	3.1	23
75	Profiles of family-focused adverse experiences through childhood and early adolescence: The ROOTS project a community investigation of adolescent mental health. <i>BMC Psychiatry</i> , 2011, 11, 109.	2.6	92
76	General and specific components of depression and anxiety in an adolescent population. <i>BMC Psychiatry</i> , 2011, 11, 191.	2.6	47
77	Measuring patients' views: a bifactor model of distinct patient-reported outcomes in psychosis. <i>Psychological Medicine</i> , 2011, 41, 277-289.	4.5	29
78	Timing of menarche and depressive symptoms in adolescent girls from a UK cohort. <i>British Journal of Psychiatry</i> , 2011, 198, 17-23.	2.8	91
79	Polymorphisms in <i>BDNF</i> (Val66Met) and <i>5-HTTLPR</i> , morning cortisol and subsequent depression in at-risk adolescents. <i>British Journal of Psychiatry</i> , 2010, 197, 365-371.	2.8	60
80	An Evaluation of the Precision of Measurement of Ryff's Psychological Well-Being Scales in a Population Sample. <i>Social Indicators Research</i> , 2010, 97, 357-373.	2.7	93
81	The effects of maternal postnatal depression and child sex on academic performance at age 16 years: a developmental approach. <i>Journal of Child Psychology and Psychiatry and Allied Disciplines</i> , 2010, 51, 1150-1159.	5.2	161
82	Cohort Profile: Risk patterns and processes for psychopathology emerging during adolescence: the ROOTS project. <i>International Journal of Epidemiology</i> , 2010, 39, 361-369.	1.9	62
83	Latent mixture models for multivariate and longitudinal outcomes. <i>Statistical Methods in Medical Research</i> , 2010, 19, 271-289.	1.5	52
84	Measuring the benefits of treatment for psychosis: validity and responsiveness of the EQ-5D. <i>British Journal of Psychiatry</i> , 2009, 195, 170-177.	2.8	41
85	Cognitive behaviour therapy for improving social recovery in psychosis: a report from the ISREP MRC Trial Platform study (Improving Social Recovery in Early Psychosis). <i>Psychological Medicine</i> , 2009, 39, 1627.	4.5	101
86	Serotonin transporter genotype, morning cortisol and subsequent depression in adolescents. <i>British Journal of Psychiatry</i> , 2009, 195, 39-45.	2.8	87
87	Is the incidence of psychotic disorder in decline? Epidemiological evidence from two decades of research. <i>International Journal of Epidemiology</i> , 2009, 38, 1255-1264.	1.9	33
88	Randomized Trial of a Parenting Intervention for Very Preterm Infants: Outcome at 2 Years. <i>Journal of Pediatrics</i> , 2009, 155, 488-494.e1.	1.8	44
89	Health-Related Quality of Life (HRQL) and Continuous Antipsychotic Treatment: 3-year Results from the Schizophrenia Health Outcomes (SOHO) Study. <i>Value in Health</i> , 2009, 12, 536-543.	0.3	51
90	A longitudinal investigation of the impact of typology of urinary incontinence on quality of life during midlife: Results from a British prospective study. <i>Maturitas</i> , 2009, 64, 246-248.	2.4	14

#	ARTICLE	IF	CITATIONS
91	Cognitive behaviour therapy for improving social recovery in psychosis: Cost-effectiveness analysis. <i>Schizophrenia Research</i> , 2009, 112, 158-163.	2.0	28
92	Outcomes of conduct problems in adolescence: 40 year follow-up of national cohort. <i>BMJ: British Medical Journal</i> , 2009, 338, a2981-a2981.	2.3	277
93	The relationship between early personality and midlife psychological well-being: evidence from a UK birth cohort study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2008, 43, 679-687.	3.1	62
94	Classifying Patients by Antipsychotic Adherence Patterns Using Latent Class Analysis: Characteristics of Nonadherent Groups in the California Medicaid (Medi-Cal) Program. <i>Value in Health</i> , 2008, 11, 48-56.	0.3	61
95	Developmental Trajectories of Sex-typed Behavior in Boys and Girls: A Longitudinal General Population Study of Children Aged 2.5-8 Years. <i>Child Development</i> , 2008, 79, 1583-1593.	3.0	107
96	Factors associated with antidepressant, anxiolytic and hypnotic use over 17 years in a national cohort. <i>Journal of Affective Disorders</i> , 2008, 110, 234-240.	4.1	27
97	Trajectories of Daytime Wetting and Soiling in a United Kingdom 4 to 9-Year-Old Population Birth Cohort Study. <i>Journal of Urology</i> , 2008, 179, 1970-1975.	0.4	49
98	Psychiatric outcomes 10 years after treatment with antidepressants or anxiolytics. <i>British Journal of Psychiatry</i> , 2008, 193, 327-331.	2.8	10
99	Randomised trial of a parenting intervention during neonatal intensive care. <i>Archives of Disease in Childhood: Fetal and Neonatal Edition</i> , 2007, 92, F438-F443.	2.8	97
100	Forty-Year Psychiatric Outcomes Following Assessment for Internalizing Disorder in Adolescence. <i>American Journal of Psychiatry</i> , 2007, 164, 126-133.	7.2	122
101	A Longitudinal Typology of Symptoms of Depression and Anxiety Over the Life Course. <i>Biological Psychiatry</i> , 2007, 62, 1265-1271.	1.3	147
102	Improvement and decline of cognitive function in schizophrenia over one year: a longitudinal investigation using latent growth modelling. <i>BMC Psychiatry</i> , 2007, 7, 16.	2.6	16
103	Analysis of cluster randomized cross-over trial data: a comparison of methods. <i>Statistics in Medicine</i> , 2007, 26, 274-289.	1.6	89
104	Biased Mentalizing in Children Aged Seven to 11: Latent Class Confirmation of Response Styles to Social Scenarios and Associations with Psychopathology. <i>Social Development</i> , 2007, 16, 181-202.	1.3	52
105	Improvements in social functioning reported by a birth cohort in mid-adult life: A person-centred analysis of GHQ-28 social dysfunction items using latent class analysis. <i>Personality and Individual Differences</i> , 2007, 42, 305-316.	2.9	18
106	A Life Course Approach to Well-Being. , 2007, , 187-205.		6
107	Psychometric evaluation and predictive validity of Ryff's psychological well-being items in a UK birth cohort sample of women. <i>Health and Quality of Life Outcomes</i> , 2006, 4, 76.	2.4	169
108	Three decades of antidepressant, anxiolytic and hypnotic use in a national population birth cohort. <i>British Journal of Psychiatry</i> , 2006, 189, 156-160.	2.8	19

#	ARTICLE	IF	CITATIONS
109	The Short Mood and Feelings Questionnaire (SMFQ): A Unidimensional Item Response Theory and Categorical Data Factor Analysis of Self-Report Ratings from a Community Sample of 7-through 11-Year-Old Children. <i>Journal of Abnormal Child Psychology</i> , 2006, 34, 365-377.	3.5	271
110	Clinical and social determinants of duration of untreated psychosis in the Å†SOP first-episode psychosis study. <i>British Journal of Psychiatry</i> , 2006, 189, 446-452.	2.8	168
111	BILOG-MG, MULTILOG, PARSCALE and TESTFACT4 programs on CD with single volume manual combining reference guide and users guide (program examples: Input and output). <i>British Journal of Mathematical and Statistical Psychology</i> , 2005, 58, 193-195.	1.4	3
112	Association Between Duration of Untreated Psychosis and Outcome in Cohorts of First-Episode Patients. <i>Archives of General Psychiatry</i> , 2005, 62, 975.	12.3	1,303
113	Deciding who gets treatment for depression and anxiety: a study of consecutive GP attenders. <i>British Journal of General Practice</i> , 2005, 55, 846-53.	1.4	23
114	The persistence of developmental markers in childhood and adolescence and risk for schizophrenic psychoses in adult life. A 34-year follow-up of the Northern Finland 1966 birth cohort. <i>Schizophrenia Research</i> , 2004, 71, 213-225.	2.0	55
115	A modelling strategy for the analysis of clinical trials with partly missing longitudinal data. <i>International Journal of Methods in Psychiatric Research</i> , 2003, 12, 139-150.	2.1	23
116	Developmental Typology of Trajectories to Nighttime Bladder Control: Epidemiologic Application of Longitudinal Latent Class Analysis. <i>American Journal of Epidemiology</i> , 2003, 157, 834-842.	3.4	76
117	Impact of the ICDâ€™10 Primary Health Care (PHC) diagnostic and management guidelines for mental disorders on detection and outcome in primary care. <i>British Journal of Psychiatry</i> , 2003, 182, 20-30.	2.8	71
118	Early developmental milestones in adult schizophrenia and other psychoses. A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Schizophrenia Research</i> , 2001, 52, 1-19.	2.0	159
119	Educational consequences of mental disorders treated in hospital. A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Psychological Medicine</i> , 2001, 31, 339-349.	4.5	50
120	Aggressive incidents in first-episode psychosis. <i>British Journal of Psychiatry</i> , 2001, 178, 433-440.	2.8	88
121	Birth order and risk for schizophrenia: a 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Acta Psychiatrica Scandinavica</i> , 2001, 104, 148-152.	4.5	35
122	Juvenile and early adulthood smoking and adult educational achievements â€™ A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Scandinavian Journal of Public Health</i> , 2001, 29, 87-95.	2.3	14
123	Three-year outcome of first-episode psychoses in an established community psychiatric service. <i>British Journal of Psychiatry</i> , 2000, 176, 210-216.	2.8	73
124	Non-linear relationship between an index of social deprivation, psychiatric admission prevalence and the incidence of psychosis. <i>Psychological Medicine</i> , 2000, 30, 177-185.	4.5	91
125	Childhood and adolescent predictors of schizophrenia in the Northern Finland 1966 Birth Cohort â€™ a descriptive life-span model. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2000, 250, 311-319.	3.2	63
126	Outcome of psychosis in people of Africanâ€™Caribbean family origin. <i>British Journal of Psychiatry</i> , 1999, 175, 43-49.	2.8	56

#	ARTICLE	IF	CITATIONS
127	Discharged from special hospital under restrictions: a comparison of the fates of psychopaths and the mentally ill. <i>Criminal Behaviour and Mental Health</i> , 1998, 8, 39-55.	0.8	53
128	The predictive validity of a diagnosis of schizophrenia. <i>British Journal of Psychiatry</i> , 1997, 170, 321-327.	2.8	71
129	The Course of Schizophrenia Over 13 Years. <i>British Journal of Psychiatry</i> , 1996, 169, 580-586.	2.8	110
130	Characteristics of Outcome in Schizophrenia at 13 Years. <i>British Journal of Psychiatry</i> , 1995, 167, 596-603.	2.8	137