

Alfredo Valido

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1641931/publications.pdf>

Version: 2024-02-01

41

papers

2,931

citations

218677

26

h-index

289244

40

g-index

41

all docs

41

docs citations

41

times ranked

4115

citing authors

#	ARTICLE	IF	CITATIONS
1	Body size in ecological networks. <i>Trends in Ecology and Evolution</i> , 2005, 20, 402-409.	8.7	931
2	Lizards as pollinators and seed dispersers: an island phenomenon. <i>Trends in Ecology and Evolution</i> , 2003, 18, 177-181.	8.7	278
3	Honeybees disrupt the structure and functionality of plant-pollinator networks. <i>Scientific Reports</i> , 2019, 9, 4711.	3.3	140
4	Impact of introduced honey bees on native pollination interactions of the endemic <i>Echium wildpretii</i> (Boraginaceae) on Tenerife, Canary Islands. <i>Biological Conservation</i> , 2004, 118, 301-311.	4.1	121
5	The signatures of Anthropocene defaunation: cascading effects of the seed dispersal collapse. <i>Scientific Reports</i> , 2016, 6, 24820.	3.3	110
6	Frugivory and Seed Dispersal by the Lizard <i>Gallotia galloti</i> (Lacertidae) in a Xeric Habitat of the Canary Islands. <i>Oikos</i> , 1994, 70, 403.	2.7	104
7	Late Neogene history of the laurel tree (<i>Laurus</i> L., Lauraceae) based on phylogeographical analyses of Mediterranean and Macaronesian populations. <i>Journal of Biogeography</i> , 2009, 36, 1270-1281.	3.0	104
8	Colour, design and reward: phenotypic integration of fleshy fruit displays. <i>Journal of Evolutionary Biology</i> , 2011, 24, 751-760.	1.7	93
9	Long-term demographic consequences of a seed dispersal disruption. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2012, 279, 3298-3303.	2.6	84
10	Geographic patterns in fruit colour diversity: do leaves constrain the colour of fleshy fruits?. <i>Oecologia</i> , 2009, 159, 337-343.	2.0	65
11	Birds see the true colours of fruits to live off the fat of the land. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2014, 281, 20132516.	2.6	65
12	Downsized mutualisms: Consequences of seed dispersers' body-size reduction for early plant recruitment. <i>Perspectives in Plant Ecology, Evolution and Systematics</i> , 2015, 17, 151-159.	2.7	59
13	Native birds and insects, and introduced honey bees visiting <i>Echium wildpretii</i> (Boraginaceae) in the Canary Islands. <i>Acta Oecologica</i> , 2002, 23, 413-419.	1.1	53
14	Bird-flower interactions in the Macaronesian islands. <i>Journal of Biogeography</i> , 2004, 31, 1945-1953.	3.0	51
15	Disturbance regimes, gap-demanding trees and seed mass related to tree height in warm temperate rain forests worldwide. <i>Biological Reviews</i> , 2013, 88, 701-744.	10.4	48
16	Opportunistic nectar-feeding birds are effective pollinators of bird-flowers from Canary Islands: experimental evidence from <i>Isoplexis canariensis</i> (Scrophulariaceae). <i>American Journal of Botany</i> , 2008, 95, 1408-1415.	1.7	47
17	Adaptation of flower and fruit colours to multiple, distinct mutualists. <i>New Phytologist</i> , 2014, 201, 678-686.	7.3	47
18	Fleshy Fruits in the Diet of Canarian Lizards <i>Gallotia galloti</i> (Lacertidae) in a Xeric Habitat of the Island of Tenerife. <i>Journal of Herpetology</i> , 2003, 37, 741-747.	0.5	40

#	ARTICLE	IF	CITATIONS
19	Functional consequences of plant-animal interactions along the mutualism-antagonism gradient. <i>Ecology</i> , 2017, 98, 1266-1276.	3.2	37
20	Digestive ecology of two omnivorous Canarian lizard species (Gallotia, Lacertidae). <i>Amphibia - Reptilia</i> , 2003, 24, 331-344.	0.5	36
21	Quantity and quality components of effectiveness in insular pollinator assemblages. <i>Oecologia</i> , 2013, 173, 179-190.	2.0	36
22	Global geographic patterns in the colours and sizes of animal-dispersed fruits. <i>Global Ecology and Biogeography</i> , 2018, 27, 1339-1351.	5.8	36
23	Persisting in defaunated landscapes: Reduced plant population connectivity after seed dispersal collapse. <i>Journal of Ecology</i> , 2018, 106, 936-947.	4.0	34
24	Macaronesia as a Fruitful Arena for Ecology, Evolution, and Conservation Biology. <i>Frontiers in Ecology and Evolution</i> , 2021, 9, .	2.2	33
25	Frugivory and Seed Dispersal by Lizards: A Global Review. <i>Frontiers in Ecology and Evolution</i> , 2019, 7, .	2.2	32
26	Indirect seed dispersal by the feral cats <i>Felis catus</i> in island ecosystems (Canary Islands). <i>Ecography</i> , 1996, 19, 3-6.	4.5	31
27	Morphological evolution and changes in foraging behaviour of island and mainland populations of Blue Tit (<i>Parus caeruleus</i>) ? a test of convergence and ecomorphological hypotheses. <i>Evolutionary Ecology</i> , 1994, 8, 25-35.	1.2	27
28	Consequences of plant-pollinator and floral-herbivore interactions on the reproductive success of the Canary Islands endemic <i>Canarina canariensis</i> (Campanulaceae). <i>American Journal of Botany</i> , 2011, 98, 1465-1474.	1.7	26
29	Pollinator shifts drive petal epidermal evolution on the Macaronesian Islands bird-flowered species. <i>Biology Letters</i> , 2016, 12, 20160022.	2.3	23
30	The role of the brown bear <i>Ursus arctos</i> as a legitimate megafaunal seed disperser. <i>Scientific Reports</i> , 2021, 11, 1282.	3.3	20
31	Habitat Distribution of Canary Chaffinches Among Islands: Competitive Exclusion or Species-Specific Habitat Preferences?. <i>Journal of Biogeography</i> , 1992, 19, 383.	3.0	17
32	Conflicting selection on <i>Cneorum tricoccon</i> (Rutaceae) seed size caused by native and alien seed dispersers. <i>Evolution; International Journal of Organic Evolution</i> , 2019, 73, 2204-2215.	2.3	17
33	Frugivory and factors influencing visitation by birds at "Baloâ™ (<i>Plocama pendula</i> Ait., Rubiaceae) plants in the Canary Islands. <i>Ecoscience</i> , 1999, 6, 531-538.	1.4	16
34	Bird-pollinated Macaronesian Lotus (Leguminosae) evolved within a group of entomophilous ancestors with post-anthesis flower color change. <i>Perspectives in Plant Ecology, Evolution and Systematics</i> , 2013, 15, 193-204.	2.7	16
35	Isolation and characterization of 20 microsatellite loci for laurel species (<i>Laurus</i>, Lauraceae). <i>American Journal of Botany</i> , 2010, 97, e26-30.	1.7	13
36	Hotspots of damage by antagonists shape the spatial structure of plant-pollinator interactions. <i>Ecology</i> , 2015, 96, 2181-2191.	3.2	11

#	ARTICLE	IF	CITATIONS
37	Impact of the introduced honeybees (<i>Apis mellifera</i> , Apidae) on Teide National Park (Tenerife, Canary) Tj ETQq1 1 0.784314 rgBT /Overde	0.4	11
38	Heterostyly in the Canarian endemic <i>Jasminum odoratissimum</i> (Oleaceae). Nordic Journal of Botany, 2003, 23, 537-539.	0.5	10
39	Isolation and characterization of 13 microsatellite loci for <i>< i>Neochamaelea pulverulenta</i></i> (Cneoraceae). Molecular Ecology Resources, 2009, 9, 1497-1500.	4.8	5
40	Sobre la presencia de <i>< i>Cneorum</i></i> (Cneoraceae) en Cuba: ¿ejemplo de disyunción biogeográfica Mediterráneo-Caribe?. Anales Del Jardin Botanico De Madrid, 2009, 66, 25-33.	0.4	4
41	X Reunión anual de Ecoflor. Ecosistemas, 2013, 22, 125-125.	0.4	0