

Murat Serilmez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1678055/publications.pdf>

Version: 2024-02-01

36
papers

513
citations

623734

14
h-index

713466

21
g-index

36
all docs

36
docs citations

36
times ranked

881
citing authors

#	ARTICLE	IF	CITATIONS
1	Can serum matrix metalloproteinase-9 and SMAD2 levels predict lamina propria invasion in bladder urothelial carcinoma?. <i>International Journal of Clinical Practice</i> , 2021, 75, e14277.	1.7	0
2	Circulating annexin A2 as a biomarker in patients with pancreatic cancer. <i>Journal of Cancer Research and Therapeutics</i> , 2020, 16, 110.	0.9	7
3	Serum nectin-2 and nectin-4 are diagnostic in lung cancer: which is superior?. <i>Wiener Klinische Wochenschrift</i> , 2019, 131, 419-426.	1.9	15
4	The Diagnostic Significance of PDGF, EphA7, CCR5, and CCL5 Levels in Colorectal Cancer. <i>Biomolecules</i> , 2019, 9, 464.	4.0	25
5	Detection of serum protein and circulating mRNA of cMET, HGF EGF and EGFR levels in lung cancer patients to guide individualized therapy. <i>Cancer Biomarkers</i> , 2019, 25, 177-184.	1.7	7
6	Diagnostic and Prognostic Significance of Carboxypeptidase A4 (CPA4) in Breast Cancer. <i>Biomolecules</i> , 2019, 9, 103.	4.0	6
7	Clinical Significance of Serum NEDD9 Levels in Patients with Pancreatic Cancer. <i>Biomolecules</i> , 2018, 8, 169.	4.0	5
8	Serum Glypican-3, Vascular Endothelial Growth Factor, and Interleukin-6 Levels in Hepatocellular Carcinoma. <i>Turk Onkoloji Dergisi</i> , 2018, , .	0.0	0
9	Circulating interleukin-18 (IL-18) is a predictor of response to gemcitabine based chemotherapy in patients with pancreatic adenocarcinoma. <i>Journal of Infection and Chemotherapy</i> , 2017, 23, 196-200.	1.7	12
10	Significance of serum neural precursor cell-expressed developmentally downregulated protein 9 in melanoma. <i>Molecular and Clinical Oncology</i> , 2017, 8, 204-208.	1.0	3
11	Clinical Significance of Serum Ykl-40 (Chitinase-3-Like-1 Protein) as a Biomarker in Melanoma: an Analysis of 112 Turkish Patients. <i>Asian Pacific Journal of Cancer Prevention</i> , 2017, 18, 1383-1387.	1.2	10
12	Role of several cytokines and adhesion molecules in the diagnosis and prediction of survival of hepatocellular carcinoma. <i>Arab Journal of Gastroenterology</i> , 2016, 17, 164-167.	0.9	6
13	Clinical significance of serum omentin-1 levels in patients with pancreatic adenocarcinoma. <i>BBA Clinical</i> , 2016, 6, 138-142.	4.1	20
14	Circulating serum levels of angiopoietin-1 and angiopoietin-2 in nasopharynx and larynx carcinoma patients. <i>Tumor Biology</i> , 2016, 37, 8979-8983.	1.8	4
15	Serum activated leukocyte cell adhesion molecule and intercellular adhesion molecule-1 in patients with gastric cancer: Can they be used as biomarkers?. <i>Biomedicine and Pharmacotherapy</i> , 2016, 77, 86-91.	5.6	14
16	Elevated circulating monocyte chemoattractant protein 1 (MCP-1/CCL-2) level may be an unfavorable predictive factor to platinum- and taxane-based combination chemotherapy in patients with gastric cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 77, 127-131.	2.3	8
17	Clinical significance of serum interleukin-29, interleukin-32, and tumor necrosis factor alpha levels in patients with gastric cancer. <i>Tumor Biology</i> , 2016, 37, 405-412.	1.8	15
18	Serum nectin-2 levels are diagnostic and prognostic in patients with colorectal carcinoma. <i>Clinical and Translational Oncology</i> , 2016, 18, 160-171.	2.4	26

#	ARTICLE	IF	CITATIONS
19	Is serum tenascin-C level potential biomarker in pancreatic adenocarcinoma?. Medical Journal of Bakirkoy, 2016, , 80-86.	0.1	0
20	D-dimer and international normalized ratio (INR) are correlated with tumor markers and disease stage in colorectal cancer patients. Cancer Biomarkers, 2015, 15, 405-411.	1.7	23
21	Diagnostic value of serum M30 and M65 in patients with nasopharyngeal carcinoma. Tumor Biology, 2015, 36, 1039-1044.	1.8	6
22	Serum levels of macrophage migration-inhibitory factor (MIF) have diagnostic, predictive and prognostic roles in epithelial ovarian cancer patients. Tumor Biology, 2014, 35, 3327-3331.	1.8	15
23	Clinical significance of serum circulating insulin-like growth factor-1 (IGF-1) mRNA in hepatocellular carcinoma. Tumor Biology, 2014, 35, 2729-2739.	1.8	6
24	Clinical significance of serum insulin-like growth factor-1 (IGF-1) and insulinlike growth factor binding protein-3 (IGFBP-3) in patients with epithelial ovarian cancer. Tumor Biology, 2014, 35, 3125-3132.	1.8	18
25	Clinical significance of serum transforming growth factor-beta 1 (TGF- β 1) levels in patients with epithelial ovarian cancer. Tumor Biology, 2014, 35, 3611-3616.	1.8	12
26	Clinical significance of serum epithelial cell adhesion molecule (EPCAM) and vascular cell adhesion molecule-1 (VCAM-1) levels in patients with epithelial ovarian cancer. Tumor Biology, 2014, 35, 3095-3102.	1.8	20
27	Increased serum level of epidermal growth factor receptor (EGFR) is associated with poor progression-free survival in patients with epithelial ovarian cancer. Cancer Chemotherapy and Pharmacology, 2014, 73, 631-637.	2.3	7
28	Clinical significance of serum hepatocyte growth factor (HGF) levels in hepatocellular carcinoma. Tumor Biology, 2014, 35, 2327-2333.	1.8	15
29	Clinical significance of serum tenascin-C levels in breast cancer. Tumor Biology, 2014, 35, 6619-6625.	1.8	17
30	The diagnostic, predictive, and prognostic role of serum epithelial cell adhesion molecule (EpCAM) and vascular cell adhesion molecule-1 (VCAM-1) levels in breast cancer. Tumor Biology, 2014, 35, 8849-8860.	1.8	16
31	Clinical and Prognostic Significance of Coagulation Assays in Gastric Cancer. Journal of Gastrointestinal Cancer, 2013, 44, 285-292.	1.3	54
32	A high serum level of M65 is associated with tumour aggressiveness and an unfavourable prognosis for epithelial ovarian cancer. Cancer Chemotherapy and Pharmacology, 2013, 72, 437-444.	2.3	6
33	Clinical and prognostic significance of coagulation assays in lung cancer. Respiratory Medicine, 2013, 107, 451-457.	2.9	71
34	Clinical significance of serum M30 and M65 levels in melanoma. Melanoma Research, 2013, 23, 390-395.	1.2	4
35	For Which Cancer Types can Neuron-Specific Enolase be Clinically Helpful in Turkish Patients?. Asian Pacific Journal of Cancer Prevention, 2013, 14, 2541-2544.	1.2	7
36	Immobilization and stabilization of β -galactosidase on Sepabeads EC-EA and EC-HA. International Journal of Biological Macromolecules, 2011, 49, 855-860.	7.5	33