

Yasuo Miyoshi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1693959/publications.pdf>

Version: 2024-02-01

140
papers

5,192
citations

94433

37
h-index

98798

67
g-index

143
all docs

143
docs citations

143
times ranked

7508
citing authors

#	ARTICLE	IF	CITATIONS
1	Abstract P4-07-26: Clinical impact of tumor infiltrating lymphocytes and neutrophil-lymphocyte ratio in estrogen receptor-positive/HER2-negative breast cancer patients with high 21-gene signature recurrence scores. <i>Cancer Research</i> , 2022, 82, P4-07-26-P4-07-26.	0.9	0
2	Harmonized pretreatment quantitative volume-based FDG-PET/CT parameters for prognosis of stage I-III breast cancer: Multicenter study. <i>Oncotarget</i> , 2021, 12, 95-105.	1.8	14
3	HERC2 inactivation abrogates nucleolar localization of RecQ helicases BLM and WRN. <i>Scientific Reports</i> , 2021, 11, 360.	3.3	8
4	Preliminary study of AI-assisted diagnosis using FDG-PET/CT for axillary lymph node metastasis in patients with breast cancer. <i>EJNMMI Research</i> , 2021, 11, 10.	2.5	20
5	Indices of peripheral leukocytes predict longer overall survival in breast cancer patients on eribulin in Japan. <i>Breast Cancer</i> , 2021, 28, 945-955.	2.9	6
6	Japanese subpopulation analysis of MONARCH 2: phase 3 study of abemaciclib plus fulvestrant for treatment of hormone receptor-positive, human epidermal growth factor receptor 2-negative breast cancer that progressed on endocrine therapy. <i>Breast Cancer</i> , 2021, 28, 1038-1050.	2.9	10
7	Investigation of a Novel S-1 Administration Schedule for Treating Metastatic and Recurrent Breast Cancer (KBCOG13). <i>Anticancer Research</i> , 2021, 41, 3121-3126.	1.1	0
8	Breast cancer diffuse liver metastasis with high liver stiffness using ultrasound elastography. <i>Acta Hepatologica Japonica</i> , 2021, 62, 647-655.	0.1	0
9	Low intratumoral genetic neutrophil-to-lymphocyte ratio (NLR) is associated with favorable tumor immune microenvironment and with survival in triple negative breast cancer (TNBC). <i>American Journal of Cancer Research</i> , 2021, 11, 5743-5755.	1.4	1
10	Prognostic Significance of Neutrophil-to-lymphocyte Ratio in Luminal Breast Cancers With Low Levels of Tumour-infiltrating Lymphocytes. <i>Anticancer Research</i> , 2020, 40, 2871-2880.	1.1	16
11	High absolute lymphocyte counts are associated with longer overall survival in patients with metastatic breast cancer treated with eribulin but not with treatment of physician's choice in the EMBRACE study. <i>Breast Cancer</i> , 2020, 27, 706-715.	2.9	41
12	C-Reactive Protein and Absolute Lymphocyte Count Can Predict Overall Survival of Patients Treated With Eribulin. <i>Anticancer Research</i> , 2020, 40, 4147-4156.	1.1	10
13	Association Between FOXP3/CD8 Lymphocyte Ratios and Tumor Infiltrating Lymphocyte Levels in Different Breast Cancer Subtypes. <i>Anticancer Research</i> , 2020, 40, 2141-2150.	1.1	11
14	Significant association between high serum CCL5 levels and better disease-free survival of patients with early breast cancer. <i>Cancer Science</i> , 2020, 111, 209-218.	3.9	23
15	Baseline neutrophil-to-lymphocyte ratio and c-reactive protein predict efficacy of treatment with bevacizumab plus paclitaxel for locally advanced or metastatic breast cancer. <i>Oncotarget</i> , 2020, 11, 86-98.	1.8	15
16	The GALNT6-GALS3BP axis promotes breast cancer cell growth. <i>International Journal of Oncology</i> , 2020, 56, 581-595.	3.3	9
17	A Single-centre, Retrospective, Observational Analysis of Fulvestrant for Recurrent/metastatic Breast Cancer According to Metastatic Site. <i>Anticancer Research</i> , 2019, 39, 5653-5662.	1.1	8
18	High Serum Levels of Interleukin-18 Are Associated With Worse Outcomes in Patients With Breast Cancer. <i>Anticancer Research</i> , 2019, 39, 5009-5018.	1.1	17

#	ARTICLE	IF	CITATIONS
19	Participants in a randomized controlled trial had longer overall survival than non-participants: a prospective cohort study. <i>Breast Cancer Research and Treatment</i> , 2019, 176, 631-635.	2.5	4
20	Prognostic significance of tumor-infiltrating lymphocytes may differ depending on Ki67 expression levels in estrogen receptor-positive/HER2-negative operated breast cancers. <i>Breast Cancer</i> , 2019, 26, 738-747.	2.9	21
21	Significance of Metabolic Tumor Volume at Baseline and Reduction of Mean Standardized Uptake Value in 18F-FDG-PET/CT Imaging for Predicting Pathological Complete Response in Breast Cancers Treated with Preoperative Chemotherapy. <i>Annals of Surgical Oncology</i> , 2019, 26, 2175-2183.	1.5	16
22	Significance of baseline neutrophil-to-lymphocyte ratio for progression-free survival of patients with HER2-positive breast cancer treated with trastuzumab emtansine. <i>Scientific Reports</i> , 2019, 9, 1811.	3.3	22
23	Abstract 5313: Frequent downregulation of SALL3 by genetic and epigenetic alterations is involved in progression and chemoresistance of triple negative breast cancers. , 2019, , .		0
24	Abstract 3813: PHB2 inactivation by AKAP-BIG3 is required for progression of HER2-overexpressing breast cancer. , 2019, , .		0
25	Prognostic value of F-FDG PET/CT prior to breast cancer treatment. Comparison with magnetic resonance spectroscopy and diffusion weighted imaging. <i>Hellenic Journal of Nuclear Medicine</i> , 2019, 22, 25-35.	0.3	3
26	Randomized phase II study of anastrozole plus tegafur-uracil as neoadjuvant therapy for ER-positive breast cancer in postmenopausal Japanese women (Neo-ACET BC). <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 81, 755-762.	2.3	8
27	Significant Association Between Low Baseline Neutrophil-to-Lymphocyte Ratio and Improved Progression-free Survival of Patients With Locally Advanced or Metastatic Breast Cancer Treated With Eribulin But Not With Nab-Paclitaxel. <i>Clinical Breast Cancer</i> , 2018, 18, 400-409.	2.4	44
28	Abundant tumor infiltrating lymphocytes after primary systemic chemotherapy predicts poor prognosis in estrogen receptor-positive/HER2-negative breast cancers. <i>Breast Cancer Research and Treatment</i> , 2018, 168, 135-145.	2.5	27
29	Mechanism of resistance to endocrine therapy in breast cancer: the important role of PI3K/Akt/mTOR in estrogen receptor-positive, HER2-negative breast cancer. <i>Breast Cancer</i> , 2018, 25, 392-401.	2.9	134
30	Prognostic value of FDG-PET and DWI in breast cancer. <i>Annals of Nuclear Medicine</i> , 2018, 32, 44-53.	2.2	34
31	Improved prognosis of low baseline neutrophil-to-lymphocyte ratio is significantly exclusive in breast cancer patients with high absolute counts of lymphocytes. <i>Molecular and Clinical Oncology</i> , 2018, 10, 275-284.	1.0	12
32	Actionable Gene Alterations in an Asian Population With Triple-Negative Breast Cancer. <i>JCO Precision Oncology</i> , 2018, 2, 1-13.	3.0	3
33	Frequent downregulation of LRRC26 by epigenetic alterations is involved in the malignant progression of triple-negative breast cancer. <i>International Journal of Oncology</i> , 2018, 52, 1539-1558.	3.3	8
34	HERC2 Facilitates BLM and WRN Helicase Complex Interaction with RPA to Suppress G-Quadruplex DNA. <i>Cancer Research</i> , 2018, 78, 6371-6385.	0.9	41
35	Predictive impact of absolute lymphocyte counts for progression-free survival in human epidermal growth factor receptor 2-positive advanced breast cancer treated with pertuzumab and trastuzumab plus eribulin or nab-paclitaxel. <i>BMC Cancer</i> , 2018, 18, 982.	2.6	33
36	Response to neoadjuvant chemotherapy for breast cancer judged by PERCIST – multicenter study in Japan. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2018, 45, 1661-1671.	6.4	23

#	ARTICLE	IF	CITATIONS
37	High levels of serum CA15â€³ and residual invasive tumor size are associated with poor prognosis for breast cancer patients with nonâ€³pathological complete response after neoadjuvant chemotherapy. <i>Journal of Surgical Oncology</i> , 2018, 118, 228-237.	1.7	4
38	Independent prognostic impact of preoperative serum carcinoembryonic antigen and cancer antigen 15-3 levels for early breast cancer subtypes. <i>World Journal of Surgical Oncology</i> , 2018, 16, 26.	1.9	17
39	Fbxo22-mediated KDM4B degradation determines selective estrogen receptor modulator activity in breast cancer. <i>Journal of Clinical Investigation</i> , 2018, 128, 5603-5619.	8.2	39
40	Tumor mutation burden in triple negative breast cancer patients in Japan.. <i>Journal of Clinical Oncology</i> , 2018, 36, e13111-e13111.	1.6	12
41	Assessment of tumor response to neoadjuvant chemotherapy in patients with breast cancer using MRI and FDG-PET/CT-RECIST 1.1 vs. PERCIST 1.0. <i>Nagoya Journal of Medical Science</i> , 2018, 80, 183-197.	0.3	15
42	C-reactive Protein Level on Postoperative Day One is Associated with Chronic Postsurgical Pain After Mastectomy. <i>Anesthesiology and Pain Medicine</i> , 2018, 8, e79331.	1.3	8
43	Abundant options to avoid toxicity and alternative strategies for human epidermal growth factor receptor 2-positive and hormone receptor-positive advanced breast cancer. <i>Translational Cancer Research</i> , 2018, 7, S514-S518.	1.0	0
44	A claims data analysis of single institution for comorbidity among various types of cancer.. <i>Journal of Clinical Oncology</i> , 2018, 36, e18843-e18843.	1.6	0
45	Abstract 1837: Overcoming trastuzumab resistance in HER2-overexpressing breast cancer by utilizing PHB2, a tumor suppressor of multiple resistance pathways. , 2018, , .		0
46	Abstract 5315: Frequent downregulation of <i>SALL3</i> by recurrent genetic and epigenetic alterations is involved in triple-negative breast cancers. <i>Cancer Research</i> , 2018, 78, 5315-5315.	0.9	1
47	Different patterns of change in bone turnover markers during treatment with bone-modifying agents for breast cancer patients with bone metastases. <i>Breast Cancer</i> , 2017, 24, 245-253.	2.9	8
48	A-kinase anchoring protein BIG3 coordinates oestrogen signalling in breast cancer cells. <i>Nature Communications</i> , 2017, 8, 15427.	12.8	14
49	Stapled BIG3 helical peptide ERAP potentiates anti-tumour activity for breast cancer therapeutics. <i>Scientific Reports</i> , 2017, 7, 1821.	3.3	11
50	Body mass index and menopausal disorders during menopause affect vasomotor symptoms of postmenopausal Japanese breast cancer patients treated with anastrozole: a prospective multicenter cohort study of patient-reported outcomes. <i>Breast Cancer</i> , 2017, 24, 528-534.	2.9	3
51	Tumor size and proliferative marker geminin rather than Ki67 expression levels significantly associated with maximum uptake of 18F-deoxyglucose levels on positron emission tomography for breast cancers. <i>PLoS ONE</i> , 2017, 12, e0184508.	2.5	6
52	Integrated analysis of somatic mutations and immune microenvironment of multiple regions in breast cancers. <i>Oncotarget</i> , 2017, 8, 62029-62038.	1.8	28
53	Prognostic significance of preoperative 18F-FDG PET/CT for breast cancer subtypes. <i>Breast</i> , 2016, 30, 5-12.	2.2	31
54	HP1 regulates the localization of FANCI at sites of DNA doubleâ€³strand breaks. <i>Cancer Science</i> , 2016, 107, 1406-1415.	3.9	14

#	ARTICLE	IF	CITATIONS
55	Impact of biomarker changes during neoadjuvant chemotherapy for clinical response in patients with residual breast cancers. <i>International Journal of Clinical Oncology</i> , 2016, 21, 254-261.	2.2	19
56	Clonal expansion of antitumor T cells in breast cancer correlates with response to neoadjuvant chemotherapy. <i>International Journal of Oncology</i> , 2016, 49, 471-478.	3.3	32
57	Prognostic significance of geminin expression levels in Ki67-high subset of estrogen receptor-positive and HER2-negative breast cancers. <i>Breast Cancer</i> , 2016, 23, 224-230.	2.9	9
58	Present and future role of FDG-PET/CT imaging in the management of breast cancer. <i>Japanese Journal of Radiology</i> , 2016, 34, 167-180.	2.4	25
59	Correlation of the SUVmax of FDG-PET and ADC values of diffusion-weighted MR imaging with pathologic prognostic factors in breast carcinoma. <i>European Journal of Radiology</i> , 2016, 85, 943-949.	2.6	61
60	Diagnostic and prognostic value of 18F-FDG PET/CT for axillary lymph node staging in patients with breast cancer. <i>Japanese Journal of Radiology</i> , 2016, 34, 220-228.	2.4	14
61	Risk factors for joint symptoms in postmenopausal Japanese breast cancer patients treated with anastrozole: a prospective multicenter cohort study of patient-reported outcomes. <i>International Journal of Clinical Oncology</i> , 2016, 21, 262-269.	2.2	19
62	Overall survival of participants compared to non-participants in a randomized-controlled trial (SELECT BC): A prospective cohort study.. <i>Journal of Clinical Oncology</i> , 2016, 34, 2527-2527.	1.6	0
63	High levels at baseline of serum pyridinoline crosslinked carboxyterminal telopeptide of type I collagen are associated with worse prognosis for breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2015, 154, 521-531.	2.5	6
64	Therapeutic advances in $BIG \rightarrow 3 \leftarrow PHB \rightarrow 2$ inhibition targeting the crosstalk between estrogen and growth factors in breast cancer. <i>Cancer Science</i> , 2015, 106, 550-558.	3.9	11
65	BIG3 Inhibits the Estrogen-Dependent Nuclear Translocation of PHB2 via Multiple Karyopherin-Alpha Proteins in Breast Cancer Cells. <i>PLoS ONE</i> , 2015, 10, e0127707.	2.5	19
66	Class I histone deacetylase inhibitors inhibit the retention of $BRCA \rightarrow 1$ and $53 \leftarrow BP \rightarrow 1$ at the site of DNA damage. <i>Cancer Science</i> , 2015, 106, 1050-1056.	3.9	28
67	Activation of mTOR/S6K But Not MAPK Pathways Might Be Associated With High Ki-67, ER+, and HER2 ⁺ Breast Cancer. <i>Clinical Breast Cancer</i> , 2015, 15, 197-203.	2.4	14
68	Interaction of BARD1 and HP1 Is Required for BRCA1 Retention at Sites of DNA Damage. <i>Cancer Research</i> , 2015, 75, 1311-1321.	0.9	83
69	Association between 18F-FDG uptake and molecular subtype of breast cancer. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2015, 42, 1371-1377.	6.4	62
70	Clinicopathological characteristics of breast cancer and trends in the management of breast cancer patients in Japan: Based on the Breast Cancer Registry of the Japanese Breast Cancer Society between 2004 and 2011. <i>Breast Cancer</i> , 2015, 22, 235-244.	2.9	58
71	High Ki-67 Expression and Low Progesterone Receptor Expression Could Independently Lead to a Worse Prognosis for Postmenopausal Patients with Estrogen Receptor-Positive and HER2-Negative Breast Cancer. <i>Clinical Breast Cancer</i> , 2015, 15, 204-211.	2.4	33
72	C4.4A highly expressed in HER2-positive human breast cancers may indicate a good prognosis. <i>Breast Cancer</i> , 2015, 22, 366-373.	2.9	7

#	ARTICLE	IF	CITATIONS
73	Association of body mass index with risk of luminal A but not luminal B estrogen receptor-positive and HER2-negative breast cancer for postmenopausal Japanese women. <i>Breast Cancer</i> , 2015, 22, 399-405.	2.9	10
74	Abstract P5-09-05: Prediction of bone metastases of breast cancer using combined markers of bone metabolism and inflammation. , 2015, , .		0
75	Abstract P1-12-04: Factors influencing on discontinuation of adjuvant anastrozole in postmenopausal Japanese breast cancer patients: Results from a prospective multicenter cohort study of patient-reported outcomes. , 2015, , .		0
76	Class I HDAC inhibitors inhibit the retention of BRCA1 and 53BP1 at the site of DNA damage.. <i>Journal of Clinical Oncology</i> , 2015, 33, e13552-e13552.	1.6	1
77	Biological characteristics of luminal subtypes in pre- and postmenopausal estrogen receptor-positive and HER2-negative breast cancers. <i>Breast Cancer</i> , 2014, 21, 52-57.	2.9	9
78	Influence of body mass index on clinicopathological factors including estrogen receptor, progesterone receptor, and Ki67 expression levels in breast cancers. <i>International Journal of Clinical Oncology</i> , 2014, 19, 467-472.	2.2	14
79	Strategy for treatment of isolated contralateral supraclavicular recurrence in patient with breast cancer after sentinel lymph node biopsy without axillary lymph node dissection. <i>International Cancer Conference Journal</i> , 2014, 3, 133-139.	0.5	0
80	Involvement of B3GALNT2 overexpression in the cell growth of breast cancer. <i>International Journal of Oncology</i> , 2014, 44, 427-434.	3.3	11
81	Xanthohumol suppresses oestrogen-signalling in breast cancer through the inhibition of BIG3-PHB2 interactions. <i>Scientific Reports</i> , 2014, 4, 7355.	3.3	68
82	Effect of body mass index and menopausal disorders during menopause on vasomotor symptoms of postmenopausal Japanese breast cancer patients treated with anastrozole: A prospective multicenter cohort study of patient-reported outcomes.. <i>Journal of Clinical Oncology</i> , 2014, 32, 9619-9619.	1.6	0
83	Phase II study of S-1 in combination with trastuzumab for HER2-positive metastatic breast cancer. <i>Anticancer Research</i> , 2014, 34, 3583-8.	1.1	2
84	Targeting BIG3-PHB2 interaction to overcome tamoxifen resistance in breast cancer cells. <i>Nature Communications</i> , 2013, 4, 2443.	12.8	63
85	Molecular features of triple negative breast cancer cells by genome-wide gene expression profiling analysis. <i>International Journal of Oncology</i> , 2013, 42, 478-506.	3.3	104
86	Phase I clinical trial of multi-antigen peptide vaccines therapy using cancer-testis antigens for patients with advanced or recurrent breast cancer.. <i>Journal of Clinical Oncology</i> , 2012, 30, e13037-e13037.	1.6	3
87	Mechanisms of estrogen receptor- α upregulation in breast cancers. <i>Medical Molecular Morphology</i> , 2010, 43, 193-196.	1.0	36
88	Predictive factors for anthracycline-based chemotherapy for human breast cancer. <i>Breast Cancer</i> , 2010, 17, 103-109.	2.9	15
89	Prediction of hormone sensitivity for breast cancers. <i>Breast Cancer</i> , 2010, 17, 86-91.	2.9	13
90	Association of loss of BRCA1 expression with centrosome aberration in human breast cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2009, 135, 421-430.	2.5	23

#	ARTICLE	IF	CITATIONS
91	Breast cancer risk assessment for possible tailored screening for Japanese women. <i>Breast Cancer</i> , 2009, 16, 243-247.	2.9	5
92	Growth-inhibitory effect of adiponectin via adiponectin receptor 1 on human breast cancer cells through inhibition of S-phase entry without inducing apoptosis. <i>Breast Cancer Research and Treatment</i> , 2008, 112, 405-410.	2.5	84
93	Basal-like subtype and BRCA1 dysfunction in breast cancers. <i>International Journal of Clinical Oncology</i> , 2008, 13, 395-400.	2.2	32
94	Low nuclear grade but not cell proliferation predictive of pathological complete response to docetaxel in human breast cancers. <i>Journal of Cancer Research and Clinical Oncology</i> , 2008, 134, 561-567.	2.5	11
95	Potential of Reduction in Total Tumor Volume Measured with 3D-MRI as a Prognostic Factor for Locally-Advanced Breast Cancer Patients Treated with Primary Chemotherapy. <i>Breast Journal</i> , 2008, 14, 523-531.	1.0	17
96	Quantitative assessment of mammographic density and breast cancer risk for Japanese women. <i>Breast</i> , 2008, 17, 27-35.	2.2	33
97	Cosmetic outcome and patient satisfaction after skin-sparing mastectomy for breast cancer with immediate reconstruction of the breast. <i>Surgery</i> , 2008, 143, 414-425.	1.9	96
98	Connexin26 expression is associated with aggressive phenotype in human papillary and follicular thyroid cancers. <i>Cancer Letters</i> , 2008, 262, 248-256.	7.2	29
99	Topoisomerase IIalpha-positive and BRCA1-negative phenotype: Association with favorable response to epirubicin-based regimens for human breast cancers. <i>Cancer Letters</i> , 2008, 264, 44-53.	7.2	12
100	Prognostic Significance of CD55 Expression in Breast Cancer. <i>Clinical Cancer Research</i> , 2008, 14, 4780-4786.	7.0	55
101	Clinicopathologic Analysis of Breast Cancers with PIK3CA Mutations in Japanese Women. <i>Clinical Cancer Research</i> , 2007, 13, 408-414.	7.0	138
102	Demonstration of Adiponectin Receptors 1 and 2 mRNA expression in human breast cancer cells. <i>Cancer Letters</i> , 2007, 250, 229-236.	7.2	77
103	Connexin26 expression is associated with lymphatic vessel invasion and poor prognosis in human breast cancer. <i>Breast Cancer Research and Treatment</i> , 2007, 106, 11-17.	2.5	72
104	Low LATS2 mRNA level can predict favorable response to epirubicin plus cyclophosphamide, but not to docetaxel, in breast cancers. <i>Journal of Cancer Research and Clinical Oncology</i> , 2007, 133, 501-509.	2.5	18
105	Quantitative analysis of aromatase, sulfatase and 17 β -HSD1 mRNA expression in soft tissue metastases of breast cancer. <i>Cancer Letters</i> , 2006, 243, 23-31.	7.2	19
106	Preoperative Evaluation of Residual Tumor Extent by Three-Dimensional Magnetic Resonance Imaging in Breast Cancer Patients Treated with Neoadjuvant Chemotherapy. <i>Breast Journal</i> , 2006, 12, 130-137.	1.0	77
107	High expression of ubiquitin carboxy-terminal hydrolase-L1 and -L3 mRNA predicts early recurrence in patients with invasive breast cancer. <i>Cancer Science</i> , 2006, 97, 523-529.	3.9	85
108	Association of GSTP1 CpG Islands Hypermethylation with Poor Prognosis in Human Breast Cancers. <i>Breast Cancer Research and Treatment</i> , 2006, 100, 169-176.	2.5	57

#	ARTICLE	IF	CITATIONS
109	High expression of leptin receptor mRNA in breast cancer tissue predicts poor prognosis for patients with high, but not low, serum leptin levels. <i>International Journal of Cancer</i> , 2006, 118, 1414-1419.	5.1	157
110	Prediction of response to docetaxel by immunohistochemical analysis of CYP3A4 expression in human breast cancers. <i>Breast Cancer</i> , 2005, 12, 11-15.	2.9	48
111	Differentiation of Follicular Thyroid Adenoma from Carcinoma by Means of Gene Expression Profiling with Adapter-Tagged Competitive Polymerase Chain Reaction. <i>Oncology</i> , 2005, 69, 428-435.	1.9	21
112	Prediction of Docetaxel Response in Human Breast Cancer by Gene Expression Profiling. <i>Journal of Clinical Oncology</i> , 2005, 23, 422-431.	1.6	263
113	Down-Regulation of <i>LATS1</i> and <i>LATS2</i> mRNA Expression by Promoter Hypermethylation and Its Association with Biologically Aggressive Phenotype in Human Breast Cancers. <i>Clinical Cancer Research</i> , 2005, 11, 1380-1385.	7.0	278
114	High Thioredoxin Expression Is Associated with Resistance to Docetaxel in Primary Breast Cancer. <i>Clinical Cancer Research</i> , 2005, 11, 8425-8430.	7.0	162
115	Down-Regulation of Intratumoral Aromatase Messenger RNA Levels by Docetaxel in Human Breast Cancers. <i>Clinical Cancer Research</i> , 2004, 10, 8163-8169.	7.0	14
116	Association of p53 genetic polymorphism (Arg72Pro) with estrogen receptor positive breast cancer risk in Japanese women. <i>Cancer Letters</i> , 2004, 210, 197-203.	7.2	55
117	Prognostic significance of intra-tumoral estradiol level in breast cancer patients. <i>Cancer Letters</i> , 2004, 216, 115-121.	7.2	11
118	Increased Expression of BRCA1 mRNA Predicts Favorable Response to Anthracycline-Containing Chemotherapy in Breast Cancers. <i>Breast Cancer Research and Treatment</i> , 2003, 78, 45-50.	2.5	35
119	Current status of endocrine therapy for breast cancer. <i>Breast Cancer</i> , 2003, 10, 105-111.	2.9	5
120	Polymorphisms of estrogen synthesizing and metabolizing genes and breast cancer risk in Japanese women. <i>Biomedicine and Pharmacotherapy</i> , 2003, 57, 471-481.	5.6	58
121	Association of CYP17 genetic polymorphism with intra-tumoral estradiol concentrations but not with CYP17 messenger RNA levels in breast cancer tissue. <i>Cancer Letters</i> , 2003, 195, 81-86.	7.2	14
122	Association of BRCA2 polymorphism at codon 784 (Met/Val) with breast cancer risk and prognosis. <i>Clinical Cancer Research</i> , 2003, 9, 1376-80.	7.0	26
123	Association of serum estrone levels with estrogen receptor-positive breast cancer risk in postmenopausal Japanese women. <i>Clinical Cancer Research</i> , 2003, 9, 2229-33.	7.0	26
124	Prediction of response to docetaxel by quantitative analysis of class I and III beta-tubulin isotype mRNA expression in human breast cancers. <i>Clinical Cancer Research</i> , 2003, 9, 2992-7.	7.0	86
125	Association of serum adiponectin levels with breast cancer risk. <i>Clinical Cancer Research</i> , 2003, 9, 5699-704.	7.0	317
126	Molecular classification of primary breast tumors possessing distinct prognostic properties. <i>Human Molecular Genetics</i> , 2002, 11, 199-206.	2.9	54

#	ARTICLE	IF	CITATIONS
127	HighBRCA2 mRNA expression predicts poor prognosis in breast cancer patients. International Journal of Cancer, 2002, 98, 879-882.	5.1	23
128	Mutational analysis of the class I β -tubulin gene in human breast cancer. International Journal of Cancer, 2002, 101, 46-51.	5.1	49
129	Prediction of response to docetaxel by CYP3A4 mRNA expression in breast cancer tissues. International Journal of Cancer, 2002, 97, 129-132.	5.1	93
130	Quantitative analysis of estrogen receptor- α and - β messenger RNA expression in human pancreatic cancers by real-time polymerase chain reaction. Cancer Letters, 2001, 170, 91-97.	7.2	26
131	Decreased expression ofBRCA2 mRNA predicts favorable response to docetaxel in breast cancer. International Journal of Cancer, 2001, 95, 255-259.	5.1	57
132	Association of centrosomal kinaseSTK15/BTAK mRNA expression with chromosomal instability in human breast cancers. International Journal of Cancer, 2001, 92, 370-373.	5.1	214
133	Involvement of up-regulation of 17 β -hydroxysteroid dehydrogenase type 1 in maintenance of intratumoral high estradiol levels in postmenopausal breast cancers. International Journal of Cancer, 2001, 94, 685-689.	5.1	132
134	Bone marrow micrometastases detected by RT-PCR for mammaglobin can be an alternative prognostic factor of breast cancer. Breast Cancer Research and Treatment, 2001, 67, 169-175.	2.5	27
135			