

Daryn R Michael

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1699082/publications.pdf>

Version: 2024-02-01

15
papers

593
citations

759233

12
h-index

996975

15
g-index

18
all docs

18
docs citations

18
times ranked

1197
citing authors

#	ARTICLE	IF	CITATIONS
1	IL-33 Reduces Macrophage Foam Cell Formation. <i>Journal of Immunology</i> , 2010, 185, 1222-1229.	0.8	165
2	ERK Is Integral to the IFN- γ -Mediated Activation of STAT1, the Expression of Key Genes Implicated in Atherosclerosis, and the Uptake of Modified Lipoproteins by Human Macrophages. <i>Journal of Immunology</i> , 2010, 185, 3041-3048.	0.8	89
3	The Human Hyaluronan Synthase 2 (HAS2) Gene and Its Natural Antisense RNA Exhibit Coordinated Expression in the Renal Proximal Tubular Epithelial Cell. <i>Journal of Biological Chemistry</i> , 2011, 286, 19523-19532.	3.4	78
4	Sp1 and Sp3 Mediate Constitutive Transcription of the Human Hyaluronan Synthase 2 Gene. <i>Journal of Biological Chemistry</i> , 2006, 281, 18043-18050.	3.4	42
5	Differential regulation of macropinocytosis in macrophages by cytokines: Implications for foam cell formation and atherosclerosis. <i>Cytokine</i> , 2013, 64, 357-361.	3.2	35
6	Liver X Receptors, Atherosclerosis and Inflammation. <i>Current Atherosclerosis Reports</i> , 2012, 14, 284-293.	4.8	32
7	Eicosapentaenoic Acid and Docosahexaenoic Acid Regulate Modified LDL Uptake and Macropinocytosis in Human Macrophages. <i>Lipids</i> , 2011, 46, 1053-1061.	1.7	30
8	Probiotics reduce self-reported symptoms of upper respiratory tract infection in overweight and obese adults: should we be considering probiotics during viral pandemics?. <i>Gut Microbes</i> , 2021, 13, 1-9.	9.8	28
9	A Unique Combination of Nutritionally Active Ingredients Can Prevent Several Key Processes Associated with Atherosclerosis In Vitro. <i>PLoS ONE</i> , 2016, 11, e0151057.	2.5	24
10	The expression of a disintegrin and metalloproteinase with thrombospondin motifs 4 in human macrophages is inhibited by the anti-atherogenic cytokine transforming growth factor- β 2 and requires Smads, p38 mitogen-activated protein kinase and c-Jun. <i>International Journal of Biochemistry and Cell Biology</i> , 2011, 43, 805-811.	2.8	18
11	The Phosphoinositide 3-Kinase Signaling Pathway is Involved in the Control of Modified Low-Density Lipoprotein Uptake by Human Macrophages. <i>Lipids</i> , 2015, 50, 253-260.	1.7	13
12	Macrophages, lipid metabolism and gene expression in atherogenesis: a therapeutic target of the future?. <i>Clinical Lipidology</i> , 2012, 7, 37-48.	0.4	12
13	Protective effects of a unique combination of nutritionally active ingredients on risk factors and gene expression associated with atherosclerosis in C57BL/6J mice fed a high fat diet. <i>Food and Function</i> , 2021, 12, 3657-3671.	4.6	12
14	Protein Kinase C Is Involved in the Induction of ATP-Binding Cassette Transporter A1 Expression by Liver X Receptor/Retinoid X Receptor Agonist in Human Macrophages. <i>Journal of Cellular Biochemistry</i> , 2015, 116, 2032-2038.	2.6	9
15	Identifying the factors influencing outcome in probiotic studies in overweight and obese patients: host or microbiome?. <i>Gut</i> , 2021, 70, 225-226.	12.1	6