
## Rebecca Kandiyali

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1709050/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Telephone triage for management of same-day consultation requests in general practice (the ESTEEM) Tj ETQq1	1 0.78431	4 rgBT /Over
	1859-1868.	10.7	100
2	Clinical effectiveness and cost-effectiveness of physiotherapy and occupational therapy versus no therapy in mild to moderate Parkinson's disease: a large pragmatic randomised controlled trial (PD) Tj ETQqC	) 0 <b>208</b> gBT /	Oøerlock 10

3	Mycophenolate Mofetil for First-Line Treatment of Immune Thrombocytopenia. New England Journal of Medicine, 2021, 385, 885-895.	27.0	53
4	The clinical effectiveness and cost-effectiveness of telephone triage for managing same-day consultation requests in general practice: a cluster randomised controlled trial comparing general practitioner-led and nurse-led management systems with usual care (the ESTEEM trial). Health Technology Assessment, 2015, 19, 1-212.	2.8	50
5	Limit of detection of troponin discharge strategy versus usual care: randomised controlled trial. Heart, 2020, 106, 1586-1594.	2.9	39
6	A pilot randomised trial to assess the methods and procedures for evaluating the clinical effectiveness and cost-effectiveness of Exercise Assisted Reduction then Stop (EARS) among disadvantaged smokers. Health Technology Assessment, 2014, 18, 1-324.	2.8	36
7	Physical activity and the prevention, reduction, and treatment of alcohol and other drug use across the lifespan (The PHASE review): A systematic review. Mental Health and Physical Activity, 2020, 19, 100360.	1.8	28
8	Involving Patients in Health Economics Research: "The PACTS Principles― Patient, 2021, 14, 429-434.	2.7	23
9	Trial protocol: a multicentre randomised trial of first-line treatment pathways for newly diagnosed immune thrombocytopenia: standard steroid treatment versus combined steroid and mycophenolate. The FLIGHT trial. BMJ Open, 2018, 8, e024427.	1.9	20
10	Considerations for Individual-Level Versus Whole-School Physical Activity Interventions: Stakeholder Perspectives. International Journal of Environmental Research and Public Health, 2021, 18, 7628.	2.6	16
11	Study protocol for †The Project About Loneliness and Social networks (PALS)': a pragmatic, randomised trial comparing a facilitated social network intervention (Genie) with a wait-list control for lonely and socially isolated people. BMJ Open, 2019, 9, e028718.	1.9	13
12	Physical activity and the prevention, reduction, and treatment of alcohol and/or substance use across the lifespan (The PHASE review): protocol for a systematic review. Systematic Reviews, 2018, 7, 9.	5.3	12
13	Working with Patients and Members of the Public: Informing Health Economics in Child Health Research. PharmacoEconomics - Open, 2019, 3, 133-141.	1.8	9
14	Protocol for a cluster randomised controlled trial of a Peer-Led physical Activity iNtervention for Adolescent girls (PLAN-A). BMC Public Health, 2019, 19, 644.	2.9	9
15	Randomised controlled trial of the Limit of Detection of Troponin and ECG Discharge (LoDED) strategy versus usual care in adult patients with chest pain attending the emergency department: study protocol. BMJ Open, 2018, 8, e025339.	1.9	8
16	Assessing the construct validity and responsiveness of Preference-Based Measures (PBMs) in cataract surgery patients. Quality of Life Research, 2020, 29, 1935-1946.	3.1	8
17	Predictive values of referrals for transient ischaemic attack from first-contact health care: a systematic review. British Journal of General Practice, 2017, 67, e871-e880.	1.4	6
18	Effectiveness and cost-effectiveness of the PLAN-A intervention, a peer led physical activity program for adolescent girls: results of a cluster randomised controlled trial. International Journal of Behavioral Nutrition and Physical Activity, 2021, 18, 63.	4.6	6

Rebecca Kandiyali

#	Article	IF	CITATIONS
19	The management of small area burns and unexpected illness after burn in children under five years of age — A costing study in the English healthcare setting. Burns, 2018, 44, 188-194.	1.9	5
20	TULIP: a randomised controlled trial of surgical versus non-surgical treatment of lateral compression injuries of the pelvis with complete sacral fractures (LC1) in the non-fragility fracture patient—a feasibility study protocol. BMJ Open, 2020, 10, e036588.	1.9	4
21	Costs and Cost-Effectiveness of User-Testing of Health Professionals' Guidelines to Reduce the Frequency of Intravenous Medicines Administration Errors by Nurses in the United Kingdom: A Probabilistic Model Based on Voriconazole Administration. Applied Health Economics and Health Policy. 2022, 20. 91-104.	2.1	3
22	Physical Activity and Psychosocial Characteristics of the Peer Supporters in the PLAN-A Study—A Latent Class Analysis. International Journal of Environmental Research and Public Health, 2020, 17, 7980.	2.6	2
23	Cost-effectiveness and value of information analysis of a low-friction environment following skin graft in patients with burn injury. Pilot and Feasibility Studies, 2020, 6, 8.	1.2	2
24	The SILKIE (Skin graftIng Low friKtIon Environment) study: a non-randomised proof-of-concept and feasibility study on the impact of low-friction nursing environment on skin grafting success rates in adult and paediatric burns. BMJ Open, 2018, 8, e021886.	1.9	2
25	Peer-led physical activity intervention for girls aged 13 to 14 years: PLAN-A cluster RCT. Public Health Research, 2022, 10, 1-154.	1.3	1