

# Watcharin Loilome

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1711743/publications.pdf>

Version: 2024-02-01

118  
papers

2,874  
citations

136950

32  
h-index

223800

46  
g-index

124  
all docs

124  
docs citations

124  
times ranked

3472  
citing authors

#	ARTICLE	IF	CITATIONS
1	Increased activation of PI3K/AKT signaling pathway is associated with cholangiocarcinoma metastasis and PI3K/mTOR inhibition presents a possible therapeutic strategy. <i>Tumor Biology</i> , 2013, 34, 3637-3648.	1.8	100
2	Cohort profile: cholangiocarcinoma screening and care program (CASCAP). <i>BMC Cancer</i> , 2015, 15, 459.	2.6	93
3	Activated macrophages promote Wnt/ $\beta$ -catenin signaling in cholangiocarcinoma cells. <i>Tumor Biology</i> , 2014, 35, 5357-5367.	1.8	87
4	Mechanisms of oxysterol-induced carcinogenesis. <i>Lipids in Health and Disease</i> , 2011, 10, 44.	3.0	69
5	Cholangiocarcinoma: a guide for the nonspecialist. <i>International Journal of General Medicine</i> , 2019, Volume 12, 13-23.	1.8	67
6	Glioblastoma cell growth is suppressed by disruption of fibroblast growth factor pathway signaling. <i>Journal of Neuro-Oncology</i> , 2009, 94, 359-366.	2.9	65
7	Circulating miR-192 in liver fluke-associated cholangiocarcinoma patients: a prospective prognostic indicator. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2014, 21, 864-872.	2.6	65
8	Prolonged oxidative stress down-regulates Early B cell factor 1 with inhibition of its tumor suppressive function against cholangiocarcinoma genesis. <i>Redox Biology</i> , 2018, 14, 637-644.	9.0	62
9	Resveratrol interrupts the pro-invasive communication between cancer associated fibroblasts and cholangiocarcinoma cells. <i>Cancer Letters</i> , 2018, 430, 160-171.	7.2	60
10	Altered gene expression in <i>Opisthorchis viverrini</i> -associated cholangiocarcinoma in hamster model. <i>Molecular Carcinogenesis</i> , 2006, 45, 279-287.	2.7	59
11	Myristoylated alanine-rich C kinase substrate phosphorylation promotes cholangiocarcinoma cell migration and metastasis via the protein kinase C-dependent pathway. <i>Cancer Science</i> , 2010, 101, 658-665.	3.9	59
12	Histological confirmation of periductal fibrosis from ultrasound diagnosis in cholangiocarcinoma patients. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2014, 21, 316-322.	2.6	58
13	Quantitative Changes in Tumor-Associated M2 Macrophages Characterize Cholangiocarcinoma and their Association with Metastasis. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 16, 3043-3050.	1.2	58
14	Tumor necrosis factor- $\alpha$ (TNF- $\alpha$ ) stimulates the epithelial-mesenchymal transition regulator Snail in cholangiocarcinoma. <i>Medical Oncology</i> , 2012, 29, 3083-3091.	2.5	57
15	CD44 variant-dependent redox status regulation in liver fluke-associated cholangiocarcinoma: A target for cholangiocarcinoma treatment. <i>Cancer Science</i> , 2016, 107, 991-1000.	3.9	57
16	Urinary microRNA-192 and microRNA-21 as potential indicators for liver fluke-associated cholangiocarcinoma risk group. <i>Parasitology International</i> , 2017, 66, 479-485.	1.3	52
17	A Comprehensive Public Health Conceptual Framework and Strategy to Effectively Combat Cholangiocarcinoma in Thailand. <i>PLoS Neglected Tropical Diseases</i> , 2016, 10, e0004293.	3.0	51
18	Current Perspectives on Opisthorchiasis Control and Cholangiocarcinoma Detection in Southeast Asia. <i>Frontiers in Medicine</i> , 2018, 5, 117.	2.6	51

#	ARTICLE	IF	CITATIONS
19	Advances in the Diagnosis of Human Opisthorchiasis: Development of Opisthorchis viverrini Antigen Detection in Urine. PLoS Neglected Tropical Diseases, 2015, 9, e0004157.	3.0	50
20	Inhibition of <scp> </scp>-type amino acid transporter 1 activity as a new therapeutic target for cholangiocarcinoma treatment. Tumor Biology, 2017, 39, 101042831769454.	1.8	50
21	High Expression of HIF-1 $\pm$ , BNIP3 and PI3KC3: Hypoxia-Induced Autophagy Predicts Cholangiocarcinoma Survival and Metastasis. Asian Pacific Journal of Cancer Prevention, 2014, 15, 5873-5878.	1.2	49
22	Dihydroartemisinin induces apoptosis and autophagy $\hat{\epsilon}$ dependent cell death in cholangiocarcinoma through a DAPK1 $\hat{\epsilon}$ BECLIN1 pathway. Molecular Carcinogenesis, 2018, 57, 1735-1750.	2.7	48
23	Characterization of 5-Fluorouracil-Resistant Cholangiocarcinoma Cell Lines. Chemotherapy, 2008, 54, 343-351.	1.6	47
24	PRKAR1A is overexpressed and represents a possible therapeutic target in human cholangiocarcinoma. International Journal of Cancer, 2011, 129, 34-44.	5.1	47
25	Loss of E-cadherin promotes migration and invasion of cholangiocarcinoma cells and serves as a potential marker of metastasis. Tumor Biology, 2014, 35, 8645-8652.	1.8	47
26	BMP-7 blocks the effects of TGF- $\hat{2}$ -induced EMT in cholangiocarcinoma. Tumor Biology, 2014, 35, 9667-9676.	1.8	43
27	Platelet-derived growth factor may be a potential diagnostic and prognostic marker for cholangiocarcinoma. Tumor Biology, 2012, 33, 1785-1802.	1.8	38
28	<scp>STAT</scp>s profiling reveals predominantly $\hat{\epsilon}$ activated <scp>STAT3</scp> in cholangiocarcinoma genesis and progression. Journal of Hepato-Biliary-Pancreatic Sciences, 2014, 21, 767-776.	2.6	38
29	Opisthorchiasis and cholangiocarcinoma in Southeast Asia: an unresolved problem. International Journal of General Medicine, 2017, Volume 10, 227-237.	1.8	38
30	Liver fluke-induced hepatic oxysterols stimulate DNA damage and apoptosis in cultured human cholangiocytes. Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis, 2012, 731, 48-57.	1.0	36
31	Tumor necrosis factor $\hat{\pm}$ modulates epithelial mesenchymal transition mediators <scp>ZEB2</scp> and <scp>S100A4</scp> to promote cholangiocarcinoma progression. Journal of Hepato-Biliary-Pancreatic Sciences, 2014, 21, 703-711.	2.6	36
32	Opisthorchis viverrini Infection Activates the PI3K/AKT/PTEN and Wnt/ $\hat{2}$ -catenin Signaling Pathways in a Cholangiocarcinogenesis Model. Asian Pacific Journal of Cancer Prevention, 2015, 15, 10463-10468.	1.2	36
33	Survey of activated kinase proteins reveals potential targets for cholangiocarcinoma treatment. Tumor Biology, 2013, 34, 3519-3528.	1.8	34
34	Cancer-Associated Fibroblast-Derived IL-6 Determines Unfavorable Prognosis in Cholangiocarcinoma by Affecting Autophagy-Associated Chemoresponse. Cancers, 2021, 13, 2134.	3.7	33
35	Increased EphB2 expression predicts cholangiocarcinoma metastasis. Tumor Biology, 2014, 35, 10031-10041.	1.8	32
36	Xanthohumol inhibits STAT3 activation pathway leading to growth suppression and apoptosis induction in human cholangiocarcinoma cells. Oncology Reports, 2016, 35, 2065-2072.	2.6	32

#	ARTICLE	IF	CITATIONS
37	Risk biomarkers for assessment and chemoprevention of liver fluke-associated cholangiocarcinoma. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2014, 21, 309-315.	2.6	31
38	Upregulation of transferrin receptor-1 induces cholangiocarcinoma progression via induction of labile iron pool. <i>Tumor Biology</i> , 2017, 39, 101042831771765.	1.8	31
39	Expression of oxysterol binding protein isoforms in opisthorchiasis-associated cholangiocarcinoma: A potential molecular marker for tumor metastasis. <i>Parasitology International</i> , 2012, 61, 136-139.	1.3	28
40	Establishment of cholangiocarcinoma cell lines from patients in the endemic area of liver fluke infection in Thailand. <i>Tumor Biology</i> , 2017, 39, 101042831772592.	1.8	27
41	Cytokines released from activated human macrophages induce epithelial mesenchymal transition markers of cholangiocarcinoma cells. <i>Asian Pacific Journal of Cancer Prevention</i> , 2012, 13 Suppl, 115-8.	1.2	25
42	Comparing the performance of urine and copro-antigen detection in evaluating <i>Opisthorchis viverrini</i> infection in communities with different transmission levels in Northeast Thailand. <i>PLoS Neglected Tropical Diseases</i> , 2019, 13, e0007186.	3.0	24
43	Potential role of HIF-1-responsive microRNA210/HIF3 axis on gemcitabine resistance in cholangiocarcinoma cells. <i>PLoS ONE</i> , 2018, 13, e0199827.	2.5	22
44	Progranulin modulates cholangiocarcinoma cell proliferation, apoptosis, and motility via the PI3K/pAkt pathway. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 395-408.	2.0	21
45	Urine proteomics study reveals potential biomarkers for the differential diagnosis of cholangiocarcinoma and periductal fibrosis. <i>PLoS ONE</i> , 2019, 14, e0221024.	2.5	21
46	Anti-apoptotic phenotypes of cholestan-3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -triol-resistant human cholangiocytes: Characteristics contributing to the genesis of cholangiocarcinoma. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2013, 138, 368-375.	2.5	20
47	Evaluation of anticancer potential of Thai medicinal herb extracts against cholangiocarcinoma cell lines. <i>PLoS ONE</i> , 2019, 14, e0216721.	2.5	20
48	Nimotuzumab Inhibits Cholangiocarcinoma Cell Metastasis via Suppression of the Epithelial-Mesenchymal Transition Process. <i>Anticancer Research</i> , 2017, 37, 3591-3597.	1.1	20
49	Oxidized alpha-1 antitrypsin as a predictive risk marker of opisthorchiasis-associated cholangiocarcinoma. <i>Tumor Biology</i> , 2013, 34, 695-704.	1.8	19
50	Increase in L-type amino acid transporter 1 expression during cholangiocarcinogenesis caused by liver fluke infection and its prognostic significance. <i>Parasitology International</i> , 2017, 66, 471-478.	1.3	19
51	Diagnostic performance of urinary IgG antibody detection: A novel approach for population screening of strongyloidiasis. <i>PLoS ONE</i> , 2018, 13, e0192598.	2.5	19
52	Discovery and Qualification of Serum Protein Biomarker Candidates for Cholangiocarcinoma Diagnosis. <i>Journal of Proteome Research</i> , 2019, 18, 3305-3316.	3.7	18
53	Discovery of Serotransferrin Glycoforms: Novel Markers for Diagnosis of Liver Periductal Fibrosis and Prediction of Cholangiocarcinoma. <i>Biomolecules</i> , 2019, 9, 538.	4.0	17
54	Smartphone-based fluorescent ELISA with simple fluorescent enhancement strategy for <i>Opisthorchis viverrini</i> (Ov) antigen detection in urine samples. <i>Sensors and Actuators B: Chemical</i> , 2021, 348, 130705.	7.8	17

#	ARTICLE	IF	CITATIONS
55	Genetic and environmental determinants of risk for cholangiocarcinoma in Thailand. <i>World Journal of Gastrointestinal Pathophysiology</i> , 2014, 5, 570.	1.0	17
56	Synchrotron FTIR microspectroscopy revealed apoptosis-induced biomolecular changes of cholangiocarcinoma cells treated with ursolic acid. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2020, 1864, 129708.	2.4	16
57	Overexpression of a panel of cancer stem cell markers enhances the predictive capability of the progression and recurrence in the early stage cholangiocarcinoma. <i>Journal of Translational Medicine</i> , 2020, 18, 64.	4.4	16
58	Targeting Fatty Acid Synthase Modulates Metabolic Pathways and Inhibits Cholangiocarcinoma Cell Progression. <i>Frontiers in Pharmacology</i> , 2021, 12, 696961.	3.5	16
59	Teleconsultation ultrasonography: a new weapon to combat cholangiocarcinoma. <i>ESMO Open</i> , 2017, 2, e000231.	4.5	15
60	Zileuton suppresses cholangiocarcinoma cell proliferation and migration through inhibition of the Akt signaling pathway. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 7019-7029.	2.0	15
61	Smartphone-based portable fluorescence sensor with gold nanoparticle mediation for selective detection of nitrite ions. <i>Food Chemistry</i> , 2022, 384, 132478.	8.2	15
62	Opisthorchis viverrini-antigen induces expression of MARCKS during inflammation-associated cholangiocarcinogenesis. <i>Parasitology International</i> , 2012, 61, 140-144.	1.3	14
63	Upregulation of endothelial nitric oxide synthase (eNOS) and its upstream regulators in Opisthorchis viverrini associated cholangiocarcinoma and its clinical significance. <i>Parasitology International</i> , 2017, 66, 486-493.	1.3	14
64	Characterisation of the Urinary Metabolic Profile of Liver Fluke-Associated Cholangiocarcinoma. <i>Journal of Clinical and Experimental Hepatology</i> , 2019, 9, 657-675.	0.9	14
65	Sulfasalazine modifies metabolic profiles and enhances cisplatin chemosensitivity on cholangiocarcinoma cells in in vitro and in vivo models. <i>Cancer &amp; Metabolism</i> , 2021, 9, 11.	5.0	14
66	Increased expression of TLR-2, COX-2, and SOD-2 genes in the peripheral blood leukocytes of opisthorchiasis patients induced by Opisthorchis viverrini antigen. <i>Parasitology Research</i> , 2012, 110, 1969-1977.	1.6	13
67	Imbalanced adaptive responses associated with microsatellite instability in cholangiocarcinoma. <i>Oncology Letters</i> , 2017, 13, 639-646.	1.8	13
68	Inhibition of endothelial nitric oxide synthase in cholangiocarcinoma cell lines – a new strategy for therapy. <i>FEBS Open Bio</i> , 2018, 8, 513-522.	2.3	13
69	The Socioeconomic Burden of Cholangiocarcinoma Associated With Opisthorchis viverrini Sensu Lato Infection in Northeast Thailand. <i>Advances in Parasitology</i> , 2018, 102, 141-163.	3.2	13
70	A panel of protein kinase high expression is associated with postoperative recurrence in cholangiocarcinoma. <i>BMC Cancer</i> , 2020, 20, 154.	2.6	13
71	Chloroquine Exerts Anti-metastatic Activities Under Hypoxic Conditions in Cholangiocarcinoma Cells. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 16, 2031-2035.	1.2	13
72	Immunosuppressive Prednisolone Enhances Early Cholangiocarcinoma in Syrian Hamsters with Liver Fluke Infection and Administration of N-nitrosodimethylamine. <i>Pathology and Oncology Research</i> , 2013, 19, 55-62.	1.9	12

#	ARTICLE	IF	CITATIONS
73	Characterisation of the Serum Metabolic Signature of Cholangiocarcinoma in a United Kingdom Cohort. <i>Journal of Clinical and Experimental Hepatology</i> , 2020, 10, 17-29.	0.9	12
74	AuNPs-LISA, an efficient detection assay for <i>Opisthorchis viverrini</i> (Ov) antigen in urine. <i>Talanta</i> , 2020, 209, 120592.	5.5	12
75	Antifibrotic effect of xanthohumol in combination with praziquantel is associated with altered redox status and reduced iron accumulation during liver fluke-associated cholangiocarcinogenesis. <i>PeerJ</i> , 2018, 6, e4281.	2.0	12
76	PRKAR1A overexpression is associated with increased ECPKA autoantibody in liver fluke-associated cholangiocarcinoma: application for assessment of the risk group. <i>Tumor Biology</i> , 2012, 33, 2289-2298.	1.8	11
77	Combination of Praziquantel and Aspirin Minimizes Liver Pathology of Hamster <i>Opisthorchis viverrini</i> Infection Associated Cholangiocarcinoma. <i>Pathology and Oncology Research</i> , 2016, 22, 57-65.	1.9	11
78	&lt;p&gt;In vitro and in vivo Anti-Tumor Effects of Pan-HER Inhibitor Varlitinib on Cholangiocarcinoma Cell Lines&lt;/p&gt;. <i>Drug Design, Development and Therapy</i> , 2020, Volume 14, 2319-2334.	4.3	11
79	A comparison of the proportion of early stage cholangiocarcinoma found in an ultrasound-screening program compared to walk-in patients. <i>Hpb</i> , 2020, 22, 874-883.	0.3	11
80	Integration of global metabolomics and lipidomics approaches reveals the molecular mechanisms and the potential biomarkers for postoperative recurrence in early-stage cholangiocarcinoma. <i>Cancer &amp; Metabolism</i> , 2021, 9, 30.	5.0	11
81	Upregulation of TCTP is associated with cholangiocarcinoma progression and metastasis. <i>Oncology Letters</i> , 2017, 14, 5973-5979.	1.8	10
82	A fluorescence AuNPs-LISA: A new approach for <i>Opisthorchis viverrini</i> (Ov) antigen detection with a simple fluorescent enhancement strategy by surfactant micelle in urine samples. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2021, 254, 119633.	3.9	10
83	Association of CYP39A1, RUNX2 and Oxidized Alpha-1 Antitrypsin Expression in Relation to Cholangiocarcinoma Progression. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 15, 10187-10192.	1.2	10
84	Anti-inflammatory agents suppress the prostaglandin E2 production and migration ability of cholangiocarcinoma cell lines. <i>Asian Pacific Journal of Cancer Prevention</i> , 2012, 13 Suppl, 47-51.	1.2	10
85	PGE2 signaling and its biosynthesis-related enzymes in cholangiocarcinoma progression. <i>Tumor Biology</i> , 2014, 35, 8051-8064.	1.8	9
86	Circulating TGF- $\beta$ 1 as the potential epithelial mesenchymal transition-biomarker for diagnosis of cholangiocarcinoma. <i>Journal of Gastrointestinal Oncology</i> , 2020, 11, 304-318.	1.4	9
87	CD44 modulates metabolic pathways and altered ROS-mediated Akt signal promoting cholangiocarcinoma progression. <i>PLoS ONE</i> , 2021, 16, e0245871.	2.5	9
88	Potential of Selenium Compounds as New Anticancer Agents for Cholangiocarcinoma. <i>Anticancer Research</i> , 2016, 36, 5981-5988.	1.1	9
89	<i>ARID1A</i> alterations and their clinical significance in cholangiocarcinoma. <i>PeerJ</i> , 2020, 8, e10464.	2.0	9
90	In vitro and molecular chemosensitivity in human cholangiocarcinoma tissues. <i>PLoS ONE</i> , 2019, 14, e0222140.	2.5	8

#	ARTICLE	IF	CITATIONS
91	Mass Spectrometry: A Guide for the Clinician. <i>Journal of Clinical and Experimental Hepatology</i> , 2019, 9, 597-606.	0.9	8
92	Diagnostic and Prognostic Value of Circulating Cell-Free DNA for Cholangiocarcinoma. <i>Diagnostics</i> , 2021, 11, 999.	2.6	8
93	Milk fat globule epidermal growth factor 8 serves a novel biomarker of opisthorchiasis-associated cholangiocarcinoma. <i>Tumor Biology</i> , 2014, 35, 1985-1995.	1.8	7
94	Bacterial challenge-associated metabolic phenotypes in <i>Hermetia illucens</i> defining nutritional and functional benefits. <i>Scientific Reports</i> , 2021, 11, 23316.	3.3	7
95	Inhibitory effect of NVP-BKM120 on cholangiocarcinoma cell growth. <i>Oncology Letters</i> , 2018, 16, 1627-1633.	1.8	6
96	Serum IgG as a Marker for <i>Opisthorchis viverrini</i> -Associated Cholangiocarcinoma Correlated with HER2 Overexpression. <i>International Journal of General Medicine</i> , 2020, Volume 13, 1271-1283.	1.8	6
97	Modification of the AJCC/UICC 8th edition staging system for intrahepatic cholangiocarcinoma: proposal for an alternative staging system from cholangiocarcinoma-prevalent Northeast Thailand. <i>Hpb</i> , 2022, 24, 1944-1956.	0.3	6
98	Performance of Mini Parasep® SF stool concentrator kit, Kato-Katz, and formalin-ethyl acetate concentration methods for diagnosis of opisthorchiasis in Northeast Thailand. <i>Parasites and Vectors</i> , 2022, 15, .	2.5	6
99	Suppression of 14-3-3 $\eta$ in cholangiocarcinoma cells inhibits proliferation through attenuated Akt activity, enhancing chemosensitivity to gemcitabine. <i>Oncology Letters</i> , 2018, 15, 347-353.	1.8	5
100	PRIMA-1 <sup>MET</sup> Induces Cellular Senescence and Apoptotic Cell Death in Cholangiocarcinoma Cells. <i>Cancer Genomics and Proteomics</i> , 2019, 16, 543-552.	2.0	5
101	Curative effect of xanthohumol supplementation during liver fluke-associated cholangiocarcinogenesis: Potential involvement of Autophagy. <i>Journal of Traditional and Complementary Medicine</i> , 2020, 10, 230-235.	2.7	5
102	High Levels of Serum IgG for <i>Opisthorchis viverrini</i> and CD44 Expression Predict Worse Prognosis for Cholangiocarcinoma Patients after Curative Resection. <i>International Journal of General Medicine</i> , 2021, Volume 14, 2191-2204.	1.8	5
103	Arctigenin inhibits cholangiocarcinoma progression by regulating cell migration and cell viability via the N-cadherin and apoptosis pathway. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2021, 394, 2049-2059.	3.0	5
104	Therapeutic targeting of ARID1A and PI3K/AKT pathway alterations in cholangiocarcinoma. <i>PeerJ</i> , 2022, 10, e12750.	2.0	5
105	Metabolic Phenotyping Predicts Gemcitabine and Cisplatin Chemosensitivity in Patients With Cholangiocarcinoma. <i>Frontiers in Public Health</i> , 2022, 10, 766023.	2.7	5
106	Impaired antioxidant enzyme activity and increased DNA repair enzyme expression in hamster liver tissues related to cholangiocarcinoma development. <i>Asian Pacific Journal of Cancer Prevention</i> , 2012, 13 Suppl, 59-64.	1.2	5
107	Promoter hypermethylation of early B cell factor 1 (EBF1) is associated with cholangiocarcinoma progression. <i>Journal of Cancer</i> , 2021, 12, 2673-2686.	2.5	4
108	Metabolic Profiling of Praziquantel-mediated Prevention of <i>Opisthorchis viverrini</i> -induced Cholangiocyte Transformation in the Hamster Model of Cholangiocarcinoma. <i>Cancer Genomics and Proteomics</i> , 2021, 18, 29-42.	2.0	4



#	ARTICLE	IF	CITATIONS
109	Elevated Levels of Urinary 8-oxodG Correlate with Persistent Periductal Fibrosis after Praziquantel Treatment in Chronic Opisthorchiasis. <i>American Journal of Tropical Medicine and Hygiene</i> , 2018, 98, 1763-1769.	1.4	4
110	A Promising Serum Autoantibody Marker, Anti-Heat Shock Protein 90 $\alpha$ , for Cholangiocarcinoma. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 16, 5779-5785.	1.2	4
111	Therapeutic challenges at the preclinical level for targeted drug development for <i>Opisthorchis viverrini</i> -associated cholangiocarcinoma. <i>Expert Opinion on Investigational Drugs</i> , 2021, 30, 985-1006.	4.1	3
112	<sup>1</sup> H NMR metabolic phenotyping of <i>Dipterocarpus alatus</i> as a novel tool for age and growth determination. <i>PLoS ONE</i> , 2020, 15, e0243432.	2.5	3
113	Evaluation of a short term effect of praziquantel treatment in opisthorchiasis-induced hepatobiliary inflammation by urinary 8-oxodG. <i>Acta Tropica</i> , 2019, 189, 124-128.	2.0	2
114	Metabolic Changes of Cholangiocarcinoma Cells in Response to Coniferyl Alcohol Treatment. <i>Biomolecules</i> , 2021, 11, 476.	4.0	2
115	Mapping of population disparities in the cholangiocarcinoma urinary metabolome. <i>Scientific Reports</i> , 2021, 11, 21286.	3.3	2
116	Suppression of PRKAR1A expression enhances anti-proliferative and apoptotic effects of protein kinase inhibitors and chemotherapeutic drugs on cholangiocarcinoma cells. <i>Asian Pacific Journal of Cancer Prevention</i> , 2012, 13 Suppl, 143-7.	1.2	2
117	Protein Kinases as Targets for <i>Opisthorchis viverrini</i> - Associated Cholangiocarcinoma Therapy. <i>Current Pharmaceutical Design</i> , 2017, 23, 4281-4289.	1.9	1
118	Lipidomic Analyses Uncover Apoptotic and Inhibitory Effects of Pyrvinium Pamoate on Cholangiocarcinoma Cells via Mitochondrial Membrane Potential Dysfunction. <i>Frontiers in Public Health</i> , 2021, 9, 766455.	2.7	1