

Masaki Izumo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1723587/publications.pdf>

Version: 2024-02-01

19
papers

249
citations

1163117

8
h-index

940533

16
g-index

21
all docs

21
docs citations

21
times ranked

404
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Results of PROspect trial to Elucidate the utility of Echocardiography-based Cardiac Output in acute heart failure (PREDICT). <i>Journal of Cardiology</i> , 2022, 80, 218-225. | 1.9 | 2 |
| 2 | Dynamic Secondary Mitral Regurgitation: Current Evidence and Challenges for the Future. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, 883450. | 2.4 | 1 |
| 3 | Prognostic impact of transcatheter mitral valve repair in patients with exercise-induced secondary mitral regurgitation. <i>European Heart Journal Cardiovascular Imaging</i> , 2021, 22, 530-538. | 1.2 | 12 |
| 4 | McConnell's sign assessed by point-of-care cardiac ultrasound associated with in-hospital mortality of COVID-19 patients with respiratory failure. <i>Journal of Echocardiography</i> , 2021, 19, 67-69. | 0.8 | 4 |
| 5 | The efficiency of exercise stress echocardiography for evaluating symptomatic mitral regurgitation. <i>European Heart Journal - Case Reports</i> , 2021, 5, ytab006. | 0.6 | 0 |
| 6 | Prognostic significance of right ventricular function during exercise in asymptomatic/minimally symptomatic patients with nonobstructive hypertrophic cardiomyopathy. <i>Echocardiography</i> , 2021, 38, 916-923. | 0.9 | 3 |
| 7 | Resting echocardiographic predictors for true-severe aortic stenosis in patients with low-gradient severe aortic stenosis: A dobutamine stress echocardiography study. <i>Echocardiography</i> , 2021, 38, 1731-1740. | 0.9 | 1 |
| 8 | Geometry of Tricuspid Valve Apparatus in Patients with Mitral Regurgitation due to Fibroelastic Deficiency versus Barlow Disease: A Real-Time Three-dimensional Transesophageal Echocardiography Study. <i>Journal of the American Society of Echocardiography</i> , 2020, 33, 1095-1105. | 2.8 | 5 |
| 9 | Value of Transvalvular Flow Rate during Exercise in Asymptomatic Patients with Aortic Stenosis. <i>Journal of the American Society of Echocardiography</i> , 2020, 33, 438-448. | 2.8 | 10 |
| 10 | Prognostic value of exercise stress echocardiography in patients with secondary mitral regurgitation: a long-term follow-up study. <i>Journal of Echocardiography</i> , 2019, 17, 147-156. | 0.8 | 14 |
| 11 | Geometry of the left ventricular outflow tract assessed by 3D TEE in patients with aortic stenosis: impact of upper septal hypertrophy on measurements of Doppler-derived left ventricular stroke volume. <i>Journal of Echocardiography</i> , 2018, 16, 162-172. | 0.8 | 7 |
| 12 | Prognostic implications in patients with symptomatic aortic stenosis and preserved ejection fraction: Japanese multicenter aortic stenosis, retrospective (JUST-R) registry. <i>Journal of Cardiology</i> , 2017, 69, 110-118. | 1.9 | 7 |
| 13 | Effect of aortic regurgitant jet direction on mitral valve leaflet remodeling: a real-time three-dimensional transesophageal echocardiography study. <i>Scientific Reports</i> , 2017, 7, 8884. | 3.3 | 7 |
| 14 | Reliability of Aortic Stenosis Severity Classified by 3-Dimensional Echocardiography in the Prediction of Cardiovascular Events. <i>American Journal of Cardiology</i> , 2016, 118, 410-417. | 1.6 | 9 |
| 15 | Exercise echocardiography for structural heart disease. <i>Journal of Echocardiography</i> , 2016, 14, 21-29. | 0.8 | 3 |
| 16 | Value of anatomical aortic valve area using real-time three-dimensional transoesophageal echocardiography in patients with aortic stenosis: a comparison between tricuspid and bicuspid aortic valves. <i>European Heart Journal Cardiovascular Imaging</i> , 2015, 16, 1120-1128. | 1.2 | 9 |
| 17 | Comparison of Left Ventricular Outflow Geometry and Aortic Valve Area in Patients With Aortic Stenosis by 2-Dimensional Versus 3-Dimensional Echocardiography. <i>American Journal of Cardiology</i> , 2012, 109, 1626-1631. | 1.6 | 94 |
| 18 | Changes in mitral regurgitation and left ventricular geometry during exercise affect exercise capacity in patients with systolic heart failure. <i>European Journal of Echocardiography</i> , 2011, 12, 54-60. | 2.3 | 28 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Three-dimensional echocardiographic assessments of exercise-induced changes in left ventricular shape and dyssynchrony in patients with dynamic functional mitral regurgitation. <i>European Journal of Echocardiography</i> , 2009, 10, 961-967. | 2.3 | 33 |