

Ewa Āojkowska

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1744252/publications.pdf>

Version: 2024-02-01

117
papers

4,116
citations

117625

34
h-index

144013

57
g-index

128
all docs

128
docs citations

128
times ranked

2854
citing authors

#	ARTICLE	IF	CITATIONS
1	Heterogenicity within the LPS Structure in Relation to the Chosen Genomic and Physiological Features of the Plant Pathogen <i>Pectobacterium parmentieri</i> . <i>International Journal of Molecular Sciences</i> , 2022, 23, 2077.	4.1	7
2	Application of pulse-modulated radio-frequency atmospheric pressure glow discharge for degradation of doxycycline from a flowing liquid solution. <i>Scientific Reports</i> , 2022, 12, 7354.	3.3	3
3	The occurrence of bacteria from different species of <i>Pectobacteriaceae</i> on seed potato plantations in Poland. <i>European Journal of Plant Pathology</i> , 2021, 159, 309-325.	1.7	17
4	Molecular Interactions of <i>Pectobacterium</i> and <i>Dickeya</i> with Plants. , 2021, , 85-147.		12
5	Comparative Genomics, from the Annotated Genome to Valuable Biological Information: A Case Study. <i>Methods in Molecular Biology</i> , 2021, 2242, 91-112.	0.9	0
6	Diseases Caused by <i>Pectobacterium</i> and <i>Dickeya</i> Species Around the World. , 2021, , 215-261.		25
7	Identification and Quantification of Coumarins by UHPLC-MS in <i>Arabidopsis thaliana</i> Natural Populations. <i>Molecules</i> , 2021, 26, 1804.	3.8	9
8	The First Polish Isolate of a Novel Species <i>Pectobacterium aquaticum</i> Originates from a Pomeranian Lake. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 5041.	2.6	6
9	Genome-Wide Analyses of the Temperature-Responsive Genetic Loci of the Pectinolytic Plant Pathogenic <i>Pectobacterium atrosepticum</i> . <i>International Journal of Molecular Sciences</i> , 2021, 22, 4839.	4.1	4
10	Interplay between Coumarin Accumulation, Iron Deficiency and Plant Resistance to <i>Dickeya</i> spp.. <i>International Journal of Molecular Sciences</i> , 2021, 22, 6449.	4.1	10
11	Implementation of a Non-Thermal Atmospheric Pressure Plasma for Eradication of Plant Pathogens from a Surface of Economically Important Seeds. <i>International Journal of Molecular Sciences</i> , 2021, 22, 9256.	4.1	9
12	PacBio-Based Protocol for Bacterial Genome Assembly. <i>Methods in Molecular Biology</i> , 2021, 2242, 3-14.	0.9	1
13	Isolation, Detection and Characterization of <i>Pectobacterium</i> and <i>Dickeya</i> Species. , 2021, , 149-173.		5
14	Effects of stressful physico-chemical factors on the fitness of the plant pathogenic bacterium <i>Dickeya solani</i> . <i>European Journal of Plant Pathology</i> , 2020, 156, 519-535.	1.7	5
15	Comparative genomics and pangenome-oriented studies reveal high homogeneity of the agronomically relevant enterobacterial plant pathogen <i>Dickeya solani</i> . <i>BMC Genomics</i> , 2020, 21, 449.	2.8	16
16	The metabolic shift in highly and weakly virulent <i>Dickeya solani</i> strains is more affected by temperature than by mutations in genes encoding global virulence regulators. <i>FEMS Microbiology Ecology</i> , 2020, 96, .	2.7	2
17	The structure of the O-polysaccharide isolated from pectinolytic gram-negative bacterium <i>Dickeya aquatica</i> IFB0154 is different from the O-polysaccharides of other <i>Dickeya</i> species. <i>Carbohydrate Research</i> , 2020, 497, 108135.	2.3	7
18	3-Chloroplumbagin Induces Cell Death in Breast Cancer Cells Through MAPK-Mediated Mcl-1 Inhibition. <i>Frontiers in Pharmacology</i> , 2019, 10, 784.	3.5	14

#	ARTICLE	IF	CITATIONS
19	Metabolic Modeling of <i>Pectobacterium parmentieri</i> SCC3193 Provides Insights into Metabolic Pathways of Plant Pathogenic Bacteria. <i>Microorganisms</i> , 2019, 7, 101.	3.6	10
20	Plumbagin Increases Paclitaxel-Induced Cell Death and Overcomes Paclitaxel Resistance in Breast Cancer Cells through ERK-Mediated Apoptosis Induction. <i>Journal of Natural Products</i> , 2019, 82, 878-885.	3.0	27
21	Rapid eradication of bacterial phytopathogens by atmospheric pressure glow discharge generated in contact with a flowing liquid cathode. <i>Biotechnology and Bioengineering</i> , 2018, 115, 1581-1593.	3.3	15
22	Genotypic and phenotypic variability of <i>Pectobacterium</i> strains causing blackleg and soft rot on potato in Turkey. <i>European Journal of Plant Pathology</i> , 2018, 152, 143-155.	1.7	19
23	Scopoletin 8-hydroxylase: a novel enzyme involved in coumarin biosynthesis and iron-deficiency responses in <i>Arabidopsis</i> . <i>Journal of Experimental Botany</i> , 2018, 69, 1735-1748.	4.8	86
24	Interplay of classic Exp and specific Vfm quorum sensing systems on the phenotypic features of <i>Dickeya solani</i> strains exhibiting different virulence levels. <i>Molecular Plant Pathology</i> , 2018, 19, 1238-1251.	4.2	30
25	Population Structure and Biodiversity of <i>Pectobacterium parmentieri</i> Isolated from Potato Fields in Temperate Climate. <i>Plant Disease</i> , 2018, 102, 154-164.	1.4	37
26	Antibacterial Activity of Fructose-Stabilized Silver Nanoparticles Produced by Direct Current Atmospheric Pressure Glow Discharge towards Quarantine Pests. <i>Nanomaterials</i> , 2018, 8, 751.	4.1	29
27	Comparison of Highly and Weakly Virulent <i>Dickeya solani</i> Strains, With a View on the Pangenome and Panregulon of This Species. <i>Frontiers in Microbiology</i> , 2018, 9, 1940.	3.5	50
28	High genomic variability in the plant pathogenic bacterium <i>Pectobacterium parmentieri</i> deciphered from de novo assembled complete genomes. <i>BMC Genomics</i> , 2018, 19, 751.	2.8	28
29	Application of Silver Nanostructures Synthesized by Cold Atmospheric Pressure Plasma for Inactivation of Bacterial Phytopathogens from the Genera <i>Dickeya</i> and <i>Pectobacterium</i> . <i>Materials</i> , 2018, 11, 331.	2.9	21
30	Characterization of <i>Dickeya</i> and <i>Pectobacterium</i> strains obtained from diseased potato plants in different climatic conditions of Norway and Poland. <i>European Journal of Plant Pathology</i> , 2017, 148, 839-851.	1.7	42
31	The uniform structure of O-polysaccharides isolated from <i>Dickeya solani</i> strains of different origin. <i>Carbohydrate Research</i> , 2017, 445, 40-43.	2.3	14
32	Plumbagin sensitizes breast cancer cells to tamoxifen-induced cell death through GRP78 inhibition and Bik upregulation. <i>Scientific Reports</i> , 2017, 7, 43781.	3.3	28
33	Molecular methods as tools to control plant diseases caused by <i>Dickeya</i> and <i>Pectobacterium</i> spp: A minireview. <i>New Biotechnology</i> , 2017, 39, 181-189.	4.4	45
34	The effect of temperature on the phenotypic features and the maceration ability of <i>Dickeya solani</i> strains isolated in Finland, Israel and Poland. <i>European Journal of Plant Pathology</i> , 2017, 147, 803-817.	1.7	33
35	Temperature-responsive genetic loci in pectinolytic plant pathogenic <i>Dickeya solani</i> . <i>Plant Pathology</i> , 2017, 66, 584-594.	2.4	37
36	A review on <i>Dickeya solani</i> , a new pathogenic bacterium causing loss in potato yield in Europe. <i>Biotechnologia</i> , 2016, 2, 109-127.	0.9	8

#	ARTICLE	IF	CITATIONS
37	Biodiversity of <i>Dickeya</i> spp. Isolated from Potato Plants and Water Sources in Temperate Climate. <i>Plant Disease</i> , 2016, 100, 408-417.	1.4	64
38	The structure of O-polysaccharides isolated from plant pathogenic bacteria <i>Pectobacterium wasabiae</i> IFB5408 and IFB5427. <i>Carbohydrate Research</i> , 2016, 426, 46-49.	2.3	18
39	Application of zinc chloride precipitation method for rapid isolation and concentration of infectious <i>Pectobacterium</i> spp. and <i>Dickeya</i> spp. lytic bacteriophages from surface water and plant and soil extracts. <i>Folia Microbiologica</i> , 2016, 61, 29-33.	2.3	11
40	Ramentaceone, a Naphthoquinone Derived from <i>Drosera</i> sp., Induces Apoptosis by Suppressing PI3K/Akt Signaling in Breast Cancer Cells. <i>PLoS ONE</i> , 2016, 11, e0147718.	2.5	20
41	Antibacterial activity of caffeine against plant pathogenic bacteria. <i>Acta Biochimica Polonica</i> , 2015, 62, 605-612.	0.5	37
42	Draft Genome Sequence of a Highly Virulent Strain of the Plant Pathogen <i>Dickeya solani</i> , IFB0099. <i>Genome Announcements</i> , 2015, 3, .	0.8	22
43	The complete genome, structural proteome, comparative genomics and phylogenetic analysis of a broad host lytic bacteriophage ĨD3 infecting pectinolytic <i>Dickeya</i> spp.. <i>Standards in Genomic Sciences</i> , 2015, 10, 68.	1.5	16
44	Salicylic acid can reduce infection symptoms caused by <i>Dickeya solani</i> in tissue culture grown potato (<i>Solanum tuberosum</i> L.) plants. <i>European Journal of Plant Pathology</i> , 2015, 141, 545-558.	1.7	48
45	Detection, identification and differentiation of <i>Pectobacterium</i> and <i>Dickeya</i> species causing potato blackleg and tuber soft rot: a review. <i>Annals of Applied Biology</i> , 2015, 166, 18-38.	2.5	166
46	Regulators Involved in <i>Dickeya solani</i> Virulence, Genetic Conservation and Functional Variability. <i>Molecular Plant-Microbe Interactions</i> , 2015, 2015, 5-16.	2.6	1
47	First Report of <i>Pectobacterium carotovorum</i> subsp. <i>brasiliense</i> Causing Soft Rot on Potato and Other Vegetables in Poland. <i>Plant Disease</i> , 2015, 99, 1271-1271.	1.4	35
48	Genomic, Proteomic and Morphological Characterization of Two Novel Broad Host Lytic Bacteriophages ĨPD10.3 and ĨPD23.1 Infecting Pectinolytic <i>Pectobacterium</i> spp. and <i>Dickeya</i> spp.. <i>PLoS ONE</i> , 2015, 10, e0119812.	2.5	90
49	Regulators Involved in <i>Dickeya solani</i> Virulence, Genetic Conservation and Functional Variability. <i>Molecular Plant-Microbe Interactions</i> , 2015, 2015, 57-68.	2.6	1
50	Complete genome sequence of a broad-host-range lytic <i>Dickeya</i> spp. bacteriophage ĨD5. <i>Archives of Virology</i> , 2014, 159, 3153-3155.	2.1	45
51	Identification of QTLs affecting scopolin and scopoletin biosynthesis in <i>Arabidopsis thaliana</i> . <i>BMC Plant Biology</i> , 2014, 14, 280.	3.6	33
52	Simultaneous detection of major blackleg and soft rot bacterial pathogens in potato by multiplex polymerase chain reaction. <i>Annals of Applied Biology</i> , 2014, 165, 474-487.	2.5	56
53	Isolation and identification of cytotoxic compounds from the rhizomes of <i>Paris quadrifolia</i> L.. <i>Pharmacognosy Magazine</i> , 2014, 10, 324.	0.6	10
54	Isolation and characterization of novel soilborne lytic bacteriophages infecting <i>Dickeya</i> spp. biovar 3 (Ĥ <i>D. solani</i> ™). <i>Plant Pathology</i> , 2014, 63, 758-772.	2.4	85

#	ARTICLE	IF	CITATIONS
55	<i>Dickeya solani</i> sp. nov., a pectinolytic plant-pathogenic bacterium isolated from potato (<i>Solanum</i>) Tj ETQq1 1 0.784314 rgBT/Overload	1.7	228
56	Characterization of <i>Pectobacterium carotovorum</i> subsp. <i>odoriferum</i> causing soft rot of stored vegetables. <i>European Journal of Plant Pathology</i> , 2014, 139, 457-469.	1.7	40
57	Effect of <i>Dionaea muscipula</i> extract and plumbagin on maceration of potato tissue by <i>Pectobacterium atrosepticum</i> . <i>Annals of Applied Biology</i> , 2014, 164, 404-414.	2.5	8
58	Regulators Involved in <i>Dickeya solani</i> Virulence, Genetic Conservation, and Functional Variability. <i>Molecular Plant-Microbe Interactions</i> , 2014, 27, 700-711.	2.6	49
59	Occurrence of <i>Pectobacterium wasabiae</i> in potato field samples. <i>European Journal of Plant Pathology</i> , 2013, 137, 149-158.	1.7	43
60	Wacław Szybalski: Lwów, Gdańsk, Madison – Life of scientist and philanthropist. <i>Gene</i> , 2013, 525, 155-157.	2.2	1
61	Multiplex detection and identification of bacterial pathogens causing potato blackleg and soft rot in Europe, using padlock probes. <i>Annals of Applied Biology</i> , 2013, 163, 378-393.	2.5	17
62	A new clade of <i>Dickeya</i> spp. plays a major role in potato blackleg outbreaks in North Finland. <i>Annals of Applied Biology</i> , 2013, 162, 231-241.	2.5	81
63	Plumbagin Induces Apoptosis in Her2-Overexpressing Breast Cancer Cells through the Mitochondrial-Mediated Pathway. <i>Journal of Natural Products</i> , 2012, 75, 747-751.	3.0	51
64	Effect of l-phenylalanine on PAL activity and production of naphthoquinone pigments in suspension cultures of <i>Arnebia euchroma</i> (Royle) Johnst. <i>In Vitro Cellular and Developmental Biology - Plant</i> , 2012, 48, 555-564.	2.1	63
65	Induction of Apoptosis in HL-60 Cells through the ROS-Mediated Mitochondrial Pathway by Ramentaceone from <i>Drosera aliciae</i> . <i>Journal of Natural Products</i> , 2012, 75, 9-14.	3.0	56
66	Enhanced production of antitumour naphthoquinones in transgenic hairy root lines of <i>Lithospermum canescens</i> . <i>Plant Cell, Tissue and Organ Culture</i> , 2012, 108, 213-219.	2.3	28
67	Triterpenoid \pm -amyrin stimulates proliferation of human keratinocytes but does not protect them against UVB damage.. <i>Acta Biochimica Polonica</i> , 2012, 59, .	0.5	15
68	Inactivation of AHLs by <i>Ochrobactrum</i> sp. A44 depends on the activity of a novel class of AHL acylase. <i>Environmental Microbiology Reports</i> , 2011, 3, 59-68.	2.4	65
69	<i>Dickeya</i> species: an emerging problem for potato production in Europe. <i>Plant Pathology</i> , 2011, 60, 385-399.	2.4	383
70	Chemical structure of the O-polysaccharide isolated from <i>Pectobacterium atrosepticum</i> SCRI 1039. <i>Carbohydrate Research</i> , 2011, 346, 2978-2981.	2.3	14
71	Polymorphism analysis of housekeeping genes for identification and differentiation of <i>Clavibacter michiganensis</i> subspecies. <i>European Journal of Plant Pathology</i> , 2011, 131, 341-354.	1.7	22
72	In vitro cultures of <i>Drosera aliciae</i> as a source of a cytotoxic naphthoquinone: ramentaceone. <i>Biotechnology Letters</i> , 2011, 33, 2309-2316.	2.2	18

#	ARTICLE	IF	CITATIONS
73	Cytotoxic Activity of Paris quadrifolia Extract and Isolated Saponin Fractions Against Human Tumor Cell Lines. Acta Biologica Cracoviensia Series Botanica, 2011, 53, .	0.5	1
74	Detection and identification of potentially toxic cyanobacteria in Polish water bodies.. Acta Biochimica Polonica, 2011, 58, .	0.5	21
75	Asymbiotic germination, seedling development and plantlet propagation of Encyclia aff. oncioides - an endangered orchid. Acta Societatis Botanicorum Poloniae, 2011, 74, 193-198.	0.8	5
76	Teratomas of Drosera capensis var. alba as a source of naphthoquinone: ramentaceone. Plant Cell, Tissue and Organ Culture, 2010, 103, 285-292.	2.3	32
77	Chromatographic analysis of simple phenols in some species from the genus <i>Salix</i>. Phytochemical Analysis, 2010, 21, 463-469.	2.4	27
78	Antibacterial and antioxidant activity of the secondary metabolites from <i>in vitro</i> cultures of the Alice sundew (<i>Drosera aliciae</i>). Biotechnology and Applied Biochemistry, 2009, 53, 175-184.	3.1	34
79	Genes responsible for coronatine synthesis in Pseudomonas syringae present in the genome of soft rot bacteria. European Journal of Plant Pathology, 2009, 124, 353-361.	1.7	21
80	Genetic transformation of Ruta graveolens L. by Agrobacterium rhizogenes: hairy root cultures a promising approach for production of coumarins and furanocoumarins. Plant Cell, Tissue and Organ Culture, 2009, 97, 59-69.	2.3	37
81	First report of bacterial soft rot on potato caused by <i>Dickeya</i> sp. (syn. <i>Erwinia</i> Tj ETQq1 1 0.784314 rgBT / Overlock 10 Tf 5	2.4	56
82	Application of RFLP analysis of recA, gyrA and rpoS gene fragments for rapid differentiation of Erwinia amylovora from Erwinia strains isolated in Korea and Japan. European Journal of Plant Pathology, 2008, 121, 161-172.	1.7	34
83	Effective biotic elicitation of Ruta graveolens L. shoot cultures by lysates from Pectobacterium atrosepticum and Bacillus sp.. Biotechnology Letters, 2008, 30, 541-545.	2.2	13
84	Stimulation of antibacterial naphthoquinones and flavonoids accumulation in carnivorous plants grown in vitro by addition of elicitors. Enzyme and Microbial Technology, 2008, 42, 216-221.	3.2	60
85	Simplex Optimized LC Analysis of Plant Coumarins and Furanocoumarins. Chromatographia, 2008, 67, 653-657.	1.3	3
86	Application of chitin and chitosan as elicitors of coumarins and furoquinolone alkaloids in <i>Ruta graveolens</i> L. (common rue). Biotechnology and Applied Biochemistry, 2008, 51, 91-96.	3.1	72
87	Identification of Ruta graveolens L. Metabolites Accumulated in the Presence of Abiotic Elicitors. Biotechnology Progress, 2008, 24, 128-133.	2.6	19
88	Rapid detection of mutagens accumulated in plant tissues using a novel Vibrio harveyi mutagenicity assay. Ecotoxicology and Environmental Safety, 2008, 70, 231-235.	6.0	4
89	Induction of apoptosis by plumbagin through reactive oxygen species-mediated inhibition of topoisomerase II. Toxicology and Applied Pharmacology, 2007, 223, 267-276.	2.8	83
90	Identification of a DNA restriction-modification system in Pectobacterium carotovorum strains isolated from Poland. Journal of Applied Microbiology, 2006, 100, 343-351.	3.1	0

#	ARTICLE	IF	CITATIONS
91	Induction of secondary metabolite production in transformed callus of <i>Ammi majus</i> L. grown after electromagnetic treatment of the culture medium. <i>Enzyme and Microbial Technology</i> , 2006, 39, 1386-1391.	3.2	23
92	Antibacterial Activity of Synthetic Peptides Against Plant Pathogenic <i>Pectobacterium</i> Species. <i>Journal of Phytopathology</i> , 2005, 153, 313-317.	1.0	19
93	Secondary metabolites in vitro cultured plants of the genus <i>Drosera</i> . <i>Phytochemical Analysis</i> , 2005, 16, 143-149.	2.4	43
94	Application of rapd in the determination of genetic fidelity in micropropagated <i>Drosera</i> plantlets. In <i>Vitro Cellular and Developmental Biology - Plant</i> , 2004, 40, 592-595.	2.1	36
95	Direct regeneration of <i>Drosera</i> from leaf explants and shoot tips. <i>Plant Cell, Tissue and Organ Culture</i> , 2003, 75, 175-178.	2.3	27
96	HPLC-DAD in identification and quantification of selected coumarins in crude extracts from plant cultures of <i>Ammi majus</i> and <i>Ruta graveolens</i> . <i>Journal of Separation Science</i> , 2003, 26, 1287-1291.	2.5	27
97	Elicitation of secondary metabolites in in vitro cultures of <i>Ammi majus</i> L.. <i>Enzyme and Microbial Technology</i> , 2003, 33, 565-568.	3.2	78
98	Establishment of a co-culture of <i>Ammi majus</i> L. and <i>Ruta graveolens</i> L. for the synthesis of furanocoumarins. <i>Plant Science</i> , 2003, 165, 1315-1319.	3.6	34
99	Genotyping of bacteria belonging to the former <i>Erwinia</i> genus by PCR-RFLP analysis of a <i>recA</i> gene fragment. <i>Microbiology (United Kingdom)</i> , 2002, 148, 583-595.	1.8	123
100	Genotypic characterisation of the <i>Erwinia</i> genus by PCR-RFLP analysis of <i>rpoS</i> gene. <i>Plant Protection Science</i> , 2002, 38, 288-290.	1.4	14
101	Establishment of hairy root cultures of <i>Ammi majus</i> . <i>Plant Science</i> , 2001, 160, 259-264.	3.6	58
102	IDENTIFICATION OF SECONDARY METABOLITES IN IN VITRO CULTURE OF AMMI MAJUS TREATED WITH ELICITORS. <i>Acta Horticulturae</i> , 2001, , 255-258.	0.2	6
103	Genetic diversity of <i>Erwinia carotovora</i> strains isolated from infected plants grown in Poland. <i>EPPO Bulletin</i> , 2000, 30, 403-407.	0.8	30
104	Expression of <i>Erwinia chrysanthemi</i> Pectinase Genes <i>pell</i> , <i>pelL</i> , and <i>pelZ</i> During Infection of Potato Tubers. <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 845-851.	2.6	20
105	Production of <i>Erwinia chrysanthemi</i> pectinases in potato tubers showing high or low level of resistance to soft-rot. <i>European Journal of Plant Pathology</i> , 1996, 102, 511-517.	1.7	11
106	Characterization of the <i>pelL</i> gene encoding a novel pectate lyase of <i>Erwinia chrysanthemi</i> 3937. <i>Molecular Microbiology</i> , 1995, 16, 1183-1195.	2.5	103
107	Comparison of the effectiveness of different methods of screening for bacterial soft rot resistance of potato tubers. <i>American Potato Journal</i> , 1994, 71, 99-113.	0.3	23
108	Resistance to <i>Erwinia</i> spp. in diploid potato with a high starch content. <i>Potato Research</i> , 1993, 36, 177-182.	2.7	26

#	ARTICLE	IF	CITATIONS
109	Use of GUS Fusion to Study the Expression of <i>Erwinia chrysanthemi</i> Pectinase Genes During Infection of Potato Tubers. <i>Molecular Plant-Microbe Interactions</i> , 1993, 6, 488.	2.6	30
110	The Role of Polyphenol Oxidase and Peroxidase in Potato Tuber Resistance to Soft Rot Caused by <i>Erwinia carotovora</i> . <i>Journal of Phytopathology</i> , 1992, 136, 319-328.	1.0	17
111	Changes of the lipid catabolism in potato tubers from cultivars differing in susceptibility to autolysis during the storage. <i>Potato Research</i> , 1989, 32, 463-470.	2.7	19
112	Screening of seedlings of wild <i>Solanum</i> species for resistance to bacterial stem rot caused by soft rot <i>Erwinias</i> . <i>American Potato Journal</i> , 1989, 66, 379-390.	0.3	19
113	Lipid composition and post-wounding degradation in potato slices from cultivars differing in susceptibility to autolysis. <i>Potato Research</i> , 1988, 31, 541-549.	2.7	6
114	Post-wounding changes in the oxygen consumption by slices from tubers of several potato cultivars. <i>Potato Research</i> , 1988, 31, 550-556.	2.7	1
115	Fertile Interspecific Somatic Hybrids of <i>Solanum</i> : A Novel Source of Resistance to <i>Erwinia</i> Soft Rot. <i>Phytopathology</i> , 1988, 78, 1216.	2.2	133
116	The effect of wound healing and of certain chemicals on electrolyte release from discs of potato by enzymes of <i>Erwinia carotovora</i> . <i>Potato Research</i> , 1984, 27, 131-143.	2.7	4
117	Regulators Involved in <i>Dickeya solani</i> Virulence, Genetic Conservation and Functional Variability. <i>Molecular Plant-Microbe Interactions</i> , 0, , MPMI-99-99-0004.	2.6	0