Young Suk Park

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1806086/publications.pdf

Version: 2024-02-01

172457 123424 4,636 181 29 61 citations h-index g-index papers 183 183 183 8274 docs citations times ranked citing authors all docs

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Comprehensive molecular characterization of clinical responses to PD-1 inhibition in metastatic gastric cancer. Nature Medicine, 2018, 24, 1449-1458. | 30.7 | 1,071 |
| 2 | Phase III Trial to Compare Adjuvant Chemotherapy With Capecitabine and Cisplatin Versus Concurrent Chemoradiotherapy in Gastric Cancer: Final Report of the Adjuvant Chemoradiotherapy in Stomach Tumors Trial, Including Survival and Subset Analyses. Journal of Clinical Oncology, 2015, 33, 3130-3136. | 1.6 | 370 |
| 3 | Oxaliplatin, fluorouracil, and leucovorin versus fluorouracil and leucovorin as adjuvant chemotherapy for locally advanced rectal cancer after preoperative chemoradiotherapy (ADORE): an open-label, multicentre, phase 2, randomised controlled trial. Lancet Oncology, The, 2014, 15, 1245-1253. | 10.7 | 336 |
| 4 | Tumor Genomic Profiling Guides Patients with Metastatic Gastric Cancer to Targeted Treatment: The VIKTORY Umbrella Trial. Cancer Discovery, 2019, 9, 1388-1405. | 9.4 | 155 |
| | Modified XELIRI (capecitabine plus irinotecan) versus FOLFIRI (leucovorin, fluorouracil, and) Tj ETQq1 1 0.784314 | | |
| 5 | colorectal cancer (AXEPT): a multicentre, open-label, randomised, non-inferiority, phase 3 trial. Lancet Oncology. The. 2018. 19. 660-671. | 10.7 | 107 |
| 6 | Oxaliplatin-Based Adjuvant Chemotherapy for Rectal Cancer After Preoperative Chemoradiotherapy (ADORE): Long-Term Results of a Randomized Controlled Trial. Journal of Clinical Oncology, 2019, 37, 3111-3123. | 1.6 | 100 |
| 7 | Prospective blinded study of somatic mutation detection in cell-free DNA utilizing a targeted 54-gene next generation sequencing panel in metastatic solid tumor patients. Oncotarget, 2015, 6, 40360-40369. | 1.8 | 85 |
| 8 | Simvastatin plus capecitabine–cisplatin versus placebo plus capecitabine–cisplatin in patients with previously untreated advanced gastric cancer: A double-blind randomised phase 3 study. European Journal of Cancer, 2014, 50, 2822-2830. | 2.8 | 79 |
| 9 | c-MET Overexpression in Colorectal Cancer: A Poor Prognostic Factor for Survival. Clinical Colorectal Cancer, 2018, 17, 165-169. | 2.3 | 71 |
| 10 | Correlating programmed death ligand 1 (PD-L1) expression, mismatch repair deficiency, and outcomes across tumor types: implications for immunotherapy. Oncotarget, 2017, 8, 77415-77423. | 1.8 | 68 |
| 11 | A Phase 1 Study of LY2874455, an Oral Selective pan-FGFR Inhibitor, in Patients with Advanced Cancer. Targeted Oncology, 2017, 12, 463-474. | 3.6 | 64 |
| 12 | Gastrointestinal malignancies harbor actionable MET exon 14 deletions. Oncotarget, 2015, 6, 28211-28222. | 1.8 | 57 |
| 13 | Genomic characterization of intrinsic and acquired resistance to cetuximab in colorectal cancer patients. Scientific Reports, 2019, 9, 15365. | 3.3 | 54 |
| 14 | Phase I Study of Ceralasertib (AZD6738), a Novel DNA Damage Repair Agent, in Combination with Weekly Paclitaxel in Refractory Cancer. Clinical Cancer Research, 2021, 27, 4700-4709. | 7.0 | 54 |
| 15 | NTRK1 rearrangement in colorectal cancer patients: evidence for actionable target using patient-derived tumor cell line. Oncotarget, 2015, 6, 39028-39035. | 1.8 | 53 |
| 16 | Patient-derived cell models as preclinical tools for genome-directed targeted therapy. Oncotarget, 2015, 6, 25619-25630. | 1.8 | 48 |
| 17 | MCT4 as a potential therapeutic target for metastatic gastric cancer with peritoneal carcinomatosis. Oncotarget, 2016, 7, 43492-43503. | 1.8 | 45 |
| 18 | Metformin enhances the response to radiotherapy in diabetic patients with rectal cancer. Journal of Cancer Research and Clinical Oncology, 2016, 142, 1377-1385. | 2.5 | 40 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | A multi-center, open-label, randomized phase III trial of first-line chemotherapy with capecitabine monotherapy versus capecitabine plus oxaliplatin in elderly patients with advanced gastric cancer. Journal of Geriatric Oncology, 2017, 8, 170-175. | 1.0 | 39 |
| 20 | Anastomotic Leak Does Not Impact Oncologic Outcomes After Preoperative Chemoradiotherapy and Resection for Rectal Cancer. Annals of Surgery, 2019, 269, 678-685. | 4.2 | 37 |
| 21 | MCT4 Expression Is a Potential Therapeutic Target in Colorectal Cancer with Peritoneal Carcinomatosis. Molecular Cancer Therapeutics, 2018, 17, 838-848. | 4.1 | 36 |
| 22 | Identification of the BRAF V600E mutation in gastroenteropancreatic neuroendocrine tumors. Oncotarget, 2016, 7, 4024-4035. | 1.8 | 36 |
| 23 | The impact of KRAS mutations on prognosis in surgically resected colorectal cancer patients with liver and lung metastases: a retrospective analysis. BMC Cancer, 2016, 16, 120. | 2.6 | 35 |
| 24 | Pazopanib, a Novel Multitargeted Kinase Inhibitor, Shows Potent <i>In Vitro</i> Antitumor Activity in Gastric Cancer Cell Lines with <i>FGFR2</i> Amplification. Molecular Cancer Therapeutics, 2014, 13, 2527-2536. | 4.1 | 34 |
| 25 | The Influence of Metastatic Lymph Node Ratio on the Treatment Outcomes in the Adjuvant Chemoradiotherapy in Stomach Tumors (ARTIST) Trial: A Phase III Trial. Journal of Gastric Cancer, 2016, 16, 105. | 2.5 | 34 |
| 26 | Large-scale clinical validation of biomarkers for pancreatic cancer using a mass spectrometry-based proteomics approach. Oncotarget, 2017, 8, 42761-42771. | 1.8 | 34 |
| 27 | Tissue recommendations for precision cancer therapy using next generation sequencing: a comprehensive single cancer center's experiences. Oncotarget, 2017, 8, 42478-42486. | 1.8 | 32 |
| 28 | Detection of novel and potentially actionable anaplastic lymphoma kinase (ALK) rearrangement in colorectal adenocarcinoma by immunohistochemistry screening. Oncotarget, 2015, 6, 24320-24332. | 1.8 | 32 |
| 29 | Effects of adjuvant radiotherapy on completely resected gastric cancer: A radiation oncologist's view of the ARTIST randomized phase III trial. Radiotherapy and Oncology, 2015, 117, 171-177. | 0.6 | 31 |
| 30 | Prospective Feasibility Study for Using Cell-Free Circulating Tumor DNA–Guided Therapy in Refractory Metastatic Solid Cancers: An Interim Analysis. JCO Precision Oncology, 2017, 1, 1-15. | 3.0 | 31 |
| 31 | Claudin 18.2 expression in various tumor types and its role as a potential target in advanced gastric cancer. Translational Cancer Research, 2020, 9, 3367-3374. | 1.0 | 26 |
| 32 | Circulating Tumor Cells are Predictive of Poor Response to Chemotherapy in Metastatic gastric cancer. International Journal of Biological Markers, 2015, 30, 382-386. | 1.8 | 25 |
| 33 | Triptolide as a novel agent in pancreatic cancer: the validation using patient derived pancreatic tumor cell line. BMC Cancer, 2018, 18, 1103. | 2.6 | 25 |
| 34 | Tumor regression grade as a clinically useful outcome predictor in patients with rectal cancer after preoperative chemoradiotherapy. Surgery, 2019, 165, 579-585. | 1.9 | 25 |
| 35 | CD133-positive tumor cell content is a predictor of early recurrence in colorectal cancer. Journal of Gastrointestinal Oncology, 2014, 5, 447-56. | 1.4 | 25 |
| 36 | Transcriptome analysis of CD133-positive stem cells and prognostic value of survivin in colorectal cancer. Cancer Genomics and Proteomics, 2014, 11, 259-66. | 2.0 | 25 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Anti-tumor efficacy of fulvestrant in estrogen receptor positive gastric cancer. Scientific Reports, 2014, 4, 7592. | 3.3 | 24 |
| 38 | A Randomized Phase 2 Study of Neoadjuvant Chemoradiaton Therapy With 5-Fluorouracil/Leucovorin or Irinotecan/S-1 in Patients With Locally Advanced Rectal Cancer. International Journal of Radiation Oncology Biology Physics, 2015, 93, 1015-1022. | 0.8 | 24 |
| 39 | A phase II study of preoperative mFOLFOX6 with short-course radiotherapy in patients with locally advanced rectal cancer and liver-only metastasis. Radiotherapy and Oncology, 2016, 118, 369-374. | 0.6 | 24 |
| 40 | Prospective phase II trial of everolimus in PIK3CA amplification/mutation and/or PTEN loss patients with advanced solid tumors refractory to standard therapy. BMC Cancer, 2017, 17, 211. | 2.6 | 24 |
| 41 | The NEXT-1 (Next generation pErsonalized tX with mulTi-omics and preclinical model) trial: prospective molecular screening trial of metastatic solid cancer patients, a feasibility analysis. Oncotarget, 2015, 6, 33358-33368. | 1.8 | 24 |
| 42 | MerTK is a novel therapeutic target in gastric cancer. Oncotarget, 2017, 8, 96656-96667. | 1.8 | 23 |
| 43 | Changes in Metabolic Syndrome Status are Associated With Altered Risk of Pancreatic Cancer: A Nationwide Cohort Study. Gastroenterology, 2022, 162, 509-520.e7. | 1.3 | 23 |
| 44 | Genomic Alterations in Biliary Tract Cancer Using Targeted Sequencing. Translational Oncology, 2016, 9, 173-178. | 3.7 | 22 |
| 45 | Phase I trial and pharmacokinetic study of tanibirumab, a fully human monoclonal antibody to vascular endothelial growth factor receptor 2, in patients with refractory solid tumors. Investigational New Drugs, 2017, 35, 782-790. | 2.6 | 22 |
| 46 | Two-week course of preoperative chemoradiotherapy followed by delayed surgery for rectal cancer: A phase II multi-institutional clinical trial (KROG 11-02). Radiotherapy and Oncology, 2014, 110, 150-154. | 0.6 | 21 |
| 47 | Direct analysis of aberrant glycosylation on haptoglobin in patients with gastric cancer. Oncotarget, 2017, 8, 11094-11104. | 1.8 | 21 |
| 48 | Activated cMET and IGF1R-Driven PI3K Signaling Predicts Poor Survival in Colorectal Cancers Independent of KRAS Mutational Status. PLoS ONE, 2014, 9, e103551. | 2.5 | 21 |
| 49 | Changes in the Mean Corpuscular Volume after Capecitabine Treatment Are Associated with Clinical Response and Survival in Patients with Advanced Gastric Cancer. Cancer Research and Treatment, 1970, 47, 72-77. | 3.0 | 20 |
| 50 | Disappearing or residual tiny (≧Âmm) colorectal liver metastases after chemotherapy on gadoxetic acid-enhanced liver MRI and diffusion-weighted imaging: Is local treatment required?. European Radiology, 2017, 27, 3088-3096. | 4.5 | 20 |
| 51 | The Clinical Impact of c-MET Over-Expression in Advanced Biliary Tract Cancer (BTC). Journal of Cancer, 2017, 8, 1395-1399. | 2.5 | 20 |
| 52 | Clinical sequencing to assess tumor mutational burden as a useful biomarker to immunotherapy in various solid tumors. Therapeutic Advances in Medical Oncology, 2021, 13, 175883592199299. | 3.2 | 20 |
| 53 | The implication of FLT3 amplification for FLT targeted therapeutics in solid tumors. Oncotarget, 2017, 8, 3237-3245. | 1.8 | 20 |
| 54 | The impact of primary tumor location in patients with metastatic colorectal cancer: a Korean Cancer Study Group CO12-04 study. Korean Journal of Internal Medicine, 2019, 34, 165-177. | 1.7 | 20 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | A Retrospective Analysis for Patients with HER2-Positive Gastric Cancer Who Were Treated with Trastuzumab-Based Chemotherapy: In the Perspectives of Ethnicity and Histology. Cancer Research and Treatment, 2016, 48, 553-560. | 3.0 | 19 |
| 56 | The Impact of Microsatellite Instability Status and Sidedness of the Primary Tumor on the Effect of Cetuximab-Containing Chemotherapy in Patients with Metastatic Colorectal Cancer. Journal of Cancer, 2017, 8, 2809-2815. | 2.5 | 18 |
| 57 | Efficacy and safety of vactosertib and pembrolizumab combination in patients with previously treated microsatellite stable metastatic colorectal cancer Journal of Clinical Oncology, 2021, 39, 3573-3573. | 1.6 | 18 |
| 58 | Gemcitabine and Docetaxel Combination for Advanced Soft Tissue Sarcoma: A Nationwide Retrospective Study. Cancer Research and Treatment, 2018, 50, 175-182. | 3.0 | 18 |
| 59 | Tumour shrinkage at 6Âweeks predicts favorable clinical outcomes in a phase III study of gemcitabine and oxaliplatin with or without erlotinib for advanced biliary tract cancer. BMC Cancer, 2015, 15, 530. | 2.6 | 17 |
| 60 | Genomic Profiling of Metastatic Gastroenteropancreatic Neuroendocrine Tumor (GEP-NET) Patients in the Personalized-Medicine Era. Journal of Cancer, 2016, 7, 1044-1048. | 2.5 | 17 |
| 61 | Phase I Trial of Anti-MET Monoclonal Antibody in MET-Overexpressed Refractory Cancer. Clinical Colorectal Cancer, 2018, 17, 140-146. | 2.3 | 17 |
| 62 | The prognostic role of tumor associated glycoprotein 72 (TAG-72) in stage II and III colorectal adenocarcinoma. Pathology Research and Practice, 2019, 215, 171-176. | 2.3 | 17 |
| 63 | S-1 plus oxaliplatin versus capecitabine plus oxaliplatin for the first-line treatment of patients with metastatic colorectal cancer: updated results from a phase 3 trial. BMC Cancer, 2014, 14, 883. | 2.6 | 16 |
| 64 | Molecular Subgroup Analysis of Clinical Outcomes in a Phase 3 Study of Gemcitabine and Oxaliplatin with or without Erlotinib in Advanced Biliary Tract Cancer. Translational Oncology, 2015, 8, 40-46. | 3.7 | 16 |
| 65 | The Impact of Cetuximab Plus AKT- or mTOR- Inhibitor in a Patient-Derived Colon Cancer Cell Model with Wild-Type RAS and PIK3CA Mutation. Journal of Cancer, 2017, 8, 2713-2719. | 2.5 | 16 |
| 66 | Association of prediabetes, diabetes, and diabetes duration with biliary tract cancer risk: A nationwide cohort study. Metabolism: Clinical and Experimental, 2021, 123, 154848. | 3.4 | 16 |
| 67 | MerTK inhibition by RXDX-106 in MerTK activated gastric cancer cell lines. Oncotarget, 2017, 8, 105727-105734. | 1.8 | 16 |
| 68 | Pilot study of sirolimus in patients with PIK3CA mutant/amplified refractory solid cancer. Molecular and Clinical Oncology, 2017, 7, 27-31. | 1.0 | 15 |
| 69 | PIK3CA mutation detection in metastatic biliary cancer using cell-free DNA. Oncotarget, 2015, 6, 40026-40035. | 1.8 | 15 |
| 70 | Prospective phase II trial of pazopanib plus CapeOX (capecitabine and oxaliplatin) in previously untreated patients with advanced gastric cancer. Oncotarget, 2016, 7, 24088-24096. | 1.8 | 15 |
| 71 | Clinical Application of Targeted Deep Sequencing in Solid-Cancer Patients and Utility for Biomarker-Selected Clinical Trials. Oncologist, 2017, 22, 1169-1177. | 3.7 | 14 |
| 72 | Proton Pump Inhibitor Use and the Efficacy of Chemotherapy in Metastatic Colorectal Cancer: A Post Hoc Analysis of a Randomized Phase III Trial (AXEPT). Oncologist, 2021, 26, e954-e962. | 3.7 | 14 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | A nCounter CNV Assay to Detect HER2 Amplification: A Correlation Study with Immunohistochemistry and In Situ Hybridization in Advanced Gastric Cancer. Molecular Diagnosis and Therapy, 2016, 20, 375-383. | 3.8 | 13 |
| 74 | <p>Everolimus for the treatment of advanced gastrointestinal or lung nonfunctional neuroendocrine tumors in East Asian patients: a subgroup analysis of the RADIANT-4 study</p> . OncoTargets and Therapy, 2019, Volume 12, 1717-1728. | 2.0 | 13 |
| 75 | Clinical Outcomes of Neoadjuvant Chemotherapy in Colorectal Cancer Patients With Synchronous Resectable Liver Metastasis: A Propensity Score Matching Analysis. Annals of Coloproctology, 2021, 37, 244-252. | 2.0 | 13 |
| 76 | Study protocol of the Asian XELIRI ProjecT (AXEPT): a multinational, randomized, non-inferiority, phase III trial of second-line chemotherapy for metastatic colorectal cancer, comparing the efficacy and safety of XELIRI with or without bevacizumab. Chinese Journal of Cancer, 2016, 35, 102. | 4.9 | 12 |
| 77 | Adjuvant Chemotherapy with or without Concurrent Radiotherapy for Patients with Stage IB Gastric Cancer: a Subgroup Analysis of the Adjuvant Chemoradiotherapy in Stomach Tumors (ARTIST) Phase III Trial. Journal of Gastric Cancer, 2018, 18, 348. | 2.5 | 12 |
| 78 | Necessity of adjuvant concurrent chemo-radiotherapy in D2-resected LN-positive gastric cancer. Radiotherapy and Oncology, 2018, 129, 306-312. | 0.6 | 12 |
| 79 | Molecular characterization of colorectal cancer patients and concomitant patient-derived tumor cell establishment. Oncotarget, 2016, 7, 19610-19619. | 1.8 | 12 |
| 80 | The efficacy of low-dose transdermal fentanyl in opioid-na \tilde{A} -ve cancer patients with moderate-to-severe pain. Korean Journal of Internal Medicine, 2015, 30, 88. | 1.7 | 12 |
| 81 | A Single Arm, Phase II Study of Simvastatin Plus XELOX and Bevacizumab as First-Line Chemotherapy in Metastatic Colorectal Cancer Patients. Cancer Research and Treatment, 2019, 51, 1128-1134. | 3.0 | 12 |
| 82 | Association between alcohol consumption and pancreatic cancer risk differs by glycaemic status: A nationwide cohort study. European Journal of Cancer, 2022, 163, 119-127. | 2.8 | 12 |
| 83 | Value of FGFR2 expression for advanced gastric cancer patients receiving pazopanib plus CapeOX (capecitabine and oxaliplatin). Journal of Cancer Research and Clinical Oncology, 2016, 142, 1231-1237. | 2.5 | 11 |
| 84 | Phase II trial of epidermal growth factor ointment for patients with Erlotinib-related skin effects. Supportive Care in Cancer, 2016, 24, 301-309. | 2.2 | 11 |
| 85 | The Role of Plasma Chromogranin A as Assessment of Treatment Response in Non-functioning Gastroenteropancreatic Neuroendocrine Tumors. Cancer Research and Treatment, 2016, 48, 153-161. | 3.0 | 11 |
| 86 | Tumor Mutational Burden as a Biomarker for Advanced Biliary Tract Cancer. Technology in Cancer Research and Treatment, 2021, 20, 153303382110623. | 1.9 | 11 |
| 87 | Exploratory biomarker analysis for treatment response in KRAS wild type metastatic colorectal cancer patients who received cetuximab plus irinotecan. BMC Cancer, 2015, 15, 747. | 2.6 | 10 |
| 88 | Regorafenib as Salvage Treatment in Korean Patients with Refractory Metastatic Colorectal Cancer. Cancer Research and Treatment, 2015, 47, 790-795. | 3.0 | 10 |
| 89 | Evaluation of quality of life using a tablet PC-based survey in cancer patients treated with radiotherapy: a multi-institutional prospective randomized crossover comparison of paper and tablet PC-based questionnaires (KROG 12–01). Supportive Care in Cancer, 2016, 24, 4399-4406. | 2.2 | 10 |
| 90 | Comprehensive molecular profiling to predict clinical outcomes in pancreatic cancer. Therapeutic Advances in Medical Oncology, 2021, 13, 175883592110384. | 3.2 | 10 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Prospective phase II trial of regional hyperthermia and whole liver irradiation for numerous chemorefractory liver metastases from colorectal cancer. Radiation Oncology Journal, 2016, 34, 34-44. | 1.5 | 10 |
| 92 | Gastroenteropancreatic Neuroendocrine Tumors with Liver Metastases in Korea: A Clinicopathological Analysis of 72 Cases in a Single Institute. Cancer Research and Treatment, 2015, 47, 738-746. | 3.0 | 10 |
| 93 | Importance of the Circumferential Extent of Tumors and Clinical Lymph Node Status as Prognostic Factors after Preoperative Chemoradiotherapy and Surgery in Patients with Rectal Cancer. Tumori, 2010, 96, 568-576. | 1.1 | 9 |
| 94 | Clinical Significance of Mucinous Rectal Adenocarcinoma following Preoperative Chemoradiotherapy and Curative Surgery. Tumori, 2016, 102, 114-121. | 1.1 | 9 |
| 95 | A randomized phase II study of gemcitabine plus Z-360, a CCK2 receptor-selective antagonist, in patients with metastatic pancreatic cancer as compared with gemcitabine plus placebo. Cancer Chemotherapy and Pharmacology, 2017, 80, 307-315. | 2.3 | 9 |
| 96 | First-in-human phase I trial of anti-hepatocyte growth factor antibody (YYB101) in refractory solid tumor patients. Therapeutic Advances in Medical Oncology, 2020, 12, 175883592092679. | 3.2 | 9 |
| 97 | A Randomized Phase II Study of Perioperative Chemotherapy Plus Bevacizumab Versus Postoperative Chemotherapy Plus Bevacizumab in Patients With Upfront Resectable Hepatic Colorectal Metastases. Clinical Colorectal Cancer, 2020, 19, e140-e150. | 2.3 | 9 |
| 98 | Clinical Significance of IGFBP-3 Methylation in Patients with Early Stage Gastric Cancer. Translational Oncology, 2015, 8, 288-294. | 3.7 | 8 |
| 99 | The Correlation Between Serum Chemokines and Clinical Outcome in Patients with Advanced Biliary Tract Cancer. Translational Oncology, 2018, 11, 353-357. | 3.7 | 8 |
| 100 | Combination of Docetaxel Plus Savolitinib in Refractory Cancer Patients: A Report on Phase I Trial. Translational Oncology, 2019, 12, 597-601. | 3.7 | 8 |
| 101 | Clinical and molecular distinctions in patients with refractory colon cancer who benefit from regorafenib treatment. Therapeutic Advances in Medical Oncology, 2020, 12, 175883592096584. | 3.2 | 8 |
| 102 | Impact of <i>UGT1A1</i> genotype on the efficacy and safety of irinotecanâ€based chemotherapy in metastatic colorectal cancer. Cancer Science, 2021, 112, 4669-4678. | 3.9 | 8 |
| 103 | Molecular analysis of the randomized phase II/III study of the anti-IGF-1R antibody dalotuzumab (MK-0646) in combination with cetuximab (Cx) and irinotecan (Ir) in the treatment of chemorefractory KRAS wild-type metastatic colorectal cancer (mCRC) Journal of Clinical Oncology, 2012, 30, 3531-3531. | 1.6 | 8 |
| 104 | Comparison of the 7th and the 8th AJCC Staging System for Non-metastatic D2-Resected Lymph Nodeâ€"Positive Gastric Cancer Treated with Different Adjuvant Protocols. Cancer Research and Treatment, 2019, 51, 876-885. | 3.0 | 8 |
| 105 | Oxaliplatin/5-fluorouracil-based adjuvant chemotherapy as a standard of care for colon cancer in clinical practice: Outcomes of the ACCElox registry. Asia-Pacific Journal of Clinical Oncology, 2015, 11, 334-342. | 1.1 | 7 |
| 106 | The impact of microsatellite instability status and sidedness of the primary tumor on the effect of bevacizumab-containing chemotherapy in patients with metastatic colorectal cancer. Journal of Cancer, 2018, 9, 1791-1796. | 2.5 | 7 |
| 107 | Prognostic Role of Carcinoembryonic Antigen Level after Preoperative Chemoradiotherapy in Patients with Rectal Cancer. Journal of Gastrointestinal Surgery, 2018, 22, 1772-1778. | 1.7 | 7 |
| 108 | The Impact of Primary Tumor Sidedness on the Effect of Regorafenib in Refractory Metastatic Colorectal Cancer. Journal of Cancer, 2019, 10, 1611-1615. | 2.5 | 7 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 109 | Capecitabine plus Oxaliplatin as a Second-Line Therapy for Advanced Biliary Tract Cancers: A Multicenter, Open-Label, Phase II Trial. Journal of Cancer, 2019, 10, 6185-6190. | 2.5 | 7 |
| 110 | Detection of Fusion Genes Using a Targeted RNA Sequencing Panel in Gastrointestinal and Rare Cancers. Journal of Oncology, 2020, 2020, 1-8. | 1.3 | 7 |
| 111 | Incidence of FGFR2 Amplification and FGFR2 Fusion in Patients with Metastatic Cancer Using Clinical Sequencing. Journal of Oncology, 2022, 2022, 1-9. | 1.3 | 7 |
| 112 | Pemetrexed Monotherapy as Salvage Treatment in Patients with Metastatic Colorectal Cancer Refractory to Standard Chemotherapy: A Phase II Single-arm Prospective Trial. Journal of Cancer, 2018, 9, 2910-2915. | 2.5 | 6 |
| 113 | TPK1 as a predictive marker for the anti-tumour effects of simvastatin in gastric cancer. Pathology Research and Practice, 2020, 216, 152820. | 2.3 | 6 |
| 114 | Programmed Death Ligand 1 Expression as a Prognostic Marker in Patients with Advanced Biliary Tract Cancer. Oncology, 2021, 99, 365-372. | 1.9 | 6 |
| 115 | Prognostic significance of survivin in rectal cancer patients treated with surgery and postoperative concurrent chemo-radiation therapy. Oncotarget, 2016, 7, 62676-62686. | 1.8 | 6 |
| 116 | Can we omit prophylactic inguinal nodal irradiation in anal cancer patients?. Radiation Oncology Journal, 2015, 33, 83. | 1.5 | 6 |
| 117 | HER2 Aberrations as a Novel Marker in Advanced Biliary Tract Cancer. Frontiers in Oncology, 2022, 12, 834104. | 2.8 | 6 |
| 118 | Oxaliplatin (3 months <i>>v</i> 6 months) With 6 Months of Fluoropyrimidine as Adjuvant Therapy in Patients With Stage II/III Colon Cancer: KCSG CO09-07. Journal of Clinical Oncology, 2022, 40, 3868-3877. | 1.6 | 6 |
| 119 | Immunohistochemical Detection of p53 Expression in Patients with Preoperative Chemoradiation for Rectal Cancer: Association with Prognosis. Yonsei Medical Journal, 2015, 56, 82. | 2.2 | 5 |
| 120 | Efficacy and Safety of FOLFIRI Regimen in Elderly Versus Nonelderly Patients with Metastatic Colorectal or Gastric Cancer. Oncologist, 2017, 22, 293-303. | 3.7 | 5 |
| 121 | Antitumor activity of sorafenib plus CDK4/6 inhibitor in pancreatic patient derived cell with KRAS mutation. Journal of Cancer, 2018, 9, 3394-3399. | 2.5 | 5 |
| 122 | The impact of primary tumor site on outcomes of treatment with etoposide and cisplatin in grade 3 gastroenteropancreatic neuroendocrine carcinoma. Journal of Cancer, 2019, 10, 3140-3144. | 2.5 | 5 |
| 123 | Systematic Evaluation of Gastric Tumor Cell Index and Two-Drug Combination Therapy via 3-Dimensional High-Throughput Drug Screening. Frontiers in Oncology, 2019, 9, 1327. | 2.8 | 5 |
| 124 | Use of Gefitinib in EGFR-Amplified Refractory Solid Tumors: An Open-Label, Single-Arm, Single-Center Prospective Pilot Study. Targeted Oncology, 2020, 15, 185-192. | 3.6 | 5 |
| 125 | The prevalence of homologous recombination deficiency (HRD) in various solid tumors and the role of HRD as a single biomarker to immune checkpoint inhibitors. Journal of Cancer Research and Clinical Oncology, 2022, 148, 2427-2435. | 2.5 | 5 |
| 126 | The impact of pathologic differentiation (well/poorly) and the degree of Ki-67 index in patients with metastatic WHO grade 3 GEP-NECs. Oncotarget, 2017, 8, 73974-73980. | 1.8 | 5 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 127 | Effect of leukocyte alteration on treatment outcomes following preoperative chemoradiotherapy in patients with rectal cancer. Radiation Oncology Journal, 2017, 35, 217-226. | 1.5 | 5 |
| 128 | Carcinoembryonic Antigen Improves the Performance of Magnetic Resonance Imaging in the Prediction of Pathologic Response after Neoadjuvant Chemoradiation for Patients with Rectal Cancer. Cancer Research and Treatment, 2020, 52, 446-454. | 3.0 | 5 |
| 129 | The use of regorafenib for patients with refractory metastatic colorectal cancer in clinical practice. OncoTargets and Therapy, 2019, Volume 12, 225-231. | 2.0 | 4 |
| 130 | Prognostic Factors of Survival with Aflibercept and FOLFIRI (fluorouracil, leucovorin, irinotecan) as Second-line Therapy for Patients with Metastatic Colorectal Cancer. Journal of Cancer, 2021, 12, 460-466. | 2.5 | 4 |
| 131 | Neutralizing antibody to FGFR2 can act as a selective biomarker and potential therapeutic agent for gastric cancer with FGFR2 amplification. American Journal of Translational Research (discontinued), 2019, 11, 4508-4515. | 0.0 | 4 |
| 132 | Whole-Genome and Transcriptome Sequencing Identified NOTCH2 and HES1 as Potential Markers of Response to Imatinib in Desmoid Tumor (Aggressive Fibromatosis): A Phase II Trial Study. Cancer Research and Treatment, 2022, 54, 1240-1255. | 3.0 | 4 |
| 133 | 332â€Novel TGF. \hat{l}^2 signatures in metastatic colorectal cancer patients treated with vactosertib in combination with pembrolizumab. , 2020, , . | | 3 |
| 134 | Phase Ib dose-escalation study of Pexa-Vec (pexastimogene devacirepvec; JX-594), an oncolytic and immunotherapeutic vaccinia virus, administered by intravenous (IV) infusions in patients with metastatic colorectal carcinoma (mCRC) Journal of Clinical Oncology, 2013, 31, 3608-3608. | 1.6 | 3 |
| 135 | Prognostic and predictive value of liver volume in colorectal cancer patients with unresectable liver metastases. Radiation Oncology Journal, 2014, 32, 77. | 1.5 | 3 |
| 136 | Placebo-controlled, double-blinded multi-center phase III trial of XELIRI/FOLFIRI plus simvastatin in metastatic colorectal cancer Journal of Clinical Oncology, 2015, 33, 3576-3576. | 1.6 | 3 |
| 137 | Light-to-Moderate Alcohol Consumption Increases the Risk of Biliary Tract Cancer in Prediabetes and Diabetes, but Not in Normoglycemic Status: A Nationwide Cohort Study. Journal of Clinical Oncology, 2022, 40, 3623-3632. | 1.6 | 3 |
| 138 | Selective colony area method for heterogeneous patient-derived tumor cell lines in anti-cancer drug screening system. PLoS ONE, 2019, 14, e0215080. | 2.5 | 2 |
| 139 | Phase I clinical trial of KML001 monotherapy in patients with advanced solid tumors. Expert Opinion on Investigational Drugs, 2020, 29, 1059-1067. | 4.1 | 2 |
| 140 | Phase I trial and pharmacokinetic study of Tanibirumab, a fully human monoclonal antibody to the vascular endothelial growth factor receptor 2 in patients with refractory solid tumors Journal of Clinical Oncology, 2015, 33, 2522-2522. | 1.6 | 2 |
| 141 | Nintedanib (N) plus best supportive care (BSC) versus placebo plus BSC for the treatment of patients (pts) with metastatic colorectal cancer (mCRC) refractory to standard therapies: Health-related quality of life (HRQoL) results of the Phase III LUME-Colon 1 study Journal of Clinical Oncology, 2017, 35, 671-671. | 1.6 | 2 |
| 142 | First-in-human phase I trial of anti-hepatocyte growth factor (HGF) antibody (YYB101) in refractory solid tumor patients: Integrative pathologic-genomic analysis and the final results Journal of Clinical Oncology, 2019, 37, 3104-3104. | 1.6 | 2 |
| 143 | Prospective phase II trial of pazopanib plus CapeOX (capecitabine and oxaliplatin) in previously untreated patients with advanced gastric cancer Journal of Clinical Oncology, 2015, 33, 4049-4049. | 1.6 | 2 |
| 144 | 74â€Tumor microenvironment based on PD-L1 and CD8 T-cell infiltration correlated with the response of MSS mCRC patients treated vactosertib in combination with pembrolizumab. , 2021, 9, A82-A82. | | 2 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Phase 1b study of vactosertib in combination with nal-IRI plus 5FU/LV in patients with metastatic pancreatic ductal adenocarcinoma who have failed first-line gemcitabine/ <i>nab</i> journal of Clinical Oncology, 2022, 40, TPS632-TPS632. | 1.6 | 2 |
| 146 | Adjuvant chemoradiation with 5-fluorouracil/leucovorin versus S-1 in gastric cancer patients following D2 lymph node dissection surgery: a feasibility study. Anticancer Research, 2014, 34, 6585-91. | 1.1 | 2 |
| 147 | To Excavate Biomarkers Predictive of the Response for Capecitabine plus RAD001 through Nanostring-Based Multigene Assay in Advanced Gastric Cancer Patients. Journal of Cancer, 2016, 7, 2173-2178. | 2.5 | 1 |
| 148 | Clinical Outcomes of Salvage Chemoradiotherapy for Locally Recurrent Biliary Tract Cancer. Tumori, 2017, 103, 345-352. | 1.1 | 1 |
| 149 | Pemetrexed/Erlotinib as a Salvage Treatment in Patients with High EGFR-Expressing Metastatic Colorectal Cancer Following Failure of Standard Chemotherapy: A Phase II Single-Arm Prospective Study. Targeted Oncology, 2020, 15, 67-73. | 3.6 | 1 |
| 150 | A phase 1 dose-escalation and dose-expansion study to assess the safety and efficacy of CKD-516, a novel vascular disrupting agent, in combination with Irinotecan in patients with previously treated metastatic colorectal cancer. Investigational New Drugs, 2021, 39, 1335-1347. | 2.6 | 1 |
| 151 | ATM Expression as a Prognostic Marker in Patients With Advanced Biliary Tract Cancer Treated With First-line Gemcitabine and Platinum Chemotherapy. In Vivo, 2021, 35, 499-505. | 1.3 | 1 |
| 152 | A randomized phase II study of neoadjuvant chemoradiotherapy with 5-FU/leucovorin or irinotecan/S1 in patients with locally advanced rectal cancer Journal of Clinical Oncology, 2013, 31, 511-511. | 1.6 | 1 |
| 153 | Interim safety analysis from a randomized, open-label, multicenter, phase III trial of capecitabine plus oxaliplatin (XELOX) versus capecitabine (X) as first-line chemotherapy in elderly patients with advanced gastric cancer (AGC) Journal of Clinical Oncology, 2014, 32, 158-158. | 1.6 | 1 |
| 154 | Phase II XELOX + lapatinib treatment in HER2-amplified gastric cancer: Monitoring with serial cell-free DNA genomics Journal of Clinical Oncology, 2017, 35, e15610-e15610. | 1.6 | 1 |
| 155 | First-in-human phase I trial of anti-hepatocyte growth factor (HGF) antibody (YYB101) in refractory solid tumor patients Journal of Clinical Oncology, 2018, 36, e14501-e14501. | 1.6 | 1 |
| 156 | A prospective multicenter phase II study of sunitinib in patients with advanced aggressive fibromatosis Journal of Clinical Oncology, 2013, 31, 10589-10589. | 1.6 | 1 |
| 157 | Determining Which Patients Require Preoperative Pelvic Radiotherapy Before Curative-Intent Surgery and/or Ablation for Metastatic Rectal Cancer. Annals of Surgical Oncology, 2022, , 1. | 1.5 | 1 |
| 158 | Phase 1b study of vactosertib in combination with oxaliplatin with 5FU/LV (FOLFOX) in patients with metastatic pancreatic cancer who have failed first-line gemcitabine/ <i>nab</i> paclitaxel Journal of Clinical Oncology, 2022, 40, e16299-e16299. | 1.6 | 1 |
| 159 | A phase I study of IMC-001, a PD-L1 blocker, in patients with metastatic or locally advanced solid tumors. Investigational New Drugs, 2021, 39, 1624-1632. | 2.6 | 0 |
| 160 | SSTR2A Protein Expression in Neuroendocrine Neoplasms of the Colorectum. Korean Journal of Pathology, 2011, 45, 276. | 1.3 | 0 |
| 161 | Perceptions of clinical trials in gastrointestinal/hepatobiliary cancer patients: A comprehensive survey in a Korean tertiary hospital Journal of Clinical Oncology, 2012, 30, e13094-e13094. | 1.6 | 0 |
| 162 | A phase Ib dose escalation study of JX-594 (TK-vaccinia virus expressing GM-CSF) administered by biweekly intravenous infusion in patients with metastatic, refractory colorectal carcinoma Journal of Clinical Oncology, 2012, 30, e13044-e13044. | 1.6 | 0 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 163 | A prospective phase II study of neoadjuvant FOLFOX6 plus cetuximab in patients with colorectal cancer and unresectable liver metastasis Journal of Clinical Oncology, 2012, 30, e14072-e14072. | 1.6 | O |
| 164 | Open-label, phase IIIb, multicenter, expanded access study of everolimus in patients with advanced neuroendocrine tumors (NET) Journal of Clinical Oncology, 2013, 31, 4138-4138. | 1.6 | 0 |
| 165 | Pathway redundancy and feedback loop as drivers of nonresponse in colorectal cancer treatments Journal of Clinical Oncology, 2013, 31, e14626-e14626. | 1.6 | 0 |
| 166 | Global clinical experience with adjuvant oxaliplatin/5-fluorouracil (5-FU)-based chemotherapy for colon cancer: Outcomes of the ACCElox registry Journal of Clinical Oncology, 2013, 31, e14624-e14624. | 1.6 | 0 |
| 167 | Molecular profiling of patient derived cells (PDCs) from metastatic cancer patients using CancerSCAN: Highly profiled models to test the efficacy of genome-directed therapy in cancer Journal of Clinical Oncology, 2015, 33, e22241-e22241. | 1.6 | 0 |
| 168 | Longitudinal follow-up of quality of life in gastrointestinal cancer patients after curative surgery in South Korea Journal of Clinical Oncology, 2016, 34, 697-697. | 1.6 | 0 |
| 169 | The clinicopathologic features and treatment of 607 hindgut neuroendocrine tumor (NET) patients at a single institution Journal of Clinical Oncology, 2016, 34, 4091-4091. | 1.6 | 0 |
| 170 | The impact of cetuximab plus AKT- or mTOR- inhibitor in patient-derived colon cancer cell model with RAS wild type and PIK3CA mutation Journal of Clinical Oncology, 2016, 34, e15153-e15153. | 1.6 | 0 |
| 171 | Programmed death (PD)-ligand 1 (L1) expression and mismatch repair (MMR) deficiency across tumor types: Candidates for checkpoint inhibitor based immunotherapy Journal of Clinical Oncology, 2017, 35, e14622-e14622. | 1.6 | 0 |
| 172 | The impact of pathologic differentiation (well/ poorly) and the degree of Ki-67 index in patients with metastatic WHO grade 3 GEP-NECs Journal of Clinical Oncology, 2017, 35, e15686-e15686. | 1.6 | 0 |
| 173 | VariantPlex panel to detect genomic aberrations in oncology patients with rare cancer type Journal of Clinical Oncology, 2018, 36, e24234-e24234. | 1.6 | 0 |
| 174 | Detection of targetable fusions using FusionPlex in oncology patients Journal of Clinical Oncology, 2018, 36, e24238-e24238. | 1.6 | 0 |
| 175 | Impact of the prior chemotherapy with two different fluoropyrimidines on the efficacy of CapeIRI or FOLFIRI in metastatic colorectal cancer: An exploratory analysis of the phase III AXEPT trial Journal of Clinical Oncology, 2019, 37, 711-711. | 1.6 | 0 |
| 176 | Detection of circulating tumor cells (CTCs) in cerebrospinal fluid of a patient with HER2-overexpressing gastric cancer and single cell analysis of intra-patient heterogeneity of CTCs. Translational Cancer Research, 2019, 8, 2107-2112. | 1.0 | 0 |
| 177 | A clinical scoring system for survival prediction in advanced gastric cancer Journal of Clinical Oncology, 2020, 38, 436-436. | 1.6 | 0 |
| 178 | Pemetrexed plus erlotinib as a salvage treatment in high EGFR-expressing metastatic colorectal cancer patients following failure of standard chemotherapy: A phase II single-arm prospective study Journal of Clinical Oncology, 2020, 38, 104-104. | 1.6 | 0 |
| 179 | 823â€Spatial analysis of tumor-infiltrating lymphocytes correlates with the response of metastatic colorectal cancer patients treated with vactosertib in combination with pembrolizumab., 2021, 9, A861-A861. | | 0 |
| 180 | Safety and effectiveness of aflibercept in combination with FOLFIRI in Korean patients with metastatic colorectal cancer who received oxaliplatin-containing regimen. Journal of Cancer Research and Clinical Oncology, 2022, , 1. | 2.5 | 0 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | ASO Visual Abstract: Determining Which Patients Require Preoperative Pelvic Radiotherapy Before Curative Intent Surgery and/or Ablation for Metastatic Rectal Cancer. Annals of Surgical Oncology, 2022, , . | 1.5 | O |