

Ana Maria Afonso

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1808061/publications.pdf>

Version: 2024-02-01

104
papers

3,907
citations

101543

36
h-index

133252

59
g-index

104
all docs

104
docs citations

104
times ranked

3381
citing authors

#	ARTICLE	IF	CITATIONS
1	Solid-phase microextraction coatings based on the metal-organic framework ZIF-8: Ensuring stable and reusable fibers. <i>Talanta</i> , 2020, 215, 120910.	5.5	36
2	Use of a pH-sensitive polymer in a microextraction and preconcentration method directly combined with high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1619, 460910.	3.7	10
3	Ionic liquid-based miniaturized aqueous biphasic system to develop an environmental-friendly analytical preconcentration method. <i>Talanta</i> , 2019, 203, 305-313.	5.5	13
4	A guanidinium ionic liquid-based surfactant as an adequate solvent to separate and preconcentrate cadmium and copper in water using <i>in situ</i> dispersive liquid-liquid microextraction. <i>Analytical Methods</i> , 2018, 10, 1529-1537.	2.7	11
5	Salt-induced ionic liquid-based microextraction using a low cytotoxic guanidinium ionic liquid and liquid chromatography with fluorescence detection to determine monohydroxylated polycyclic aromatic hydrocarbons in urine. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 4701-4713.	3.7	25
6	Guanidinium ionic liquid-based surfactants as low cytotoxic extractants: Analytical performance in an <i>in-situ</i> dispersive liquid-liquid microextraction method for determining personal care products. <i>Journal of Chromatography A</i> , 2018, 1559, 102-111.	3.7	31
7	Insights in the analytical performance of neat metal-organic frameworks in the determination of pollutants of different nature from waters using dispersive miniaturized solid-phase extraction and liquid chromatography. <i>Talanta</i> , 2018, 179, 775-783.	5.5	52
8	Influence of Ligand Functionalization of UiO-66-Based Metal-Organic Frameworks When Used as Sorbents in Dispersive Solid-Phase Analytical Microextraction for Different Aqueous Organic Pollutants. <i>Molecules</i> , 2018, 23, 2869.	3.8	40
9	A green metal-organic framework to monitor water contaminants. <i>RSC Advances</i> , 2018, 8, 31304-31310.	3.6	34
10	Vacuum-assisted headspace-solid phase microextraction for determining volatile free fatty acids and phenols. Investigations on the effect of pressure on competitive adsorption phenomena in a multicomponent system. <i>Analytica Chimica Acta</i> , 2017, 962, 41-51.	5.4	53
11	Monitoring trihalomethanes and nitrogenous disinfection by-products in blending desalinated waters using solid-phase microextraction and gas chromatography. <i>Environmental Technology (United Kingdom)</i> , 2017, 38, 911-922.	2.2	11
12	Effect of the inclusion of banana silage in the diet of goats on physicochemical and sensory characteristics of cheeses at different ripening times. <i>Small Ruminant Research</i> , 2017, 149, 52-61.	1.2	4
13	Influence of vegetable coagulant and ripening time on the lipolytic and sensory profile of cheeses made with raw goat milk from Canary breeds. <i>Food Science and Technology International</i> , 2017, 23, 254-264.	2.2	11
14	Metal-organic frameworks as novel sorbents in dispersive-based microextraction approaches. <i>TrAC - Trends in Analytical Chemistry</i> , 2017, 90, 114-134.	11.4	119
15	Monitoring trihalomethanes in chlorinated waters using a dispersive liquid-liquid microextraction method with a non-chlorinated organic solvent and gas chromatography-mass spectrometry. <i>Environmental Technology (United Kingdom)</i> , 2017, 38, 718-729.	2.2	5
16	Utilization of highly robust and selective crosslinked polymeric ionic liquid-based sorbent coatings in direct-immersion solid-phase microextraction and high-performance liquid chromatography for determining polar organic pollutants in waters. <i>Talanta</i> , 2016, 158, 125-133.	5.5	60
17	Ionic liquids and derivatives in gas chromatography. , 2016, , 45-82.		1
18	Magnetic ionic liquids as non-conventional extraction solvents for the determination of polycyclic aromatic hydrocarbons. <i>Analytica Chimica Acta</i> , 2016, 934, 106-113.	5.4	64

#	ARTICLE	IF	CITATIONS
19	A magnetic-based dispersive micro-solid-phase extraction method using the metal-organic framework HKUST-1 and ultra-high-performance liquid chromatography with fluorescence detection for determining polycyclic aromatic hydrocarbons in waters and fruit tea infusions. <i>Journal of Chromatography A</i> , 2016, 1436, 42-50.	3.7	100
20	Analytical Applications of Ionic Liquids in Chromatographic and Electrophoretic Separation Techniques. <i>Green Chemistry and Sustainable Technology</i> , 2016, , 193-233.	0.7	2
21	Ionic liquids versus ionic liquid-based surfactants in dispersive liquid-liquid microextraction for determining copper in water by flame atomic absorption spectrometry. <i>International Journal of Environmental Analytical Chemistry</i> , 2016, 96, 101-118.	3.3	31
22	The metal-organic framework HKUST-1 as efficient sorbent in a vortex-assisted dispersive micro solid-phase extraction of parabens from environmental waters, cosmetic creams, and human urine. <i>Talanta</i> , 2015, 139, 13-20.	5.5	144
23	Automated direct-immersion solid-phase microextraction using crosslinked polymeric ionic liquid sorbent coatings for the determination of water pollutants by gas chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 4615-4627.	3.7	25
24	Interfacial and aggregation behavior of dicationic and tricationic ionic liquid-based surfactants in aqueous solution. <i>Colloids and Surfaces A: Physicochemical and Engineering Aspects</i> , 2015, 469, 224-234.	4.7	32
25	A simplified vortex-assisted emulsification microextraction method for determining personal care products in environmental water samples by ultra-high-performance liquid chromatography. <i>Analytical Methods</i> , 2015, 7, 1825-1833.	2.7	12
26	Double salts of ionic-liquid-based surfactants in microextraction: application of their mixed hemimicelles as novel sorbents in magnetic-assisted micro-dispersive solid-phase extraction for the determination of phenols. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 8753-8764.	3.7	26
27	Polymeric ionic liquid coatings versus commercial solid-phase microextraction coatings for the determination of volatile compounds in cheeses. <i>Talanta</i> , 2014, 121, 153-162.	5.5	55
28	Vortex-assisted emulsification microextraction followed by in-syringe ultrasound-assisted back-microextraction to determine haloacetic acids in waters. <i>Analytical Methods</i> , 2014, 6, 4115-4123.	2.7	9
29	Multiple headspace solid-phase microextraction for quantifying volatile free fatty acids in cheeses. <i>Talanta</i> , 2014, 129, 183-190.	5.5	19
30	Ionic liquids in dispersive liquid-liquid microextraction. <i>TrAC - Trends in Analytical Chemistry</i> , 2013, 51, 87-106.	11.4	246
31	Utilization of an ionic liquid <i>in situ</i> preconcentration method for the determination of the 15 + 1 European Union polycyclic aromatic hydrocarbons in drinking water and fruit tea infusions. <i>Journal of Separation Science</i> , 2013, 36, 2496-2506.	2.5	13
32	Monitoring polycyclic aromatic hydrocarbons in seawaters and wastewaters using a dispersive liquid-liquid microextraction method. <i>Environmental Technology (United Kingdom)</i> , 2013, 34, 607-616.	2.2	11
33	An in-situ extraction-preconcentration method using ionic liquid-based surfactants for the determination of organic contaminants contained in marine sediments. <i>Talanta</i> , 2012, 99, 972-983.	5.5	57
34	Surface-bonded ionic liquid stationary phases in high-performance liquid chromatography-A review. <i>Analytica Chimica Acta</i> , 2012, 714, 20-37.	5.4	103
35	A novel preconcentration strategy for extraction methods based on common cationic surfactants: An alternative to classical coacervative extraction. <i>Journal of Chromatography A</i> , 2012, 1257, 9-18.	3.7	18
36	A novel in situ preconcentration method with ionic liquid-based surfactants resulting in enhanced sensitivity for the extraction of polycyclic aromatic hydrocarbons from toasted cereals. <i>Journal of Chromatography A</i> , 2012, 1227, 29-37.	3.7	58

#	ARTICLE	IF	CITATIONS
37	Headspace-single drop microextraction (HS-SDME) in combination with high-performance liquid chromatography (HPLC) to evaluate the content of alkyl- and methoxy-phenolic compounds in biomass smoke. <i>Talanta</i> , 2011, 85, 1265-1273.	5.5	26
38	Use of ionic liquid aggregates of 1-hexadecyl-3-butyl imidazolium bromide in a focused-microwave assisted extraction method followed by high-performance liquid chromatography with ultraviolet and fluorescence detection to determine the 15+1 EU priority PAHs in toasted cereals (â€œgofiosâ€œ). <i>Talanta</i> , 2011, 85, 1199-1206.	5.5	42
39	Factors That Affect the Content of Heterocyclic Aromatic Amines in Foods. <i>Comprehensive Reviews in Food Science and Food Safety</i> , 2011, 10, 52-108.	11.7	215
40	Ionic liquids as a tool for determination of metals and organic compounds in food analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2011, 30, 1598-1619.	11.4	63
41	In-situ ionic liquid-dispersive liquid-liquid microextraction method to determine endocrine disrupting phenols in seawaters and industrial effluents. <i>Mikrochimica Acta</i> , 2011, 174, 213-222.	5.0	59
42	Developing qualitative extraction profiles of coffee aromas utilizing polymeric ionic liquid sorbent coatings in headspace solid-phase microextraction gas chromatographyâ€“mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2011, 401, 2965-2976.	3.7	36
43	Suitability of ionic liquids as mobileâ€“phase additives in HPLC with fluorescence and UV detection for the determination of heterocyclic aromatic amines. <i>Journal of Separation Science</i> , 2010, 33, 182-190.	2.5	22
44	Determination of water pollutants by direct-immersion solid-phase microextraction using polymeric ionic liquid coatings. <i>Journal of Chromatography A</i> , 2010, 1217, 1236-1243.	3.7	105
45	Utilization of a benzyl functionalized polymeric ionic liquid for the sensitive determination of polycyclic aromatic hydrocarbons; parabens and alkylphenols in waters using solid-phase microextraction coupled to gas chromatographyâ€“flame ionization detection. <i>Journal of Chromatography A</i> , 2010, 1217, 7189-7197.	3.7	122
46	Dispersive liquidâ€“liquid microextraction versus single-drop microextraction for the determination of several endocrine-disrupting phenols from seawaters. <i>Talanta</i> , 2010, 80, 1611-1618.	5.5	130
47	Ionic liquids as desorption solvents and memory effect suppressors in heterocyclic aromatic amines determination by SPMEâ€“HPLC fluorescence. <i>Analytical and Bioanalytical Chemistry</i> , 2009, 394, 937-946.	3.7	49
48	Ionic liquids as mobile phase additives in high-performance liquid chromatography with electrochemical detection: Application to the determination of heterocyclic aromatic amines in meat-based infant foods. <i>Talanta</i> , 2009, 79, 590-597.	5.5	67
49	Determination of carbonyl compounds in smoke samples: strategies for sampling and standardization. <i>Journal of Environmental Monitoring</i> , 2009, 11, 1043.	2.1	1
50	Analytical methods applied to the determination of heterocyclic aromatic amines in foods. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 862, 15-42.	2.3	92
51	Micelle-mediated extractions using nonionic surfactant mixtures and HPLC-UV to determine endocrine-disrupting phenols in seawaters. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 391, 735-744.	3.7	27
52	The ionic liquid 1-hexadecyl-3-methylimidazolium bromide as novel extracting system for polycyclic aromatic hydrocarbons contained in sediments using focused microwave-assisted extraction. <i>Journal of Chromatography A</i> , 2008, 1182, 145-152.	3.7	87
53	Coupling the extraction efficiency of imidazolium-based ionic liquid aggregates with solid-phase microextraction-gas chromatographyâ€“mass spectrometry. <i>Journal of Chromatography A</i> , 2008, 1214, 23-29.	3.7	33
54	Exposure to heterocyclic aromatic amines from the consumption of cooked red meat and its effect on human cancer risk: A review. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2008, 25, 2-24.	2.3	72

#	ARTICLE	IF	CITATIONS
55	Metabolism and toxicology of heterocyclic aromatic amines when consumed in diet: Influence of the genetic susceptibility to develop human cancer. A review. Food Research International, 2008, 41, 327-340.	6.2	34
56	Estimation of Uncertainty in the Analysis of Carbonyl Compounds by HPLC-UV Using DNPH Derivatization. Journal of Liquid Chromatography and Related Technologies, 2007, 31, 361-381.	1.0	8
57	Evaluation of the Uncertainty Associated to the Determination of Heavy Metals in Seawater Using Graphite Furnace Atomic Absorption Spectrometry. Analytical Letters, 2007, 40, 3322-3342.	1.8	4
58	Monitoring chlorophenols in industrial effluents by solid-phase microextraction-gas chromatography-mass spectrometry. International Journal of Environmental Analytical Chemistry, 2007, 87, 159-175.	3.3	12
59	Focused Microwave-Assisted Extraction and HPLC with Electrochemical Detection to Determine Heterocyclic Amines in Meat Extracts. Journal of Liquid Chromatography and Related Technologies, 2007, 30, 27-42.	1.0	13
60	Determination of the alkyl- and methoxy-phenolic content in wood extractives by micellar solid-phase microextraction and gas chromatography-mass spectrometry. Talanta, 2007, 73, 505-513.	5.5	15
61	Focused microwave-assisted micellar extraction combined with solid-phase microextraction-gas chromatography/mass spectrometry to determine chlorophenols in wood samples. Analytica Chimica Acta, 2007, 582, 10-18.	5.4	24
62	Determination of less polar heterocyclic amines in meat extracts. Analytica Chimica Acta, 2007, 582, 259-266.	5.4	43
63	Micellar solid-phase microextraction for determining partition coefficients of substituted polycyclic aromatic hydrocarbons in micellar media: possible prediction of hydrocarbon-micelle behaviour. Analytical and Bioanalytical Chemistry, 2007, 387, 2271-2281.	3.7	5
64	Experimental Design Optimization of Solid-Phase Microextraction Conditions for the Determination of Heterocyclic Aromatic Amines by High-Performance Liquid Chromatography. Analytical Letters, 2006, 39, 405-423.	1.8	19
65	Correlations Between Phenols-Micelles Partition Coefficients and Several Molecular Descriptors. An Approach to Predict the Phenols Behaviour in MSPME. Chromatographia, 2006, 63, 167-174.	1.3	8
66	Optimization of an analytical methodology for the determination of alkyl- and methoxy-phenolic compounds by HS-SPME in biomass smoke. Analytical and Bioanalytical Chemistry, 2006, 385, 1162-1171.	3.7	17
67	Biosynthesis of Antitumoral and Bactericidal Sanguinarine. Journal of Biomedicine and Biotechnology, 2006, 2006, 1-6.	3.0	9
68	Study of the interactions between phenolic compounds and micellar media using micellar solid-phase microextraction/gas chromatography. Journal of Chromatography A, 2005, 1099, 64-74.	3.7	18
69	Emissions of polycyclic aromatic hydrocarbons from combustion of agricultural and sylvicultural debris. Atmospheric Environment, 2005, 39, 6654-6663.	4.1	48
70	Polycyclic Aromatic Hydrocarbons in Smoke Used to Smoke Cheese Produced by the Combustion of Rock Rose (Cistus monspeliensis) and Tree Heather (Erica arborea) Wood. Journal of Agricultural and Food Chemistry, 2005, 53, 176-182.	5.2	35
71	Coupling micelle-mediated extraction using mixtures of surfactants and fluorescence measurements with a fiber-optic for the screening of PAHs in seawater. Analyst, The, 2005, 130, 571-577.	3.5	13
72	Solid-phase microextraction coupled with high-performance liquid chromatography for the analysis of heterocyclic aromatic amines. Journal of Chromatography A, 2004, 1030, 87-93.	3.7	52

#	ARTICLE	IF	CITATIONS
73	Optimization of a sampling method to determine polycyclic aromatic hydrocarbons in smoke from incomplete biomass combustion. <i>Analytica Chimica Acta</i> , 2004, 524, 287-294.	5.4	15
74	Nonionic surfactant mixtures: a new cloud-point extraction approach for the determination of PAHs in seawater using HPLC with fluorimetric detection. <i>Analytica Chimica Acta</i> , 2004, 518, 165-172.	5.4	105
75	Solid-Phase Microextraction Coupled to Gas Chromatography/Mass Spectrometry for Determining Polycyclic Aromatic Hydrocarbon Micelle Partition Coefficients. <i>Analytical Chemistry</i> , 2004, 76, 4572-4578.	6.5	31
76	Micellar microwave-assisted extraction combined with solid-phase microextraction for the determination of polycyclic aromatic hydrocarbons in a certified marine sediment. <i>Analytica Chimica Acta</i> , 2003, 477, 81-91.	5.4	75
77	Reversed Phase Liquid Chromatographic Method for Separation and Determination of Positional Isomeric Mono- and Di-substituted Anilines and Phenols on an R,S-Hydroxypropyl Ether Cyclodextrin Column. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2003, 26, 1-15.	1.0	7
78	DETERMINATION OF N-NITROSODIMETHYLAMINE BY HPLC, WITH FLUORESCENCE DETECTION. A SURVEY OF N-NITROSODIMETHYLAMINE IN COMMERCIAL BEERS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2002, 25, 977-984.	1.0	9
79	Optimizing Conditions for the Extraction of Pigments in Cochineals (<i>Dactylopius coccus</i> Costa) Using Response Surface Methodology. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 6968-6974.	5.2	38
80	Fast microwave-assisted dansylation of N-nitrosamines. <i>Journal of Chromatography A</i> , 2002, 946, 133-140.	3.7	42
81	Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration. <i>Journal of Chromatography A</i> , 2002, 949, 291-299.	3.7	71
82	Ultrasonic micellar extraction of polycyclic aromatic hydrocarbons from marine sediments. <i>Talanta</i> , 2001, 54, 15-23.	5.5	36
83	Cloud-point preconcentration and HPLC determination of polycyclic aromatic hydrocarbons in marine sediments. <i>Fresenius' Journal of Analytical Chemistry</i> , 2001, 371, 526-531.	1.5	29
84	Micellar Extraction of Polycyclic Aromatic Hydrocarbons from Certified Marine Sediment. <i>International Journal of Environmental Analytical Chemistry</i> , 2001, 81, 281-294.	3.3	17
85	Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media. <i>Journal of Chromatography A</i> , 2000, 869, 515-522.	3.7	93
86	NON-LINEAR CALIBRATION IN QUANTITATIVE ANALYSIS BY HPTLC UTILIZING A FIBRE OPTIC FLUORESCENCE DETECTOR. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2000, 23, 2653-2668.	1.0	4
87	Rapid microwave-assisted dansylation of biogenic amines. <i>Journal of Chromatography A</i> , 1998, 808, 87-93.	3.7	26
88	Selective Analysis of Fluorene by Quenched Fluorescence in Cetylpyridinium Bromide Micelles. <i>Microchemical Journal</i> , 1998, 60, 101-109.	4.5	13
89	Effect of non-ionic surfactants as mobile phase additives on the fluorescence intensity of dansyl derivatives of biogenic amines in high-performance thin-layer chromatography. <i>Analyst</i> , 1998, 123, 725-729.	3.5	17
90	Quantitative Analysis of Biogenic Amines by High-Performance Thin-Layer Chromatography Utilizing a Fibre Optic Fluorescence Detector. <i>Analytical Letters</i> , 1998, 31, 475-489.	1.8	11

#	ARTICLE	IF	CITATIONS
91	Fluorescence Quenching of Polycyclic Aromatic Hydrocarbons by Cetylpyridinium Bromide: Discrimination between Alternant and Nonalternant Hydrocarbons. <i>Applied Spectroscopy</i> , 1997, 51, 380-386.	2.2	17
92	Effects of cetylpyridinium bromide micelles on the spectrofluorimetric characteristics of polycyclic aromatic hydrocarbons. <i>Talanta</i> , 1997, 44, 257-267.	5.5	18
93	Selective determination of acenaphthene in mixtures of three-ring polycyclic aromatic hydrocarbons by fluorescence quenching in micellar medium of cetylpyridinium bromide. <i>Journal of Fluorescence</i> , 1997, 7, 147-153.	2.5	5
94	Effects of cationic micelles on fluorescence of indole and indolecarboxylic acids. Analytical determinations. <i>Mikrochimica Acta</i> , 1995, 118, 153-162.	5.0	5
95	Degradation of carbaryl in natural waters: Enhanced hydrolysis rate in micellar solution. <i>Bulletin of Environmental Contamination and Toxicology</i> , 1992, 48, 171-8.	2.7	4
96	Simultaneous synchronous fluorimetric determination of benzo[a]pyrene and perylene in micellar media. <i>Analytica Chimica Acta</i> , 1991, 255, 107-111.	5.4	22
97	Spectrofluorimetric determination of carbaryl and 1-naphthol in micellar media. <i>Mikrochimica Acta</i> , 1991, 103, 171-179.	5.0	5
98	Bioluminescence, chemiluminescence. <i>Fresenius' Journal of Analytical Chemistry</i> , 1990, 337, 86-96.	1.5	4
99	1,5-bis-(2,3-dihydroxy-phenylmethylene)-thiocarbohydrazone as reagent for the spectrofluorimetric determination of nanogram amounts of gallium. <i>Mikrochimica Acta</i> , 1990, 100, 55-61.	5.0	1
100	Spectrofluorometric determination of zinc with 1,5-bis(2,3-dihydroxyphenylmethylene) thiocarbohydrazone. <i>Analytica Chimica Acta</i> , 1987, 202, 207-213.	5.4	4
101	Spectrofluorimetric determination of zinc with pyrocatechol-1-aldehyde 2-pyridylhydrazone. <i>Analyst</i> , 1986, 111, 327.	3.5	7
102	Kinetic spectrofluorimetric determination of silver, based on its catalytic effect on the oxidation of pyrocatechol-1-aldehyde 2-pyridylhydrazone by peroxodisulphate in the presence of 1,10-phenanthroline as activator. <i>Talanta</i> , 1986, 33, 779-783.	5.5	5
103	Pyrocatechol-1-Aldehyde 2-Benzothiazolylhydrazone As Reagent for the Spectrofluorimetric Determination of Nanogram Amounts of Gallium in Urine and Blood Serum. <i>Analytical Letters</i> , 1985, 18, 1003-1012.	1.8	3
104	Pyrocatechol-1-aldehyde salicyloylhydrazone as reagent for the spectrofluorimetric determination of zinc. <i>Mikrochimica Acta</i> , 1984, 83, 53-60.	5.0	2