

Salvador Resino

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1812664/publications.pdf>

Version: 2024-02-01

269
papers

4,804
citations

136950

32
h-index

189892

50
g-index

287
all docs

287
docs citations

287
times ranked

7047
citing authors

#	ARTICLE	IF	CITATIONS
1	Epidemiological trend of hospitalized acute and recurrent hepatitis C in Spain. <i>Annals of Medicine</i> , 2024, 51, 68-68.	3.8	0
2	Antibody levels to SARS-CoV-2 spike protein in mothers and children from delivery to six months later. <i>Birth</i> , 2023, 50, 418-427.	2.2	5
3	Low anti-SARS-CoV-2 S antibody levels predict increased mortality and dissemination of viral components in the blood of critical COVID-19 patients. <i>Journal of Internal Medicine</i> , 2022, 291, 232-240.	6.0	21
4	Similar humoral immune responses against the SARS-CoV-2 spike protein in HIV and non-HIV individuals after COVID-19. <i>Journal of Infection</i> , 2022, 84, 418-467.	3.3	7
5	Blood microbiome is associated with changes in portal hypertension after successful direct-acting antiviral therapy in patients with HCV-related cirrhosis. <i>Journal of Antimicrobial Chemotherapy</i> , 2022, 77, 719-726.	3.0	7
6	HCV eradication with DAAs differently affects HIV males and females: A whole miRNA sequencing characterization. <i>Biomedicine and Pharmacotherapy</i> , 2022, 145, 112405.	5.6	3
7	Differences in the hepatitis C virus cascade of care and time to initiation of therapy among vulnerable subpopulations using a mobile unit as point-of-care. <i>Liver International</i> , 2022, 42, 309-319.	3.9	7
8	Baseline and time-updated factors in preclinical development of anionic dendrimers as successful anti-HIV vaginal microbicides. <i>Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology</i> , 2022, 14, e1774.	6.1	5
9	Reply to Pati et al. <i>Journal of Infectious Diseases</i> , 2022, , .	4.0	0
10	Metabolomic changes after DAAs therapy are related to the improvement of cirrhosis and inflammation in HIV/HCV-coinfected patients. <i>Biomedicine and Pharmacotherapy</i> , 2022, 147, 112623.	5.6	6
11	Plasma miRNA profile at COVID-19 onset predicts severity status and mortality. <i>Emerging Microbes and Infections</i> , 2022, 11, 676-688.	6.5	44
12	Early innate immune response triggered by the human respiratory syncytial virus and its regulation by ubiquitination/deubiquitination processes. <i>Journal of Biomedical Science</i> , 2022, 29, 11.	7.0	6
13	High SARS-CoV-2 Viral Load and Low CCL5 Expression Levels in the Upper Respiratory Tract Are Associated With COVID-19 Severity. <i>Journal of Infectious Diseases</i> , 2022, 225, 977-982.	4.0	20
14	Misdiagnosis rate among negative COVID-19 patients in real-life with Panbio COVID-19 antigen rapid test during 2021. <i>Journal of Infection</i> , 2022, , .	3.3	0
15	High Plasma sTNF-R1 Level Is Related to Loss of Natural HIV Control in Long-Term Elite Controllers. <i>Frontiers in Cellular and Infection Microbiology</i> , 2022, 12, 858872.	3.9	1
16	Environmental factors linked to hospital admissions in young children due to acute viral lower respiratory infections: A bidirectional case-crossover study. <i>Environmental Research</i> , 2022, 212, 113319.	7.5	2
17	Negative impact of HIV infection on broad-spectrum anti-HCV neutralizing antibody titers in HCV-infected patients with advanced HCV-related cirrhosis. <i>Biomedicine and Pharmacotherapy</i> , 2022, 150, 113024.	5.6	1
18	OLFM4 polymorphisms predict septic shock survival after major surgery. <i>European Journal of Clinical Investigation</i> , 2021, 51, e13416.	3.4	3

#	ARTICLE	IF	CITATIONS
19	Strategies Targeting the Innate Immune Response for the Treatment of Hepatitis C Virus-Associated Liver Fibrosis. <i>Drugs</i> , 2021, 81, 419-443.	10.9	12
20	TRPM5 rs886277 Polymorphism Predicts Hepatic Fibrosis Progression in Non-Cirrhotic HCV-Infected Patients. <i>Journal of Clinical Medicine</i> , 2021, 10, 483.	2.4	1
21	The Challenging Road to Hepatitis C Virus Eradication. <i>Journal of Clinical Medicine</i> , 2021, 10, 611.	2.4	5
22	Successful HCV Therapy Reduces Liver Disease Severity and Inflammation Biomarkers in HIV/HCV-Coinfected Patients With Advanced Cirrhosis: A Cohort Study. <i>Frontiers in Medicine</i> , 2021, 8, 615342.	2.6	11
23	HCV eradication with IFN-based therapy does not completely restore gene expression in PBMCs from HIV/HCV-coinfected patients. <i>Journal of Biomedical Science</i> , 2021, 28, 23.	7.0	6
24	Impact of the Economic Crisis on Endocarditis Mortality in Spain: A Nationwide Study. <i>International Journal of Health Services</i> , 2021, , 002073142110123.	2.5	1
25	Prevalence and factors associated with SARS-CoV-2 seropositivity in the Spanish HIV Research Network Cohort. <i>Clinical Microbiology and Infection</i> , 2021, 27, 1678-1684.	6.0	34
26	HCV screening based on dried blood samples and linkage to care in people who use drugs: A prospective study. <i>International Journal of Drug Policy</i> , 2021, 92, 103134.	3.3	11
27	HIV screening and retention in care in people who use drugs in Madrid, Spain: a prospective study. <i>Infectious Diseases of Poverty</i> , 2021, 10, 111.	3.7	1
28	HCV Cure With Direct-Acting Antivirals Improves Liver and Immunological Markers in HIV/HCV-Coinfected Patients. <i>Frontiers in Immunology</i> , 2021, 12, 723196.	4.8	14
29	Detection of active hepatitis C in a single visit and linkage to care among marginalized people using a mobile unit in Madrid, Spain. <i>International Journal of Drug Policy</i> , 2021, 96, 103424.	3.3	10
30	Age-Adjusted Endothelial Activation and Stress Index for Coronavirus Disease 2019 at Admission Is a Reliable Predictor for 28-Day Mortality in Hospitalized Patients With Coronavirus Disease 2019. <i>Frontiers in Medicine</i> , 2021, 8, 736028.	2.6	4
31	Are Reduced Levels of Coagulation Proteins Upon Admission Linked to COVID-19 Severity and Mortality?. <i>Frontiers in Medicine</i> , 2021, 8, 718053.	2.6	7
32	IL-1R1 rs6755229 polymorphism is related to death in patients undergoing major surgery who develop septic shock: a retrospective study. <i>Infectious Diseases</i> , 2021, , 1-4.	2.8	0
33	CEACAM7 polymorphisms predict genetic predisposition to mortality in post-surgical septic shock patients. <i>Journal of Microbiology, Immunology and Infection</i> , 2021, , .	3.1	0
34	Different HCV Exposure Drives Specific miRNA Profile in PBMCs of HIV Patients. <i>Biomedicines</i> , 2021, 9, 1627.	3.2	2
35	DBP rs7041 and DHCR7 rs3829251 are Linked to CD4+ Recovery in HIV Patients on Antiretroviral Therapy. <i>Frontiers in Pharmacology</i> , 2021, 12, 773848.	3.5	0
36	Mild profile improvement of immune biomarkers in HIV/HCV-coinfected patients who removed hepatitis C after HCV treatment: A prospective study. <i>Journal of Infection</i> , 2020, 80, 99-110.	3.3	9

#	ARTICLE	IF	CITATIONS
37	Persistent HIV controllers are more prone to spontaneously clear HCV: a retrospective cohort study. <i>Journal of the International AIDS Society</i> , 2020, 23, e25607.	3.0	2
38	The PANDEMYC Score. An Easily Applicable and Interpretable Model for Predicting Mortality Associated With COVID-19. <i>Journal of Clinical Medicine</i> , 2020, 9, 3066.	2.4	31
39	Near normalization of peripheral blood markers in HIV-infected patients on long-term suppressive antiretroviral therapy: a case-control study. <i>Aids</i> , 2020, 34, 1891-1897.	2.2	4
40	Comparison of methods and characterization of small RNAs from plasma extracellular vesicles of HIV/HCV coinfecting patients. <i>Scientific Reports</i> , 2020, 10, 11140.	3.3	22
41	MTHFR rs1801133 Polymorphism Is Associated With Liver Fibrosis Progression in Chronic Hepatitis C: A Retrospective Study. <i>Frontiers in Medicine</i> , 2020, 7, 582666.	2.6	4
42	Epidemic history and baseline resistance to NS5A-specific direct acting drugs of hepatitis C virus in Spain. <i>Scientific Reports</i> , 2020, 10, 13024.	3.3	1
43	Effects of Hepatitis C Virus (HCV) Eradication on Bone Mineral Density in Human Immunodeficiency Virus/HCV-Coinfected Patients. <i>Clinical Infectious Diseases</i> , 2020, 73, e2026-e2033.	5.8	2
44	Telomere Length Increase in HIV/HCV-Coinfected Patients with Cirrhosis after HCV Eradication with Direct-Acting Antivirals. <i>Journal of Clinical Medicine</i> , 2020, 9, 2407.	2.4	5
45	Viral RNA load in plasma is associated with critical illness and a dysregulated host response in COVID-19. <i>Critical Care</i> , 2020, 24, 691.	5.8	185
46	Environmental factors are associated with hospital admissions for sepsis-related pneumonia: A bidirectional case-crossover design. <i>Environmental Research</i> , 2020, 191, 110102.	7.5	4
47	Persistence of Clinically Significant Portal Hypertension After Eradication of Hepatitis C Virus in Patients With Advanced Cirrhosis. <i>Clinical Infectious Diseases</i> , 2020, 71, 2726-2729.	5.8	23
48	Epidemiological Trend of Sepsis in Patients with Hospital Admissions Related to Hepatitis C in Spain (2000-2015): A Nationwide Study. <i>Journal of Clinical Medicine</i> , 2020, 9, 1607.	2.4	2
49	IFNL3 rs12980275 Polymorphism Predicts Septic Shock-Related Death in Patients Undergoing Major Surgery: A Retrospective Study. <i>Frontiers in Medicine</i> , 2020, 7, 186.	2.6	1
50	Epidemiological trend of hepatitis C-related liver events in Spain (2000-2015): A nationwide population-based study. <i>European Journal of Internal Medicine</i> , 2020, 75, 84-92.	2.2	7
51	Metabolic changes during respiratory syncytial virus infection of epithelial cells. <i>PLoS ONE</i> , 2020, 15, e0230844.	2.5	35
52	Hepatitis C virus vaccine design: focus on the humoral immune response. <i>Journal of Biomedical Science</i> , 2020, 27, 78.	7.0	23
53	Innate Immune Response against Hepatitis C Virus: Targets for Vaccine Adjuvants. <i>Vaccines</i> , 2020, 8, 313.	4.4	12
54	Liver Stiffness Hinders Normalization of Systemic Inflammation and Endothelial Activation after Hepatitis C Virus (HCV) Eradication in HIV/HCV Coinfected Patients. <i>Vaccines</i> , 2020, 8, 323.	4.4	5

#	ARTICLE	IF	CITATIONS
55	Plasma IP-10 and IL-6 are linked to Child-Pugh B cirrhosis in patients with advanced HCV-related cirrhosis: a cross-sectional study. <i>Scientific Reports</i> , 2020, 10, 10384.	3.3	5
56	Plasma metabolomic fingerprint of advanced cirrhosis stages among HIV/HCV-coinfected and HCV-monoinfected patients. <i>Liver International</i> , 2020, 40, 2215-2227.	3.9	11
57	Downregulation of A20 Expression Increases the Immune Response and Apoptosis and Reduces Virus Production in Cells Infected by the Human Respiratory Syncytial Virus. <i>Vaccines</i> , 2020, 8, 100.	4.4	11
58	Effects of Eradication of HCV on Cardiovascular Risk and Preclinical Atherosclerosis in HIV/HCV-Coinfected Patients. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2020, 83, 292-300.	2.1	5
59	Gender-based vulnerability in women who inject drugs in a harm reduction setting. <i>PLoS ONE</i> , 2020, 15, e0230886.	2.5	20
60	Brief Report: CYP27B1 rs10877012 T Allele Was Linked to Non-AIDS Progression in ART-Naïve HIV-Infected Patients: A Retrospective Study. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2020, 85, 659-664.	2.1	2
61	CD4 recovery is associated with genetic variation in IFN β and IL19 genes. <i>Antiviral Research</i> , 2019, 170, 104577.	4.1	7
62	National Temporal Trend Analysis of Infective Endocarditis among Patients Infected with HIV in Spain (1997-2014): A Retrospective Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 1167.	2.4	4
63	European mitochondrial haplogroups predict liver-related outcomes in patients coinfecting with HIV and HCV: a retrospective study. <i>Journal of Translational Medicine</i> , 2019, 17, 244.	4.4	6
64	Nosocomial Vs. Community-Acquired Infective Endocarditis in Spain: Location, Trends, Clinical Presentation, Etiology, and Survival in the 21st Century. <i>Journal of Clinical Medicine</i> , 2019, 8, 1755.	2.4	22
65	DBP rs16846876 and rs12512631 polymorphisms are associated with progression to AIDS naïve HIV-infected patients: a retrospective study. <i>Journal of Biomedical Science</i> , 2019, 26, 83.	7.0	2
66	Rapid decrease in titer and breadth of neutralizing anti-HCV antibodies in HIV/HCV-coinfected patients who achieved SVR. <i>Scientific Reports</i> , 2019, 9, 12163.	3.3	2
67	MicroRNA Profile of HCV Spontaneous Clarified Individuals, Denotes Previous HCV Infection. <i>Journal of Clinical Medicine</i> , 2019, 8, 849.	2.4	11
68	IL7RA rs6897932 Polymorphism is Associated with Better CD4+ T-Cell Recovery in HIV Infected Patients Starting Combination Antiretroviral Therapy. <i>Biomolecules</i> , 2019, 9, 233.	4.0	9
69	Evaluation of the diagnostic accuracy of laboratory-based screening for hepatitis C in dried blood spot samples: A systematic review and meta-analysis. <i>Scientific Reports</i> , 2019, 9, 7316.	3.3	35
70	siRNA-Mediated Simultaneous Regulation of the Cellular Innate Immune Response and Human Respiratory Syncytial Virus Replication. <i>Biomolecules</i> , 2019, 9, 165.	4.0	5
71	VDR rs2228570 Polymorphism Is Related to Non-Progression to AIDS in Antiretroviral Therapy Naïve HIV-Infected Patients. <i>Journal of Clinical Medicine</i> , 2019, 8, 311.	2.4	9
72	TNFAIP3, TNIP1, and MyD88 Polymorphisms Predict Septic-Shock-Related Death in Patients Who Underwent Major Surgery. <i>Journal of Clinical Medicine</i> , 2019, 8, 283.	2.4	5

#	ARTICLE	IF	CITATIONS
73	Impact of DARC rs12075 Variants on Liver Fibrosis Progression in Patients with Chronic Hepatitis C: A Retrospective Study. <i>Biomolecules</i> , 2019, 9, 143.	4.0	7
74	Genetic variation in CCR2 and CXCL12 genes impacts on CD4 restoration in patients initiating cART with advanced immunosuppression. <i>PLoS ONE</i> , 2019, 14, e0214421.	2.5	11
75	HCV-coinfection is related to an increased HIV-1 reservoir size in cART-treated HIV patients: a cross-sectional study. <i>Scientific Reports</i> , 2019, 9, 5606.	3.3	22
76	Prevalence of hepatitis E infection in HIV/HCV-coinfected patients in Spain (2012–2014). <i>Scientific Reports</i> , 2019, 9, 1143.	3.3	8
77	Lower expression of plasma-derived exosome miR-21 levels in HIV-1 elite controllers with decreasing CD4 T cell count. <i>Journal of Microbiology, Immunology and Infection</i> , 2019, 52, 667-671.	3.1	14
78	Genetic variants upstream of TNFAIP3 in the 6q23 region are associated with liver disease severity in HIV/HCV-coinfected patients: A cross-sectional study. <i>Infection, Genetics and Evolution</i> , 2019, 67, 112-120.	2.3	2
79	PNPLA3 rs738409 polymorphism is associated with liver fibrosis progression in patients with chronic hepatitis C: A repeated measures study. <i>Journal of Clinical Virology</i> , 2018, 103, 71-74.	3.1	10
80	Association of CD14 rs2569190 polymorphism with mortality in shock septic patients who underwent major cardiac or abdominal surgery: A retrospective study. <i>Scientific Reports</i> , 2018, 8, 2698.	3.3	7
81	Elevated liver stiffness is linked to increased biomarkers of inflammation and immune activation in HIV/hepatitis C virus-coinfected patients. <i>Aids</i> , 2018, 32, 1095-1105.	2.2	28
82	Evaluation of dried blood spot samples for screening of hepatitis C and human immunodeficiency virus in a real-world setting. <i>Scientific Reports</i> , 2018, 8, 1858.	3.3	34
83	Pegylated Interferon- α -Induced Natural Killer Cell Activation Is Associated With Human Immunodeficiency Virus-1 DNA Decline in Antiretroviral Therapy-Treated HIV-1/Hepatitis C Virus-Coinfected Patients. <i>Clinical Infectious Diseases</i> , 2018, 66, 1910-1917.	5.8	30
84	Frecuencia de sustituciones relevantes asociadas a resistencia en la región NS5A a elbasvir en el virus de la hepatitis C en pacientes con genotipo 1a en España. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2018, 36, 262-267.	0.5	2
85	Epidemiological trends of deep venous thrombosis in HIV-infected subjects (1997–2013): A nationwide population-based study in Spain. <i>European Journal of Internal Medicine</i> , 2018, 48, 69-74.	2.2	3
86	Bacterial translocation and clinical progression of HCV-related cirrhosis in HIV-infected patients. <i>Journal of Viral Hepatitis</i> , 2018, 25, 180-186.	2.0	4
87	Mitochondrial haplogroup H is related to CD4+ T cell recovery in HIV infected patients starting combination antiretroviral therapy. <i>Journal of Translational Medicine</i> , 2018, 16, 343.	4.4	6
88	Trends in pulmonary embolism in patients infected with HIV during the combination antiretroviral therapy era in Spain: A nationwide population-based study. <i>Scientific Reports</i> , 2018, 8, 12137.	3.3	4
89	The Myeloid-Epithelial-Reproductive Tyrosine Kinase (MERTK) rs4374383 Polymorphism Predicts Progression of Liver Fibrosis in Hepatitis C Virus-Infected Patients: A Longitudinal Study. <i>Journal of Clinical Medicine</i> , 2018, 7, 473.	2.4	15
90	Dysregulation of the Immune System in HIV/HCV-Coinfected Patients According to Liver Stiffness Status. <i>Cells</i> , 2018, 7, 196.	4.1	14

#	ARTICLE	IF	CITATIONS
91	Surgery for acute infective endocarditis: epidemiological data from a Spanish nationwide hospital-based registry. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2018, 27, 498-504.	1.1	13
92	The IL7RA rs6897932 polymorphism is associated with progression of liver fibrosis in patients with chronic hepatitis C: Repeated measurements design. <i>PLoS ONE</i> , 2018, 13, e0197115.	2.5	10
93	Vitamin D in Human Immunodeficiency Virus Infection: Influence on Immunity and Disease. <i>Frontiers in Immunology</i> , 2018, 9, 458.	4.8	110
94	High Plasma Levels of sTNF-R1 and CCL11 Are Related to CD4+ T-Cells Fall in Human Immunodeficiency Virus Elite Controllers With a Sustained Virologic Control. <i>Frontiers in Immunology</i> , 2018, 9, 1399.	4.8	3
95	Epidemiological trends of sepsis in the twenty-first century (2000–2013): an analysis of incidence, mortality, and associated costs in Spain. <i>Population Health Metrics</i> , 2018, 16, 4.	2.7	51
96	Prevalence of relevant NS5A resistance-associated substitutions to elbasvir in genotype 1a hepatitis C virus patients in Spain. <i>Enfermedades Infecciosas Y Microbiología Clínica (English Ed)</i> , 2018, 36, 262-267.	0.3	0
97	Mx1, OAS1 and OAS2 polymorphisms are associated with the severity of liver disease in HIV/HCV-coinfected patients: A cross-sectional study. <i>Scientific Reports</i> , 2017, 7, 41516.	3.3	22
98	IL-6 rs1800795 polymorphism is associated with septic shock-related death in patients who underwent major surgery: a preliminary retrospective study. <i>Annals of Intensive Care</i> , 2017, 7, 22.	4.6	12
99	G2-S16 dendrimer as a candidate for a microbicide to prevent HIV-1 infection in women. <i>Nanoscale</i> , 2017, 9, 9732-9742.	5.6	25
100	ADAR1 polymorphisms are related to severity of liver fibrosis in HIV/HCV-coinfected patients. <i>Scientific Reports</i> , 2017, 7, 12918.	3.3	7
101	IL7RA polymorphisms are not associated with AIDS progression. <i>European Journal of Clinical Investigation</i> , 2017, 47, 719-727.	3.4	3
102	Low frequency of NS5A relevant resistance-associated substitutions to Elbasvir among hepatitis C virus genotype 1a in Spain: a cross-sectional study. <i>Scientific Reports</i> , 2017, 7, 2892.	3.3	8
103	IL1B rs16944 polymorphism is related to septic shock and death. <i>European Journal of Clinical Investigation</i> , 2017, 47, 53-62.	3.4	17
104	CXCL9 and IL11 polymorphisms are associated with liver fibrosis in patients with chronic hepatitis C: a cross-sectional study. <i>Clinical and Translational Medicine</i> , 2017, 6, 26.	4.0	13
105	TRIM25 in the Regulation of the Antiviral Innate Immunity. <i>Frontiers in Immunology</i> , 2017, 8, 1187.	4.8	109
106	Efficacy of carbosilane dendrimers with an antiretroviral combination against HIV-1 in the presence of semen-derived enhancer of viral infection. <i>European Journal of Pharmacology</i> , 2017, 811, 155-163.	3.5	23
107	Stroke in HIV-infected individuals with and without HCV coinfection in Spain in the combination antiretroviral therapy era. <i>PLoS ONE</i> , 2017, 12, e0179493.	2.5	14
108	Liver stiffness measurement predicts liver-related events in patients with chronic hepatitis C: A retrospective study. <i>PLoS ONE</i> , 2017, 12, e0184404.	2.5	10

#	ARTICLE	IF	CITATIONS
109	Genetic Polymorphisms Associated with Liver Disease Progression in HIV/HCV-Coinfected Patients. <i>AIDS Reviews</i> , 2017, 19, 3-15.	1.0	14
110	Soluble Adhesion Molecules in Patients Coinfected with HIV and HCV: A Predictor of Outcome. <i>PLoS ONE</i> , 2016, 11, e0148537.	2.5	8
111	Environmental Factors Related to Pulmonary Tuberculosis in HIV-Infected Patients in the Combined Antiretroviral Therapy (cART) Era. <i>PLoS ONE</i> , 2016, 11, e0165944.	2.5	34
112	Trends in Epidemiology of COPD in HIV-Infected Patients in Spain (1997â€“2012). <i>PLoS ONE</i> , 2016, 11, e0166421.	2.5	11
113	Optimal vitamin D plasma levels are associated with lower bacterial DNA translocation in HIV/hepatitis c virus coinfecting patients. <i>Aids</i> , 2016, 30, 1069-1074.	2.2	7
114	<i>IL15</i> polymorphism is associated with advanced fibrosis, inflammationâ€“related biomarkers and virological response in human immunodeficiency virus/hepatitis C virus coinfection. <i>Liver International</i> , 2016, 36, 1258-1266.	3.9	5
115	Impact of patatin-like phospholipase domain-containing 3 gene polymorphism (rs738409) on severity of liver disease in HIV/hepatitis C virus-coinfecting patients. <i>Aids</i> , 2016, 30, 465-470.	2.2	12
116	Relationship of TRIM5 and TRIM22 polymorphisms with liver disease and HCV clearance after antiviral therapy in HIV/HCV coinfecting patients. <i>Journal of Translational Medicine</i> , 2016, 14, 257.	4.4	20
117	Impact of chronic hepatitis C on mortality in cirrhotic patients admitted to intensive-care unit. <i>BMC Infectious Diseases</i> , 2016, 16, 122.	2.9	5
118	Short Communication: <i>CXCL12</i> rs1029153 Polymorphism Is Associated with the Sustained Virological Response in HIV/Hepatitis C Virus-Coinfecting Patients on Hepatitis C Virus Therapy. <i>AIDS Research and Human Retroviruses</i> , 2016, 32, 226-231.	1.1	0
119	Fifteen-Year Trends in the Prevalence of Diabetes among Hospitalized HIV-Infected Patients in Spain (1997-2012). <i>PLoS ONE</i> , 2016, 11, e0161953.	2.5	3
120	NS3 Resistance-Associated Variants (RAVs) in Patients Infected with HCV Genotype 1a in Spain. <i>PLoS ONE</i> , 2016, 11, e0163197.	2.5	16
121	Relationship between ITPA polymorphisms and hemolytic anemia in HCV-infected patients after ribavirin-based therapy: a meta-analysis. <i>Journal of Translational Medicine</i> , 2015, 13, 320.	4.4	19
122	Association between IL7R polymorphisms and severe liver disease in HIV/HCV coinfecting patients: a cross-sectional study. <i>Journal of Translational Medicine</i> , 2015, 13, 206.	4.4	10
123	Trends in nontuberculous mycobacterial disease in hospitalized subjects in Spain (1997â€“2010) according to HIV infection. <i>HIV Medicine</i> , 2015, 16, 485-493.	2.2	8
124	Reply. <i>Hepatology</i> , 2015, 62, 1643-1643.	7.3	2
125	<i>IL7RA</i> polymorphisms predict the CD4+ recovery in HIV patients on cART. <i>European Journal of Clinical Investigation</i> , 2015, 45, 1192-1199.	3.4	12
126	<i>Pneumocystis</i> pneumonia in HIVâ€“positive patients in Spain: epidemiology and environmental risk factors. <i>Journal of the International AIDS Society</i> , 2015, 18, 19906.	3.0	23

#	ARTICLE	IF	CITATIONS
127	Single Nucleotide Polymorphisms of CXCL9-11 Chemokines Are Associated With Liver Fibrosis in HIV/HCV-Coinfected Patients. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2015, 68, 386-395.	2.1	11
128	TLR3 polymorphisms are associated with virologic response to hepatitis C virus (HCV) treatment in HIV/HCV coinfecting patients. <i>Journal of Clinical Virology</i> , 2015, 65, 62-67.	3.1	6
129	Association between IL7RA polymorphisms and the successful therapy against HCV in HIV/HCV-coinfecting patients. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2015, 34, 385-393.	2.9	4
130	Transcriptomic correlates of organ failure extent in sepsis. <i>Journal of Infection</i> , 2015, 70, 445-456.	3.3	81
131	Toll-like receptor 8 (TLR8) polymorphisms are associated with non-progression of chronic hepatitis C in HIV/HCV coinfecting patients. <i>Infection, Genetics and Evolution</i> , 2015, 36, 339-344.	2.3	6
132	Mitochondrial DNA haplogroups are associated with severe sepsis and mortality in patients who underwent major surgery. <i>Journal of Infection</i> , 2015, 70, 20-29.	3.3	17
133	rs7903146 Polymorphism at Transcription Factor 7 Like 2 Gene Is Associated with Total Cholesterol and Lipoprotein Profile in HIV/Hepatitis C Virus-Coinfecting Patients. <i>AIDS Research and Human Retroviruses</i> , 2015, 31, 326-334.	1.1	5
134	Relationship between European Mitochondrial Haplogroups and Chronic Renal Allograft Rejection in Patients with Kidney Transplant. <i>International Journal of Medical Sciences</i> , 2014, 11, 1129-1132.	2.5	3
135	FTOrs9939609 polymorphism is associated with metabolic disturbances and response to HCV therapy in HIV/HCV-coinfecting patients. <i>BMC Medicine</i> , 2014, 12, 198.	5.5	4
136	Association of adiponectin (ADIPOQ) rs2241766 polymorphism and dyslipidemia in HIV/HCV-coinfecting patients. <i>European Journal of Clinical Investigation</i> , 2014, 44, 453-462.	3.4	12
137	Mortality of patients infected with HIV in the intensive care unit (2005 through 2010): significant role of chronic hepatitis C and severe sepsis. <i>Critical Care</i> , 2014, 18, 475.	5.8	18
138	PPAR β Pro12Ala Polymorphism Is Associated With Sustained Virological Response in HIV/HCV-Coinfecting Patients Under HCV Therapy. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2014, 67, 113-119.	2.1	5
139	SLC30A8 rs13266634 polymorphism is related to a favorable cardiometabolic lipid profile in HIV/hepatitis C virus-coinfecting patients. <i>Aids</i> , 2014, 28, 1325-1332.	2.2	9
140	Incidence and mortality of tuberculosis disease in Spain between 1997 and 2010: Impact of human immunodeficiency virus (HIV) status. <i>Journal of Infection</i> , 2014, 68, 355-362.	3.3	12
141	Vitamin D deficiency is associated with severity of liver disease in HIV/HCV coinfecting patients. <i>Journal of Infection</i> , 2014, 68, 176-184.	3.3	28
142	CXCL9, CXCL10 and CXCL11 polymorphisms are associated with sustained virologic response in HIV/HCV-coinfecting patients. <i>Journal of Clinical Virology</i> , 2014, 61, 423-429.	3.1	13
143	IL28RA polymorphism (rs10903035) is associated with insulin resistance in HIV/HCV-coinfecting patients. <i>Journal of Viral Hepatitis</i> , 2014, 21, 189-197.	2.0	5
144	Relationship of vitamin D status with advanced liver fibrosis and response to hepatitis C virus therapy: A meta-analysis. <i>Hepatology</i> , 2014, 60, 1541-1550.	7.3	68

#	ARTICLE	IF	CITATIONS
145	PPAR β 2 Pro12Ala polymorphism was associated with favorable cardiometabolic risk profile in HIV/HCV coinfecting patients: a cross-sectional study. <i>Journal of Translational Medicine</i> , 2014, 12, 235.	4.4	11
146	European mitochondrial haplogroups are not associated with hepatitis C virus (HCV) treatment response in HIV/HCV-coinfecting patients. <i>HIV Medicine</i> , 2014, 15, 425-430.	2.2	5
147	ACSM4 Polymorphisms Are Associated With Rapid AIDS Progression in HIV-Infected Patients. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2014, 65, 27-32.	2.1	8
148	FTO rs9939609 polymorphism is associated with metabolic disturbances and response to HCV therapy in HIV/HCV-coinfecting patients. <i>BMC Medicine</i> , 2014, 12, 198.	5.5	7
149	Rate of candidiasis among HIV-infected children in Spain in the era of highly active antiretroviral therapy (1997-2008). <i>BMC Infectious Diseases</i> , 2013, 13, 115.	2.9	7
150	Meta-analysis: implications of interleukin-28B polymorphisms in spontaneous and treatment-related clearance for patients with hepatitis C. <i>BMC Medicine</i> , 2013, 11, 6.	5.5	80
151	Comment on: "Interleukin-28 polymorphisms on the SVR in the treatment of naïve chronic hepatitis C with pegylated interferon plus ribavirin: A meta-analysis". <i>Gene</i> , 2013, 522, 121.	2.2	2
152	IL28B polymorphism is associated with early hepatitis C virus (HCV) treatment failure in human immunodeficiency virus-coinfecting patients. <i>Journal of Viral Hepatitis</i> , 2013, 20, 358-366.	2.0	17
153	Association of torque teno virus (TTV) and torque teno mini virus (TTMV) with liver disease among patients coinfecting with human immunodeficiency virus and hepatitis C virus. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2013, 32, 289-297.	2.9	37
154	IL28B polymorphisms are associated with severity of liver disease in human immunodeficiency virus (HIV) patients coinfecting with hepatitis C virus. <i>Journal of Infection</i> , 2013, 66, 170-178.	3.3	13
155	HLA-E variants are associated with sustained virological response in HIV/hepatitis C virus-coinfecting patients on hepatitis C virus therapy. <i>Aids</i> , 2013, 27, 1231-1238.	2.2	15
156	Prediction of Hepatic Fibrosis in Patients Coinfecting With HIV and Hepatitis C Virus Based on Genetic Markers. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2013, 64, 434-442.	2.1	6
157	European mitochondrial haplogroups are associated with CD4+ T cell recovery in HIV-infected patients on combination antiretroviral therapy. <i>Journal of Antimicrobial Chemotherapy</i> , 2013, 68, 2349-2357.	3.0	17
158	Mitochondrial Haplogroups Are Associated With Clinical Pattern of AIDS Progression in HIV-Infected Patients. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2013, 63, 178-183.	2.1	21
159	Analysis of IL28B alleles with virologic response patterns and plasma cytokine levels in HIV/HCV-coinfecting patients. <i>Aids</i> , 2013, 27, 163-173.	2.2	12
160	Coinfection by human immunodeficiency virus and hepatitis C virus. <i>Current Opinion in Infectious Diseases</i> , 2012, 25, 564-569.	3.1	30
161	Plasma IL-6 and IL-9 predict the failure of interferon plus ribavirin therapy in HIV/HCV-coinfecting patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2012, 67, 1238-1245.	3.0	30
162	Trend of Pneumonia Incidence Among Children Infected With HIV in the Era of Highly Active Antiretroviral Therapy. <i>Pediatric Infectious Disease Journal</i> , 2012, 31, 599-601.	2.0	3

#	ARTICLE	IF	CITATIONS
163	Reduction in Mycobacterial Disease Among HIV-infected Children in the Highly Active Antiretroviral Therapy Era (1997–2008). <i>Pediatric Infectious Disease Journal</i> , 2012, 31, 278-283.	2.0	15
164	Bacterial DNA Translocation and Liver Disease Severity Among HIV-Infected Patients With Chronic Hepatitis C. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2012, 61, 552-556.	2.1	11
165	High plasma CXCL10 levels are associated with HCV-genotype 1, and higher insulin resistance, fibrosis, and HIV viral load in HIV/HCV coinfecting patients. <i>Cytokine</i> , 2012, 57, 25-29.	3.2	20
166	A combined score of pro- and anti-inflammatory interleukins improves mortality prediction in severe sepsis. <i>Cytokine</i> , 2012, 57, 332-336.	3.2	139
167	Genetic polymorphisms located in TGFB1, AGTR1, and VEGFA genes are associated to chronic renal allograft dysfunction. <i>Cytokine</i> , 2012, 58, 321-326.	3.2	17
168	Genetic polymorphisms located in genes related to immune and inflammatory processes are associated with end-stage renal disease: a preliminary study. <i>BMC Medical Genetics</i> , 2012, 13, 58.	2.1	9
169	Early natural killer cell counts in blood predict mortality in severe sepsis. <i>Critical Care</i> , 2011, 15, R243.	5.8	85
170	High plasma fractalkine (CX3CL1) levels are associated with severe liver disease in HIV/HCV co-infected patients with HCV genotype 1. <i>Cytokine</i> , 2011, 54, 244-248.	3.2	14
171	European Mitochondrial DNA Haplogroups and Metabolic Disorders in HIV/HCV-Coinfected Patients on Highly Active Antiretroviral Therapy. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2011, 58, 371-378.	2.1	22
172	Epidemiologic Trends of Cancer Diagnoses Among HIV-infected Children in Spain From 1997 to 2008. <i>Pediatric Infectious Disease Journal</i> , 2011, 30, 764-768.	2.0	21
173	Hepatitis C virus (HCV) treatment uptake and changes in the prevalence of HCV genotypes in HIV/HCV-coinfected patients. <i>Journal of Viral Hepatitis</i> , 2011, 18, 325-330.	2.0	31
174	Adipokine profiles and lipodystrophy in HIV-infected children during the first 4 years on highly active antiretroviral therapy. <i>HIV Medicine</i> , 2011, 12, 54-60.	2.2	12
175	Diagnostic accuracy of the APRI, FIB-4, and the Forns index for predicting liver fibrosis in HIV/HCV-coinfected patients: A validation study. <i>Journal of Infection</i> , 2011, 63, 402-405.	3.3	18
176	Soluble markers of inflammation are associated with Framingham scores in HIV-infected patients on suppressive antiretroviral therapy. <i>Journal of Infection</i> , 2011, 63, 382-390.	3.3	19
177	Soluble Fas and Fas ligand in HIV/HCV coinfecting patients and impact of HCV therapy. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2011, 30, 1213-1221.	2.9	3
178	Direct association between pharyngeal viral secretion and host cytokine response in severe pandemic influenza. <i>BMC Infectious Diseases</i> , 2011, 11, 232.	2.9	24
179	An artificial neural network improves the non-invasive diagnosis of significant fibrosis in HIV/HCV coinfecting patients. <i>Journal of Infection</i> , 2011, 62, 77-86.	3.3	31
180	Cardiovascular risk markers are increased in HIV-infected children with lipodystrophy syndrome. <i>Journal of Infection</i> , 2011, 62, 240-243.	3.3	4

#	ARTICLE	IF	CITATIONS
181	Sustained Virologic Response Decreases Serum Markers of Angiogenesis, Inflammation, and Fibrosis in HIV/HCV-Coinfected Patients on Hepatitis C Virus Therapy. <i>AIDS Patient Care and STDs</i> , 2011, 25, 131-133.	2.5	2
182	Sustained virological response to interferon- α plus ribavirin decreases inflammation and endothelial dysfunction markers in HIV/HCV co-infected patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2011, 66, 645-649.	3.0	29
183	European mitochondrial DNA haplogroups and liver fibrosis in HIV and hepatitis C virus coinfectd patients. <i>Aids</i> , 2011, 25, 1619-1926.	2.2	12
184	Plasma Interferon- β -Inducible Protein-10 Can Predict Virologic Response to Hepatitis C Virus Therapy in HIV/HCV-Coinfected Patients With HCV Genotype 1. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2010, 54, 219-220.	2.1	5
185	ASSOCIATION BETWEEN LIPODYSTROPHY AND LEPTIN IN HUMAN IMMUNODEFICIENCY VIRUS-1-INFECTED CHILDREN RECEIVING LOPINAVIR/RITONAVIR-BASED THERAPY. <i>Pediatric Infectious Disease Journal</i> , 2010, 29, 774-777.	2.0	4
186	Hepatitis C virus infection is associated with endothelial dysfunction in HIV/hepatitis C virus coinfectd patients. <i>Aids</i> , 2010, 24, 2059-2067.	2.2	39
187	Incidence of liver cirrhosis in HIV-infected patients with chronic hepatitis B or C in the era of highly active antiretroviral therapy. <i>Antiviral Therapy</i> , 2010, 15, 881-886.	1.0	35
188	Serum levels of adipokines in HIV/HCV co-infected patients and their association with insulin resistance and liver disease severity. <i>Journal of Infection</i> , 2010, 61, 499-501.	3.3	3
189	Can serum hyaluronic acid replace simple non-invasive indexes to predict liver fibrosis in HIV/Hepatitis C coinfectd patients?. <i>BMC Infectious Diseases</i> , 2010, 10, 244.	2.9	27
190	Diagnosis of advanced fibrosis in HIV and hepatitis C virus-coinfectd patients via a new noninvasive index: the HGM- β index. <i>HIV Medicine</i> , 2010, 11, 64-73.	2.2	13
191	CD81 expression in peripheral blood lymphocytes before and after treatment with interferon and ribavirin in HIV/HCV coinfectd patients. <i>HIV Medicine</i> , 2010, 11, 161-169.	2.2	3
192	Opportunistic infections and organ-specific diseases in HIV-1-infected children: a cohort study (1990-2006). <i>HIV Medicine</i> , 2010, 11, 245-252.	2.2	25
193	Association between plasma levels of eotaxin (CCL-11) and treatment response to interferon- α and ribavirin in HIV/HCV co-infected patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2010, 65, 303-306.	3.0	8
194	Modeling the Probability of Sustained Virological Response to Therapy with Pegylated Interferon plus Ribavirin in Patients Coinfectd with Hepatitis C Virus and HIV. <i>Clinical Infectious Diseases</i> , 2010, 51, 1209-1216.	5.8	56
195	Insulin Resistance Impairs Response to Interferon Plus Ribavirin in Patients Coinfectd With HIV and Hepatitis C Virus. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2010, 55, 176-181.	2.1	15
196	Host adaptive immunity deficiency in severe pandemic influenza. <i>Critical Care</i> , 2010, 14, R167.	5.8	145
197	First evidence of a pro-inflammatory response to severe infection with influenza virus H1N1. <i>Critical Care</i> , 2010, 14, 115.	5.8	10
198	Increased Th1, Th17 and pro-fibrotic responses in hepatitis C-infected patients are down-regulated after 12 weeks of treatment with pegylated interferon plus ribavirin. <i>European Cytokine Network</i> , 2010, 21, 84-91.	2.0	31

#	ARTICLE	IF	CITATIONS
199	Premature immunosenescence in HIV-infected patients on highly active antiretroviral therapy with low-level CD4 T cell repopulation. <i>Journal of Antimicrobial Chemotherapy</i> , 2009, 64, 579-588.	3.0	57
200	Rate and Timing of Hepatitis C Virus Relapse after a Successful Course of Pegylated Interferon plus Ribavirin in HIV-Infected and HIV-Uninfected Patients. <i>Clinical Infectious Diseases</i> , 2009, 49, 1397-1401.	5.8	35
201	Serum levels of fibrosis biomarkers measured early after liver transplantation are associated with severe hepatitis C virus recurrence. <i>Transplant Infectious Disease</i> , 2009, 11, 183-188.	1.7	26
202	High rate of infection and immune disorders in patients with hepatitis C virus after liver transplantation. <i>Transplant Infectious Disease</i> , 2009, 11, 367-372.	1.7	4
203	Insulin Resistance Is Associated With Advanced Liver Fibrosis and High Body Mass Index in HIV/HCV-Coinfected Patients. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2009, 50, 109-110.	2.1	22
204	Immunological predictors of CD4+T cell decline in antiretroviral treatment interruptions. <i>BMC Infectious Diseases</i> , 2008, 8, 20.	2.9	1
205	Negative influence of age on CD4+ cell recovery after highly active antiretroviral therapy in HIV-1-infected patients with severe immunodeficiency. <i>Journal of Infection</i> , 2008, 56, 130-136.	3.3	12
206	Association between Exposure to Nevirapine and Reduced Liver Fibrosis Progression in Patients with HIV and Hepatitis C Virus Coinfection. <i>Clinical Infectious Diseases</i> , 2008, 46, 137-143.	5.8	53
207	Lipid and Apoprotein Profile in HIV-1-Infected Patients After CD4-Guided Treatment Interruption. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2008, 48, 455-459.	2.1	12
208	Immunological Recovery and Metabolic Disorders in Severe Immunodeficiency HIV Type 1-Infected Children on Highly Active Antiretroviral Therapy. <i>AIDS Research and Human Retroviruses</i> , 2008, 24, 1477-1484.	1.1	11
209	Long-term response to highly active antiretroviral therapy with lopinavir/ritonavir in pre-treated vertically HIV-infected children. <i>Journal of Antimicrobial Chemotherapy</i> , 2007, 61, 183-190.	3.0	11
210	Short Communication: Immune Reconstitution after Autologous Peripheral Blood Stem Cell Transplantation in HIV-Infected Patients: Might Be Better Than Expected?. <i>AIDS Research and Human Retroviruses</i> , 2007, 23, 543-548.	1.1	24
211	SLOW PROGRESSION OF HUMAN IMMUNODEFICIENCY VIRUS AND HEPATITIS C VIRUS DISEASE IN A COHORT OF COINFECTED CHILDREN. <i>Pediatric Infectious Disease Journal</i> , 2007, 26, 846-849.	2.0	20
212	Functional patterns of HIV-1-specific CD4 T-cell responses in children are influenced by the extent of virus suppression and exposure. <i>Aids</i> , 2007, 21, 23-30.	2.2	20
213	Long-Term Response to Highly Active Antiretroviral Therapy in Human Immunodeficiency Virus and Hepatitis C Virus Coinfected Children. <i>Pediatric Infectious Disease Journal</i> , 2007, 26, 1061-1064.	2.0	9
214	NK Cell Increase in Neonates from the Preterm to the Full-Term Period of Gestation. <i>Neonatology</i> , 2007, 92, 158-163.	2.0	27
215	Persistence of proinflammatory response after severe respiratory syncytial virus disease in children. <i>Journal of Allergy and Clinical Immunology</i> , 2007, 119, 1547-1550.	2.9	19
216	Identification of liver fibrosis in HIV/HCV-coinfected patients using a simple predictive model based on routine laboratory data. <i>Journal of Viral Hepatitis</i> , 2007, 14, 070901052026008-???	2.0	28

#	ARTICLE	IF	CITATIONS
217	Long-Term Effects of Highly Active Antiretroviral Therapy in Pretreated, Vertically HIV Type 1-Infected Children: 6 Years of Follow-Up. <i>Clinical Infectious Diseases</i> , 2006, 42, 862-869.	5.8	73
218	Interleukin (IL)-1 β , IL-6 and IL-8 in nasal secretions: a common role for innate immunity in viral bronchial infection in infants?. <i>British Journal of Biomedical Science</i> , 2006, 63, 173-175.	1.3	5
219	LOW IMMUNOLOGIC RESPONSE TO HIGHLY ACTIVE ANTIRETROVIRAL THERAPY IN NAIVE VERTICALLY HUMAN IMMUNODEFICIENCY VIRUS TYPE 1-INFECTED CHILDREN WITH SEVERE IMMUNODEFICIENCY. <i>Pediatric Infectious Disease Journal</i> , 2006, 25, 365-368.	2.0	15
220	CD4+ T-Cell Immunodeficiency Is More Dependent on Immune Activation Than Viral Load in HIV-Infected Children on Highly Active Antiretroviral Therapy. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2006, 42, 269-276.	2.1	25
221	Impact of long-term viral suppression in CD4+ recovery of HIV-children on Highly Active Antiretroviral Therapy. <i>BMC Infectious Diseases</i> , 2006, 6, 10.	2.9	10
222	Effects of highly active antiretroviral therapy with nelfinavir in vertically HIV-1 infected children: 3 years of follow-up. Long-term response to nelfinavir in children. <i>BMC Infectious Diseases</i> , 2006, 6, 107.	2.9	8
223	Different profiles of immune reconstitution in children and adults with HIV-infection after highly active antiretroviral therapy. <i>BMC Infectious Diseases</i> , 2006, 6, 112.	2.9	29
224	Effectiveness and Safety of Abacavir, Lamivudine, and Zidovudine in Antiretroviral Therapy-Naive HIV-Infected Patients. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2006, 41, 154-159.	2.1	18
225	Clinical Outcomes Improve with Highly Active Antiretroviral Therapy in Vertically HIV Type 1-Infected Children. <i>Clinical Infectious Diseases</i> , 2006, 43, 243-252.	5.8	72
226	Interleukin-7 levels before highly active antiretroviral therapy may predict CD4+ T-cell recovery and virological failure in HIV-infected children. <i>Journal of Antimicrobial Chemotherapy</i> , 2006, 57, 798-800.	3.0	11
227	Antiretroviral activity and safety of lopinavir/ritonavir in protease inhibitor-experienced HIV-infected children with severe-moderate immunodeficiency. <i>Journal of Antimicrobial Chemotherapy</i> , 2006, 57, 579-582.	3.0	15
228	Lopinavir/r versus Nelfinavir as Salvage Therapy. <i>Pediatric Infectious Disease Journal</i> , 2005, 24, 392-393.	2.0	1
229	Mutations at codons 54 and 82 of HIV protease predict virological response of HIV-infected children on salvage lopinavir/ritonavir therapy. <i>Journal of Antimicrobial Chemotherapy</i> , 2005, 56, 1081-1086.	3.0	13
230	Immunological Changes after Highly Active Antiretroviral Therapy with Lopinavir-Ritonavir in Heavily Pretreated HIV-Infected Children. <i>AIDS Research and Human Retroviruses</i> , 2005, 21, 398-406.	1.1	10
231	Homeostatic role of IL-7 in HIV-1 infected children on HAART: Association with immunological and virological parameters. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2005, 94, 170-177.	1.5	7
232	Homeostatic role of IL-7 in HIV-1 infected children on HAART: Association with immunological and virological parameters. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2005, 94, 170-177.	1.5	9
233	Impact of Highly Active Antiretroviral Therapy on CD4+ T Cells and Viral Load of Children with AIDS: A Population-Based Study. <i>AIDS Research and Human Retroviruses</i> , 2004, 20, 927-931.	1.1	6
234	CD38 Expression in CD8+T Cells Predicts Virological Failure in HIV Type 1-Infected Children Receiving Antiretroviral Therapy. <i>Clinical Infectious Diseases</i> , 2004, 38, 412-417.	5.8	48

#	ARTICLE	IF	CITATIONS
235	Extensive Implementation of Highly Active Antiretroviral Therapy Shows Great Effect on Survival and Surrogate Markers in Vertically HIV-Infected Children. <i>Clinical Infectious Diseases</i> , 2004, 38, 1605-1612.	5.8	34
236	Positive virological outcome after lopinavir/ritonavir salvage therapy in protease inhibitor-experienced HIV-1-infected children: a prospective cohort study. <i>Journal of Antimicrobial Chemotherapy</i> , 2004, 54, 921-931.	3.0	21
237	Virological phenotype switches under salvage therapy with lopinavir-ritonavir in heavily pretreated HIV-1 vertically infected children. <i>Aids</i> , 2004, 18, 247-255.	2.2	24
238	HIV-infected children with moderate/severe immune-suppression: changes in the immune system after highly active antiretroviral therapy. <i>Clinical and Experimental Immunology</i> , 2004, 137, 570-577.	2.6	41
239	Immunity in HIV-1-Infected Adults with a Previous State of Moderate-Severe Immune-Suppression and More Than 500 CD4+ T Cell After Highly Active Antiretroviral Therapy. <i>Journal of Clinical Immunology</i> , 2004, 24, 379-388.	3.8	16
240	Salvage Lopinavir-Ritonavir Therapy in Human Immunodeficiency Virus-Infected Children. <i>Pediatric Infectious Disease Journal</i> , 2004, 23, 923-930.	2.0	41
241	Characterizing the immune system after long-term undetectable viral load in HIV-1-infected children. <i>Journal of Clinical Immunology</i> , 2003, 23, 279-289.	3.8	21
242	Increased interleukin-7 plasma levels are associated with recovery of CD4+ T cells in HIV-infected children. <i>Journal of Clinical Immunology</i> , 2003, 23, 401-406.	3.8	21
243	Stimulated proliferative responses in vertically HIV-infected children on HAART correlate with clinical and immunological markers. <i>Clinical and Experimental Immunology</i> , 2003, 131, 130-137.	2.6	5
244	Preserved immune system in long-term asymptomatic vertically HIV-1 infected children. <i>Clinical and Experimental Immunology</i> , 2003, 132, 105-112.	2.6	26
245	Recovery of T-cell subsets after antiretroviral therapy in HIV-infected children. <i>European Journal of Clinical Investigation</i> , 2003, 33, 619-627.	3.4	6
246	Neuroprotective effects of early antiretrovirals in vertical HIV infection. <i>Pediatric Neurology</i> , 2003, 29, 218-221.	2.1	32
247	Low Blood CD8+ T-Lymphocytes and High Circulating Monocytes Are Predictors of HIV-1-Associated Progressive Encephalopathy in Children. <i>Pediatrics</i> , 2003, 111, e168-e175.	2.1	56
248	Viral Load and CD4+ T Lymphocyte Response to Highly Active Antiretroviral Therapy in Human Immunodeficiency Virus Type 1-Infected Children: An Observational Study. <i>Clinical Infectious Diseases</i> , 2003, 37, 1216-1225.	5.8	54
249	CD8+ T-Cell Numbers Predict the Response to Antiviral Therapy in HIV-1-Infected Children. <i>Pediatric Research</i> , 2003, 53, 309-312.	2.3	7
250	CD8+ T-Cell Numbers Predict the Response to Antiviral Therapy in HIV-1-Infected Children. <i>Pediatric Research</i> , 2003, 53, 309-312.	2.3	6
251	Characterizing Immune Reconstitution after Long-Term Highly Active Antiretroviral Therapy in Pediatric AIDS. <i>AIDS Research and Human Retroviruses</i> , 2002, 18, 1395-1406.	1.1	32
252	Impact of antiretroviral protocols on dynamics of AIDS progression markers. <i>Archives of Disease in Childhood</i> , 2002, 86, 119-124.	1.9	19

#	ARTICLE	IF	CITATIONS
253	Dynamics of progression markers in a non-study population of human immunodeficiency virus-1 vertically infected infants with different antiretroviral treatments. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2002, 91, 776-782.	1.5	4
254	Dynamics of progression markers in a non-study population of human immunodeficiency virus-1 vertically infected infants with different antiretroviral treatments. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2002, 91, 776-782.	1.5	4
255	Immunological recovery after 3 years' antiretroviral therapy in HIV-1-infected children. <i>Aids</i> , 2002, 16, 483-486.	2.2	21
256	Naïve and memory CD4+ T cells and T cell activation markers in HIV-1 infected children on HAART. <i>Clinical and Experimental Immunology</i> , 2001, 125, 266-273.	2.6	35
257	Antibodies to an Epitope from the Cha Human Autoantigen Are Markers of Chagas' Disease. <i>Vaccine Journal</i> , 2001, 8, 1039-1043.	2.6	32
258	Association of CD8+T Lymphocyte Subsets with the Most Commonly Used Markers to Monitor HIV Type 1 Infection in Children Treated with Highly Active Antiretroviral Therapy. <i>AIDS Research and Human Retroviruses</i> , 2001, 17, 525-532.	1.1	22
259	Disruption in cytokine and chemokine production by T-cells in vertically HIV-1-infected children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2001, 90, 989-997.	1.5	11
260	Disruption in cytokine and chemokine production by T-cells in vertically HIV-1-infected children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2001, 90, 989-997.	1.5	8
261	Impaired interleukin-5 (IL-5) production by T cells as a prognostic marker of disease progression in human immunodeficiency virus type 1 (HIV-1)-infected children. <i>European Cytokine Network</i> , 2001, 12, 253-9.	2.0	7
262	Tumor necrosis factor-alpha and nitric oxide in vertically HIV-1-infected children: implications for pathogenesis. <i>European Cytokine Network</i> , 2001, 12, 437-44.	2.0	10
263	Correlación entre carga viral elevada y concentraciones de TNF- α y cICAM-1 en el plasma de niños infectados por el VIH-1. <i>Anales De Pediatr�a</i> , 2000, 52, 501-506.	0.2	5
264	Clinical Relevance of Cytokine Production in HIV-1 Infection in Children on Antiretroviral Therapy. <i>Scandinavian Journal of Immunology</i> , 2000, 52, 634-640.	2.7	0
265	Clinical Relevance of Cytokine Production in HIV-1 Infection in Children on Antiretroviral Therapy. <i>Scandinavian Journal of Immunology</i> , 2000, 52, 634-640.	2.7	11
266	Predictive Markers of Clinical Outcome in Vertically HIV-1-Infected Infants. A Prospective Longitudinal Study. <i>Pediatric Research</i> , 2000, 47, 509-515.	2.3	51
267	Correlation of Viral Load and CD8 T-Lymphocytes with Development of Neurological Manifestations in Vertically HIV-1-Infected Infants. A Prospective Longitudinal Study. <i>Neuropediatrics</i> , 1999, 30, 197-204.	0.6	20
268	Metabolic Profiling at COVID-19 Onset Shows Disease Severity and Sex-Specific Dysregulation. <i>Frontiers in Immunology</i> , 0, 13, .	4.8	14
269	Hepatitis E Virus Seroprevalence is Associated with Neurodegenerative Disorders in Older People with Dementia: A Case-Control Study. <i>Journal of Infectious Diseases</i> , 0, , .	4.0	2