

Tine Jepsen Nielsen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1835265/publications.pdf>

Version: 2024-02-01

146
papers

10,393
citations

38742

50
h-index

36028

97
g-index

147
all docs

147
docs citations

147
times ranked

13364
citing authors

#	ARTICLE	IF	CITATIONS
1	Prenatal Valproate Exposure and Risk of Autism Spectrum Disorders and Childhood Autism. JAMA - Journal of the American Medical Association, 2013, 309, 1696.	7.4	1,009
2	A Population-Based Study of Measles, Mumps, and Rubella Vaccination and Autism. New England Journal of Medicine, 2002, 347, 1477-1482.	27.0	595
3	Risk Factors for Autism: Perinatal Factors, Parental Psychiatric History, and Socioeconomic Status. American Journal of Epidemiology, 2005, 161, 916-925.	3.4	584
4	Epilepsy and risk of suicide: a population-based caseâ€“control study. Lancet Neurology, The, 2007, 6, 693-698.	10.2	427
5	Life expectancy and cardiovascular mortality in persons with schizophrenia. Current Opinion in Psychiatry, 2012, 25, 83-88.	6.3	426
6	Preâ€“pregnancy weight and the risk of stillbirth and neonatal death. BJOG: an International Journal of Obstetrics and Gynaecology, 2005, 112, 403-408.	2.3	367
7	Long-term risk of epilepsy after traumatic brain injury in children and young adults: a population-based cohort study. Lancet, The, 2009, 373, 1105-1110.	13.7	324
8	Selective serotonin reuptake inhibitors in pregnancy and congenital malformations: population based cohort study. BMJ: British Medical Journal, 2009, 339, b3569-b3569.	2.3	263
9	Long-term risk of dementia among people with traumatic brain injury in Denmark: a population-based observational cohort study. Lancet Psychiatry,the, 2018, 5, 424-431.	7.4	228
10	Risk for schizophrenia and schizophrenia-like psychosis among patients with epilepsy: population based cohort study. BMJ: British Medical Journal, 2005, 331, 23.	2.3	198
11	Mortality and life expectancy in persons with severe unipolar depression. Journal of Affective Disorders, 2016, 193, 203-207.	4.1	180
12	Health of children born to mothers who had preeclampsia: a population-based cohort study. American Journal of Obstetrics and Gynecology, 2009, 201, 269.e1-269.e10.	1.3	165
13	Validation of epilepsy diagnoses in the Danish National Hospital Register. Epilepsy Research, 2007, 75, 162-170.	1.6	163
14	Incidence and prevalence of epilepsy in Denmark. Epilepsy Research, 2007, 76, 60-65.	1.6	156
15	Attention-deficit/hyperactivity disorder in the offspring following prenatal maternal bereavement: a nationwide follow-up study in Denmark. European Child and Adolescent Psychiatry, 2010, 19, 747-753.	4.7	156
16	Effect of Depression and Diabetes Mellitus on the Risk for Dementia. JAMA Psychiatry, 2015, 72, 612.	11.0	145
17	MMR Vaccination and Febrile Seizures. JAMA - Journal of the American Medical Association, 2004, 292, 351.	7.4	144
18	Antidepressant exposure in pregnancy and risk of autism spectrum disorders. Clinical Epidemiology, 2013, 5, 449.	3.0	140

#	ARTICLE	IF	CITATIONS
19	Common variants associated with general and MMR vaccine-related febrile seizures. <i>Nature Genetics</i> , 2014, 46, 1274-1282.	21.4	128
20	Prenatal Stress Exposure Related to Maternal Bereavement and Risk of Childhood Overweight. <i>PLoS ONE</i> , 2010, 5, e11896.	2.5	120
21	The Association Between Perceived Stress and Mortality Among People With Multimorbidity: A Prospective Population-Based Cohort Study. <i>American Journal of Epidemiology</i> , 2016, 184, 199-210.	3.4	119
22	The construct validity of the Perceived Stress Scale. <i>Journal of Psychosomatic Research</i> , 2016, 84, 22-30.	2.6	114
23	Long-term Risk of Dementia in Persons With Schizophrenia. <i>JAMA Psychiatry</i> , 2015, 72, 1095.	11.0	107
24	Paternal age and congenital malformations. <i>Human Reproduction</i> , 2005, 20, 3173-3177.	0.9	106
25	The Long-Term Risk of Epilepsy after Febrile Seizures in Susceptible Subgroups. <i>American Journal of Epidemiology</i> , 2007, 165, 911-918.	3.4	106
26	Incidence of Suicide Among Persons Who Had a Parent Who Died During Their Childhood. <i>JAMA Psychiatry</i> , 2015, 72, 1227.	11.0	102
27	Myocardial Infarction and Risk of Suicide. <i>Circulation</i> , 2010, 122, 2388-2393.	1.6	98
28	Mortality after Parental Death in Childhood: A Nationwide Cohort Study from Three Nordic Countries. <i>PLoS Medicine</i> , 2014, 11, e1001679.	8.4	98
29	Association of Psychiatric and Neurologic Comorbidity With Mortality Among Persons With Autism Spectrum Disorder in a Danish Population. <i>JAMA Pediatrics</i> , 2016, 170, 243.	6.2	95
30	A Nationwide Study on the Risk of Autism After Prenatal Stress Exposure to Maternal Bereavement. <i>Pediatrics</i> , 2009, 123, 1102-1107.	2.1	88
31	Prenatal Antidepressant Exposure and Risk of Spontaneous Abortion – A Population-Based Study. <i>PLoS ONE</i> , 2013, 8, e72095.	2.5	87
32	Association of Streptococcal Throat Infection With Mental Disorders. <i>JAMA Psychiatry</i> , 2017, 74, 740.	11.0	83
33	Risk of Febrile Seizures and Epilepsy After Vaccination With Diphtheria, Tetanus, Acellular Pertussis, Inactivated Poliovirus, and Haemophilus Influenzae Type b. <i>JAMA - Journal of the American Medical Association</i> , 2012, 307, 823-31.	7.4	82
34	A new questionnaire to identify bodily distress in primary care: The “BDS checklist”™. <i>Journal of Psychosomatic Research</i> , 2015, 78, 536-545.	2.6	78
35	Death in children with febrile seizures: a population-based cohort study. <i>Lancet</i> , The, 2008, 372, 457-463.	13.7	72
36	The Danish National Hospital Register is a valuable study base for epidemiologic research in febrile seizures. <i>Journal of Clinical Epidemiology</i> , 2006, 59, 61-66.	5.0	71

#	ARTICLE	IF	CITATIONS
37	Paternal age and mortality in children. <i>European Journal of Epidemiology</i> , 2008, 23, 443-447.	5.7	69
38	Bodily distress syndrome: A new diagnosis for functional disorders in primary care?. <i>BMC Family Practice</i> , 2015, 16, 180.	2.9	68
39	Preeclampsia and Risk for Epilepsy in Offspring. <i>Pediatrics</i> , 2008, 122, 1072-1078.	2.1	65
40	Apgar Scores and Long-Term Risk of Epilepsy. <i>Epidemiology</i> , 2006, 17, 296-301.	2.7	64
41	Gestational Age, Birth Weight, Intrauterine Growth, and the Risk of Epilepsy. <i>American Journal of Epidemiology</i> , 2007, 167, 262-270.	3.4	64
42	Paternal Age and Epilepsy in the Offspring. <i>European Journal of Epidemiology</i> , 2005, 20, 1003-1005.	5.7	61
43	Register-based studies on febrile seizures in Denmark. <i>Brain and Development</i> , 2009, 31, 372-377.	1.1	60
44	Prevalence of depression, quality of life and antidepressant treatment in the Danish General Suburban Population Study. <i>Nordic Journal of Psychiatry</i> , 2014, 68, 507-512.	1.3	60
45	Fetal Growth and Preterm Birth in Children Exposed to Maternal or Paternal Rheumatoid Arthritis: A Nationwide Cohort Study. <i>Arthritis and Rheumatology</i> , 2014, 66, 3265-3273.	5.6	60
46	Use of phenytoin, phenobarbital, or diazepam during pregnancy and risk of congenital abnormalities: a case-control study. <i>Pharmacoepidemiology and Drug Safety</i> , 2007, 16, 181-188.	1.9	56
47	Serious Mental Illness and Risk for Hospitalizations and Rehospitalizations for Ambulatory Care-sensitive Conditions in Denmark. <i>Medical Care</i> , 2016, 54, 90-97.	2.4	56
48	Birth outcomes after prenatal exposure to antiepileptic drugs: A population-based study. <i>Epilepsia</i> , 2014, 55, 1714-1721.	5.1	55
49	Prenatal Exposure to Maternal Infections and Epilepsy in Childhood: A Population-Based Cohort Study. <i>Pediatrics</i> , 2008, 121, e1100-e1107.	2.1	54
50	Risk Factors for Febrile Convulsions. <i>Epidemiology</i> , 2002, 13, 282-287.	2.7	52
51	Problems and challenges in relation to the treatment of patients with multimorbidity: General practitioners' views and attitudes. <i>Scandinavian Journal of Primary Health Care</i> , 2015, 33, 121-126.	1.5	52
52	Prenatal Exposure to Cigarettes, Alcohol, and Coffee and the Risk for Febrile Seizures. <i>Pediatrics</i> , 2005, 116, 1089-1094.	2.1	51
53	Paternal Age and Preterm Birth. <i>Epidemiology</i> , 2005, 16, 259-262.	2.7	51
54	Risk of suicide, deliberate self-harm and psychiatric illness after the loss of a close relative: A nationwide cohort study. <i>World Psychiatry</i> , 2017, 16, 193-199.	10.4	50

#	ARTICLE	IF	CITATIONS
55	Impact of socioeconomic position and distance on mental health care utilization: a nationwide Danish follow-up study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017, 52, 1405-1413.	3.1	48
56	Use of antiepileptic drugs during pregnancy and risk of spontaneous abortion and stillbirth: population based cohort study. <i>BMJ, The</i> , 2014, 349, g5159-g5159.	6.0	47
57	Prenatal Exposure to Bereavement and Type-2 Diabetes: A Danish Longitudinal Population Based Study. <i>PLoS ONE</i> , 2012, 7, e43508.	2.5	46
58	Chronic care management in Danish general practice - a cross-sectional study of workload and multimorbidity. <i>BMC Family Practice</i> , 2012, 13, 52.	2.9	45
59	Long-Term Health Outcomes in Children Born to Mothers with Diabetes: A Population-Based Cohort Study. <i>PLoS ONE</i> , 2012, 7, e36727.	2.5	45
60	Early Life Disease Programming during the Preconception and Prenatal Period: Making the Link between Stressful Life Events and Type-1 Diabetes. <i>PLoS ONE</i> , 2010, 5, e11523.	2.5	42
61	Cohort Profile: The Nordic Perinatal Bereavement Cohort. <i>International Journal of Epidemiology</i> , 2011, 40, 1161-1167.	1.9	42
62	Psychological distress, cardiovascular complications and mortality among people with screen-detected type 2 diabetes: follow-up of the ADDITION-Denmark trial. <i>Diabetologia</i> , 2014, 57, 710-717.	6.3	39
63	Adverse pregnancy outcomes after exposure to methylphenidate or atomoxetine during pregnancy. <i>Clinical Epidemiology</i> , 2015, 7, 139.	3.0	39
64	Patient characteristics and frequency of bodily distress syndrome in primary care: a cross-sectional study. <i>British Journal of General Practice</i> , 2015, 65, e617-e623.	1.4	38
65	Psychosocial Stress Related to the Loss of a Close Relative the Year Before or During Pregnancy and Risk of Preeclampsia. <i>Hypertension</i> , 2013, 62, 183-189.	2.7	37
66	Association of Mortality With the Death of a Sibling in Childhood. <i>JAMA Pediatrics</i> , 2017, 171, 538.	6.2	37
67	Depression, antidepressants, and the risk of non-valvular atrial fibrillation: A nationwide Danish matched cohort study. <i>European Journal of Preventive Cardiology</i> , 2019, 26, 187-195.	1.8	37
68	Binge Drinking During Pregnancy and Risk of Seizures in Childhood: A Study Based on the Danish National Birth Cohort. <i>American Journal of Epidemiology</i> , 2008, 169, 313-322.	3.4	36
69	Febrile seizures and risk of schizophrenia. <i>Schizophrenia Research</i> , 2005, 73, 343-349.	2.0	35
70	Thirty-Day Mortality After Infection Among Persons With Severe Mental Illness: A Population-Based Cohort Study in Denmark. <i>American Journal of Psychiatry</i> , 2015, 172, 776-783.	7.2	35
71	Pregnancy outcomes among female hairdressers who participated in the Danish National Birth Cohort. <i>Scandinavian Journal of Work, Environment and Health</i> , 2006, 32, 61-66.	3.4	34
72	Depressive Symptoms and Risk of New Cardiovascular Events or Death in Patients with Myocardial Infarction: A Population-Based Longitudinal Study Examining Health Behaviors and Health Care Interventions. <i>PLoS ONE</i> , 2013, 8, e74393.	2.5	33

#	ARTICLE	IF	CITATIONS
73	Long-term mortality in children and young adults with epilepsy – A population-based cohort study. <i>Epilepsy Research</i> , 2015, 114, 81-88.	1.6	33
74	Low Apgar Scores and Risk of Childhood Attention Deficit Hyperactivity Disorder. <i>Journal of Pediatrics</i> , 2011, 158, 775-779.	1.8	32
75	Maternal Bereavement During Pregnancy and the Risk of Stillbirth: A Nationwide Cohort Study in Sweden. <i>American Journal of Epidemiology</i> , 2013, 177, 219-227.	3.4	32
76	Prenatal stress and epilepsy in later life: A nationwide follow-up study in Denmark. <i>Epilepsy Research</i> , 2008, 81, 52-57.	1.6	31
77	The prognosis of bodily distress syndrome: a cohort study in primary care. <i>General Hospital Psychiatry</i> , 2015, 37, 560-566.	2.4	31
78	Stress and epilepsy: A population-based cohort study of epilepsy in parents who lost a child. <i>Epilepsy and Behavior</i> , 2007, 11, 324-328.	1.7	30
79	Prenatal Stress and Cerebral Palsy: A Nationwide Cohort Study in Denmark. <i>Psychosomatic Medicine</i> , 2009, 71, 615-618.	2.0	30
80	Epilepsy and febrile seizures in children of treated and untreated subfertile couples. <i>Human Reproduction</i> , 2007, 22, 215-220.	0.9	29
81	Association between perceived stress, multimorbidity and primary care health services: a Danish population-based cohort study. <i>BMJ Open</i> , 2018, 8, e018323.	1.9	29
82	Prenatal Maternal Bereavement and Congenital Heart Defects in Offspring: A Registry-Based Study. <i>Pediatrics</i> , 2013, 131, e1225-e1230.	2.1	28
83	Post-Myocardial Infarction Anxiety or Depressive Symptoms and Risk of New Cardiovascular Events or Death. <i>Psychosomatic Medicine</i> , 2014, 76, 739-746.	2.0	28
84	Depression and risk of hospitalisations and rehospitalisations for ambulatory care-sensitive conditions in Denmark: a population-based cohort study. <i>BMJ Open</i> , 2015, 5, e009878.	1.9	27
85	Variation in general practice prostate-specific antigen testing and prostate cancer outcomes: An ecological study. <i>International Journal of Cancer</i> , 2015, 136, 435-442.	5.1	27
86	Mental health status and risk of new cardiovascular events or death in patients with myocardial infarction: a population-based cohort study. <i>BMJ Open</i> , 2013, 3, e003045.	1.9	26
87	Long-term risk of atrial fibrillation after the death of a partner. <i>Open Heart</i> , 2016, 3, e000367.	2.3	26
88	MMR Vaccination and Autism. <i>Drug Safety</i> , 2004, 27, 831-840.	3.2	25
89	The 5-minute Apgar score as a predictor of childhood cancer: a population-based cohort study in five million children. <i>BMJ Open</i> , 2012, 2, e001095.	1.9	25
90	Socioeconomic Position, Type 2 Diabetes and Long-Term Risk of Death. <i>PLoS ONE</i> , 2015, 10, e0124829.	2.5	24

#	ARTICLE	IF	CITATIONS
91	Risk of Fetal Death after Treatment with Antipsychotic Medications during Pregnancy. PLoS ONE, 2015, 10, e0132280.	2.5	23
92	Assessing fetal growth impairments based on family data as a tool for identifying high-risk babies. An example with neonatal mortality. BMC Pregnancy and Childbirth, 2007, 7, 28.	2.4	22
93	Breastfeeding and Risk of Epilepsy in Childhood: A Birth Cohort Study. Journal of Pediatrics, 2011, 158, 924-929.	1.8	22
94	Diseases in children born to mothers with preeclampsia: a population-based sibling cohort study. American Journal of Obstetrics and Gynecology, 2011, 204, 157.e1-157.e5.	1.3	22
95	Rehabilitation status three months after first-time myocardial infarction. Scandinavian Journal of Primary Health Care, 2011, 29, 210-215.	1.5	22
96	Ten-Year Mortality after a Breast Cancer Diagnosis in Women with Severe Mental Illness: A Danish Population-Based Cohort Study. PLoS ONE, 2016, 11, e0158013.	2.5	22
97	Risk of epilepsy and autism in full and half siblings – A population-based cohort study. Epilepsia, 2016, 57, 2011-2018.	5.1	21
98	Risk of diabetic complications and subsequent mortality among individuals with schizophrenia and diabetes - a population-based register study. Schizophrenia Research, 2020, 218, 99-106.	2.0	21
99	Prenatal exposure to elevated maternal body temperature and risk of epilepsy in childhood: a population-based pregnancy cohort study. Paediatric and Perinatal Epidemiology, 2011, 25, 53-59.	1.7	20
100	Screening for depression in patients with myocardial infarction by general practitioners. European Journal of Preventive Cardiology, 2013, 20, 800-806.	1.8	19
101	Chronic-disease patients and their use of out-of-hours primary health care: a cross-sectional study. BMC Family Practice, 2014, 15, 114.	2.9	18
102	Prevaccination epidemiology of herpes zoster in Denmark: Quantification of occurrence and risk factors. Vaccine, 2017, 35, 5589-5596.	3.8	18
103	Frequency of health care utilization in the year prior to completed suicide: A Danish nationwide matched comparative study. PLoS ONE, 2019, 14, e0214605.	2.5	18
104	The association of unipolar depression with thirty-day mortality after hospitalization for infection: A population-based cohort study in Denmark. Journal of Psychosomatic Research, 2016, 89, 32-38.	2.6	17
105	Cancer Mortality in People Treated with Antidepressants before Cancer Diagnosis: A Population Based Cohort Study. PLoS ONE, 2015, 10, e0138134.	2.5	17
106	Psychiatric disorders following fetal death: a population-based cohort study. BMJ Open, 2014, 4, e005187-e005187.	1.9	16
107	Partner Bereavement and Risk of Herpes Zoster: Results from Two Population-Based Case-Control Studies in Denmark and the United Kingdom. Clinical Infectious Diseases, 2017, 64, ciw840.	5.8	16
108	Attendance of routine childcare visits in primary care for children of mothers with depression: a nationwide population-based cohort study. British Journal of General Practice, 2018, 68, e97-e104.	1.4	16

#	ARTICLE	IF	CITATIONS
109	Paternal Age and Apgar Scores of Newborn Infants. <i>Epidemiology</i> , 2006, 17, 473-474.	2.7	15
110	Prenatal Stress and Risk of Febrile Seizures in Children: A Nationwide Longitudinal Study in Denmark. <i>Journal of Autism and Developmental Disorders</i> , 2009, 39, 1047-1052.	2.7	15
111	Algorithm linking patients and general practices in Denmark using the Danish National Health Service Register. <i>Clinical Epidemiology</i> , 2016, Volume 8, 273-283.	3.0	15
112	Prenatal exposure to antidepressants and risk of epilepsy in childhood. <i>Pharmacoepidemiology and Drug Safety</i> , 2016, 25, 1320-1330.	1.9	15
113	Perceived Stress, Multimorbidity, and Risk for Hospitalizations for Ambulatory Care-sensitive Conditions. <i>Medical Care</i> , 2017, 55, 131-139.	2.4	15
114	Gestational Age, Birth Weight, and Risk for Injuries in Childhood. <i>Epidemiology</i> , 2010, 21, 650-657.	2.7	14
115	Hospital-based herpes zoster diagnoses in Denmark: rate, patient characteristics, and all-cause mortality. <i>BMC Infectious Diseases</i> , 2016, 16, 99.	2.9	13
116	The construct validity of the Major Depression Inventory: A Rasch analysis of a self-rating scale in primary care. <i>Journal of Psychosomatic Research</i> , 2017, 97, 70-81.	2.6	13
117	Use of general practice before and after mild traumatic brain injury: a nationwide population-based cohort study in Denmark. <i>BMJ Open</i> , 2017, 7, e017735.	1.9	13
118	Psychiatric conditions and general practitioner attendance prior to HPV vaccination and the risk of referral to a specialized hospital setting because of suspected adverse events following HPV vaccination: a register-based, matched case–control study. <i>Clinical Epidemiology</i> , 2017, Volume 9, 465-473.	3.0	13
119	Apgar-score in children prenatally exposed to antiepileptic drugs: a population-based cohort study. <i>BMJ Open</i> , 2015, 5, e007425.	1.9	12
120	Does perceived stress increase the risk of atrial fibrillation? A population-based cohort study in Denmark. <i>American Heart Journal</i> , 2017, 188, 26-34.	2.7	12
121	The criterion validity of the web-based Major Depression Inventory when used on clinical suspicion of depression in primary care. <i>Clinical Epidemiology</i> , 2017, Volume 9, 355-365.	3.0	12
122	Mood Disorders and Risk of Herpes Zoster in 2 Population-Based Case-Control Studies in Denmark and the United Kingdom. <i>American Journal of Epidemiology</i> , 2018, 187, 1019-1028.	3.4	12
123	Bereavement in Early Life and Later Childhood Overweight. <i>Obesity Facts</i> , 2012, 5, 881-889.	3.4	11
124	Post-Stroke Mortality, Stroke Severity, and Preadmission Antipsychotic Medicine Use - A Population-Based Cohort Study. <i>PLoS ONE</i> , 2014, 9, e84103.	2.5	11
125	Use of primary health care prior to a postpartum psychiatric episode. <i>Scandinavian Journal of Primary Health Care</i> , 2015, 33, 127-133.	1.5	10
126	Socioeconomic position and cardiovascular risk factors among people with screen-detected Type 2 DM: Six-year follow-up of the ADDITION-Denmark trial. <i>Primary Care Diabetes</i> , 2014, 8, 322-329.	1.8	9

#	ARTICLE	IF	CITATIONS
127	Early treatment with talk therapy or antidepressants in severely bereaved people and risk of suicidal behavior and psychiatric illness: an instrumental variable analysis. <i>Clinical Epidemiology</i> , 2018, Volume 10, 1013-1026.	3.0	9
128	Socioeconomic position and contact to general practice among persons with diabetes. <i>Primary Care Diabetes</i> , 2012, 6, 313-318.	1.8	8
129	Daytime use of general practice and use of the Out-of-Hours Primary Care Service for patients with chronic disease: a cohort study. <i>BMC Family Practice</i> , 2014, 15, 156.	2.9	8
130	Increased utilisation of primary healthcare in persons exposed to severe stress in prenatal life: a national population-based study in Denmark. <i>BMJ Open</i> , 2015, 5, e005657-e005657.	1.9	8
131	Prenatal exposure to antiepileptic drugs and use of primary healthcare during childhood: a population-based cohort study in Denmark. <i>BMJ Open</i> , 2017, 7, e012836.	1.9	8
132	Selective serotonin reuptake inhibitors and risk of epilepsy after traumatic brain injury – A population based cohort study. <i>PLoS ONE</i> , 2019, 14, e0219137.	2.5	8
133	Glucocorticoid use and risk of suicide: a Danish population-based case-control study. <i>World Psychiatry</i> , 2021, 20, 142-143.	10.4	8
134	Maternal Use of Antibiotics and the Risk of Childhood Febrile Seizures: A Danish Population-Based Cohort. <i>PLoS ONE</i> , 2013, 8, e61148.	2.5	8
135	Healthcare Contacts after Myocardial Infarction According to Mental Health and Socioeconomic Position: A Population-Based Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0134557.	2.5	7
136	The Risk of Gestational Diabetes Mellitus Following Bereavement: A Cohort Study from Denmark and Sweden. <i>Paediatric and Perinatal Epidemiology</i> , 2015, 29, 271-280.	1.7	7
137	Maternal bereavement shortly before or during pregnancy and risk of postpartum psychotic illness: a population-based study from Denmark and Sweden. <i>Clinical Epidemiology</i> , 2019, Volume 11, 285-298.	3.0	7
138	Intake of marine n-3 fatty acids during pregnancy and risk for epilepsy in the offspring: A population-based cohort study. <i>Epilepsy Research</i> , 2010, 91, 267-272.	1.6	6
139	Premature mortality in persons with epilepsy and schizophrenia: A population-based nationwide cohort study. <i>Epilepsia</i> , 2019, 60, 1200-1208.	5.1	6
140	Risk of substance use disorders following prenatal or postnatal exposure to bereavement. <i>Drug and Alcohol Dependence</i> , 2013, 132, 277-282.	3.2	5
141	The 30-day prognosis of chronic-disease patients after contact with the out-of-hours service in primary healthcare. <i>Scandinavian Journal of Primary Health Care</i> , 2014, 32, 208-216.	1.5	4
142	Use of primary healthcare for persons with epilepsy. <i>Epilepsy and Behavior</i> , 2018, 80, 285-290.	1.7	2
143	Health care utilization in general practice after HPV vaccination – A Danish nationwide register-based cohort study. <i>PLoS ONE</i> , 2017, 12, e0184658.	2.5	2
144	Laszlo et al. Respond to "Maternal Stress and Stillbirth". <i>American Journal of Epidemiology</i> , 2013, 177, 230-231.	3.4	1

#	ARTICLE	IF	CITATIONS
145	O5.7. RISK OF DIABETIC COMPLICATIONS AND SUBSEQUENT MORTALITY AMONG INDIVIDUALS WITH SCHIZOPHRENIA AND DIABETES MELLITUS: A NATIONWIDE POPULATION-BASED REGISTER STUDY. Schizophrenia Bulletin, 2018, 44, S88-S89.	4.3	1
146	Traumatic brain injury and dementia – Authors' reply. Lancet Psychiatry, the, 2018, 5, 783.	7.4	0