

# Berhane T Weldegergis

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1839382/publications.pdf>

Version: 2024-02-01

37  
papers

2,137  
citations

201674

27  
h-index

330143

37  
g-index

37  
all docs

37  
docs citations

37  
times ranked

2661  
citing authors

#	ARTICLE	IF	CITATIONS
1	Symbiotic polydnavirus and venom reveal parasitoid to its hyperparasitoids. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, 5205-5210.	7.1	54
2	Covariation and phenotypic integration in chemical communication displays: biosynthetic constraints and eco-evolutionary implications. <i>New Phytologist</i> , 2018, 220, 739-749.	7.3	101
3	Do apes smell like humans? The role of skin bacteria and volatiles of primates in mosquito host selection. <i>Journal of Experimental Biology</i> , 2018, 221, .	1.7	24
4	Does Aphid Infestation Interfere with Indirect Plant Defense against Lepidopteran Caterpillars in Wild Cabbage?. <i>Journal of Chemical Ecology</i> , 2017, 43, 493-505.	1.8	12
5	Qualitative and Quantitative Differences in Herbivore-Induced Plant Volatile Blends from Tomato Plants Infested by Either <i>Tuta absoluta</i> or <i>Bemisia tabaci</i> . <i>Journal of Chemical Ecology</i> , 2017, 43, 53-65.	1.8	63
6	Response of a Predatory ant to Volatiles Emitted by Aphid- and Caterpillar-Infested Cucumber and Potato Plants. <i>Journal of Chemical Ecology</i> , 2017, 43, 1007-1022.	1.8	19
7	Terpenoid biosynthesis in <i>Arabidopsis</i> attacked by caterpillars and aphids: effects of aphid density on the attraction of a caterpillar parasitoid. <i>Oecologia</i> , 2017, 185, 699-712.	2.0	10
8	Symbionts protect aphids from parasitic wasps by attenuating herbivore-induced plant volatiles. <i>Nature Communications</i> , 2017, 8, 1860.	12.8	96
9	Integrating Insect Life History and Food Plant Phenology: Flexible Maternal Choice Is Adaptive. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1263.	4.1	6
10	Compatible and incompatible pathogen-plant interactions differentially affect plant volatile emissions and the attraction of parasitoid wasps. <i>Functional Ecology</i> , 2016, 30, 1779-1789.	3.6	31
11	Attractiveness of volatiles from different body parts to the malaria mosquito <i>Anopheles coluzzii</i> is affected by deodorant compounds. <i>Scientific Reports</i> , 2016, 6, 27141.	3.3	43
12	Trading direct for indirect defense? Phytochrome B inactivation in tomato attenuates direct anti-herbivore defenses whilst enhancing volatile-mediated attraction of predators. <i>New Phytologist</i> , 2016, 212, 1057-1071.	7.3	59
13	Volatile-mediated foraging behaviour of three parasitoid species under conditions of dual insect herbivore attack. <i>Animal Behaviour</i> , 2016, 111, 197-206.	1.9	50
14	Attraction of egg-killing parasitoids toward induced plant volatiles in a multi-herbivore context. <i>Oecologia</i> , 2015, 179, 163-174.	2.0	45
15	Altered Volatile Profile Associated with Precopulatory Mate Guarding Attracts Spider Mite Males. <i>Journal of Chemical Ecology</i> , 2015, 41, 187-193.	1.8	9
16	To be in time: egg deposition enhances plant-mediated detection of young caterpillars by parasitoids. <i>Oecologia</i> , 2015, 177, 477-486.	2.0	29
17	Rhizobacterial colonization of roots modulates plant volatile emission and enhances the attraction of a parasitoid wasp to host-infested plants. <i>Oecologia</i> , 2015, 178, 1169-1180.	2.0	83
18	Parasitism overrides herbivore identity allowing hyperparasitoids to locate their parasitoid host using herbivore-induced plant volatiles. <i>Molecular Ecology</i> , 2015, 24, 2886-2899.	3.9	40

#	ARTICLE	IF	CITATIONS
19	Drought stress affects plant metabolites and herbivore preference but not host location by its parasitoids. <i>Oecologia</i> , 2015, 177, 701-713.	2.0	75
20	Understanding the Long-Lasting Attraction of Malaria Mosquitoes to Odor Baits. <i>PLoS ONE</i> , 2015, 10, e0121533.	2.5	17
21	Virulence Factors of Geminivirus Interact with MYC2 to Subvert Plant Resistance and Promote Vector Performance. <i>Plant Cell</i> , 2014, 26, 4991-5008.	6.6	224
22	Caterpillar-induced plant volatiles remain a reliable signal for foraging wasps during dual attack with a plant pathogen or non-host insect herbivore. <i>Plant, Cell and Environment</i> , 2014, 37, 1924-1935.	5.7	66
23	Synergism in the effect of prior jasmonic acid application on herbivore-induced volatile emission by Lima bean plants: transcription of a monoterpene synthase gene and volatile emission. <i>Journal of Experimental Botany</i> , 2014, 65, 4821-4831.	4.8	29
24	Effect of Sequential Induction by <i>Mamestra brassicae</i> L. and <i>Tetranychus urticae</i> Koch on Lima Bean Plant Indirect Defense. <i>Journal of Chemical Ecology</i> , 2014, 40, 977-985.	1.8	8
25	Body Odors of Parasitized Caterpillars Give Away the Presence of Parasitoid Larvae to Their Primary Hyperparasitoid Enemies. <i>Journal of Chemical Ecology</i> , 2014, 40, 986-995.	1.8	22
26	Canopy light cues affect emission of constitutive and methyl jasmonate-induced volatile organic compounds in <i>Scirpus thalassia</i> . <i>New Phytologist</i> , 2013, 200, 861-874.	7.3	78
27	Non-pathogenic rhizobacteria interfere with the attraction of parasitoids to aphid-induced plant volatiles via jasmonic acid signalling. <i>Plant, Cell and Environment</i> , 2013, 36, 393-404.	5.7	110
28	Genetic engineering of plant volatile terpenoids: effects on a herbivore, a predator and a parasitoid. <i>Pest Management Science</i> , 2013, 69, 302-311.	3.4	43
29	Hyperparasitoids Use Herbivore-Induced Plant Volatiles to Locate Their Parasitoid Host. <i>PLoS Biology</i> , 2012, 10, e1001435.	5.6	168
30	Plant Volatiles Induced by Herbivore Egg Deposition Affect Insects of Different Trophic Levels. <i>PLoS ONE</i> , 2012, 7, e43607.	2.5	152
31	Neonates know better than their mothers when selecting a host plant. <i>Oikos</i> , 2012, 121, 1923-1934.	2.7	46
32	Herbivore-Mediated Effects of Glucosinolates on Different Natural Enemies of a Specialist Aphid. <i>Journal of Chemical Ecology</i> , 2012, 38, 100-115.	1.8	77
33	Solid phase extraction in combination with comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry for the detailed investigation of volatiles in South African red wines. <i>Analytica Chimica Acta</i> , 2011, 701, 98-111.	5.4	68
34	Chemometric investigation of the volatile content of young South African wines. <i>Food Chemistry</i> , 2011, 128, 1100-1109.	8.2	33
35	Characterisation of volatile components of Pinotage wines using comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry (GC-TOFMS). <i>Food Chemistry</i> , 2011, 129, 188-199.	8.2	81
36	Analysis of Volatiles in Pinotage Wines by Stir Bar Sorptive Extraction and Chemometric Profiling. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 10225-10236.	5.2	31

#	ARTICLE	IF	CITATIONS
37	Application of a Headspace Sorptive Extraction Method for the Analysis of Volatile Components in South African Wines. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 8696-8702.	5.2	35