

Åystein Hetlevik

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1874619/publications.pdf>

Version: 2024-02-01

29
papers

432
citations

840776

11
h-index

839539

18
g-index

30
all docs

30
docs citations

30
times ranked

432
citing authors

#	ARTICLE	IF	CITATIONS
1	Continuity in general practice as predictor of mortality, acute hospitalisation, and use of out-of-hours care: a registry-based observational study in Norway. <i>British Journal of General Practice</i> , 2022, 72, e84-e90.	1.4	106
2	Continuity of care for patients with chronic disease: a registry-based observational study from Norway. <i>Family Practice</i> , 2022, 39, 570-578.	1.9	7
3	Reasons for acute referrals to hospital from general practitioners and out-of-hours doctors in Norway: a registry-based observational study. <i>BMC Health Services Research</i> , 2022, 22, 78.	2.2	2
4	Factors associated with self-rated health in primary care in the South-Western health zone of Malawi. , 2022, 23, 88.		3
5	Patient experiences with depression care in general practice: a qualitative questionnaire study. <i>Scandinavian Journal of Primary Health Care</i> , 2022, 40, 253-260.	1.5	5
6	Construction and validation of a morbidity index based on the International Classification of Primary Care. <i>Scandinavian Journal of Primary Health Care</i> , 2022, 40, 305-312.	1.5	8
7	Variation in general practitioners' depression care following certification of sickness absence: a registry-based cohort study. <i>Family Practice</i> , 2021, 38, 238-245.	1.9	5
8	Continuity of care, measurement and association with hospital admission and mortality: a registry-based longitudinal cohort study. <i>BMJ Open</i> , 2021, 11, e051958.	1.9	6
9	Self-rated health in adolescence as a predictor of "multi-illness" in early adulthood: A prospective registry-based Norwegian HUNT study. <i>SSM - Population Health</i> , 2020, 11, 100604.	2.7	12
10	General practitioners' and out-of-hours doctors' role as gatekeeper in emergency admissions to somatic hospitals in Norway: registry-based observational study. <i>BMC Health Services Research</i> , 2019, 19, 568.	2.2	23
11	Patient-reported depression treatment and future treatment preferences: an observational study in general practice. <i>Family Practice</i> , 2019, 36, 771-777.	1.9	10
12	The effect of team collaboration and continuity of care on health and disability among rehabilitation patients: a longitudinal survey-based study from western Norway. <i>Quality of Life Research</i> , 2019, 28, 2773-2785.	3.1	8
13	Adolescent self-rated health predicts general practice attendance in adulthood: Results from the Young-HUNT1 survey. <i>Scandinavian Journal of Public Health</i> , 2019, 47, 37-44.	2.3	23
14	Adolescents consulting general practitioners for psychological problems—a nationwide, register-based study in Norway. <i>Family Practice</i> , 2019, 36, 77-83.	1.9	6
15	GP-diagnosed internalizing and externalizing problems and dropout from secondary school: a cross-sectional study. <i>European Journal of Public Health</i> , 2018, 28, 474-479.	0.3	12
16	Quality of primary care from patients' perspective: a cross sectional study of outpatients' experience in public health facilities in rural Malawi. <i>BMC Health Services Research</i> , 2018, 18, 872.	2.2	13
17	Development and validation of a Malawian version of the primary care assessment tool. <i>BMC Family Practice</i> , 2018, 19, 63.	2.9	17
18	GP follow-up after first diagnosing psychological problems among adolescents: a nationwide register-based study. <i>British Journal of General Practice</i> , 2018, 68, e794-e802.	1.4	2

#	ARTICLE	IF	CITATIONS
19	Relational coordination in interprofessional teams and its effect on patient-reported benefit and continuity of care: a prospective cohort study from rehabilitation centres in Western Norway. <i>BMC Health Services Research</i> , 2018, 18, 719.	2.2	24
20	Communication and Relational Ties in Inter-Professional Teams in Norwegian Specialized Health Care: A Multicentre Study of Relational Coordination. <i>International Journal of Integrated Care</i> , 2018, 18, 9.	0.2	14
21	Patient perspectives on continuity of care: adaption and preliminary psychometric assessment of a Norwegian version of the Nijmegen Continuity Questionnaire (NCQ-N). <i>BMC Health Services Research</i> , 2017, 17, 760.	2.2	13
22	GP consultations for common mental disorders and subsequent sickness certification: register-based study of the employed population in Norway. <i>Family Practice</i> , 2016, 33, 656-662.	1.9	13
23	GP utilisation by education level among adults with COPD or asthma: a cross-sectional register-based study. <i>Npj Primary Care Respiratory Medicine</i> , 2016, 26, 16027.	2.6	7
24	Point-of-care testing with CRP in primary care: a registry-based observational study from Norway. <i>BMC Family Practice</i> , 2015, 16, 170.	2.9	12
25	Use of GP services by patients with schizophrenia: a national cross-sectional register-based study. <i>BMC Health Services Research</i> , 2015, 15, 66.	2.2	19
26	Personal continuity of care in Norwegian general practice: A national cross-sectional study. <i>Scandinavian Journal of Primary Health Care</i> , 2012, 30, 214-221.	1.5	24
27	Does socioeconomic status of list populations affect GP practice? A register-based study of 2201 Norwegian GPs. <i>European Journal of General Practice</i> , 2012, 18, 212-218.	2.0	2
28	Norwegian GPs' participation in multidisciplinary meetings: A register-based study from 2007. <i>BMC Health Services Research</i> , 2010, 10, 309.	2.2	6
29	Young people and their GP: a register-based study of 1717 Norwegian GPs. <i>Family Practice</i> , 2010, 27, 3-8.	1.9	22