

Robert S Hodges

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1883905/publications.pdf>

Version: 2024-02-01

158
papers

12,506
citations

16451

64
h-index

25787

108
g-index

160
all docs

160
docs citations

160
times ranked

9616
citing authors

#	ARTICLE	IF	CITATIONS
1	Overcoming prostate cancer drug resistance with a novel organosilicon small molecule. <i>Neoplasia</i> , 2021, 23, 1261-1274.	5.3	4
2	Deuteration of nonexchangeable protons on proteins affects their thermal stability, side-chain dynamics, and hydrophobicity. <i>Protein Science</i> , 2020, 29, 1641-1654.	7.6	21
3	De Novo Designed Amphipathic α -Helical Antimicrobial Peptides Incorporating Dab and Dap Residues on the Polar Face To Treat the Gram-Negative Pathogen, <i>Acinetobacter baumannii</i> . <i>Journal of Medicinal Chemistry</i> , 2019, 62, 3354-3366.	6.4	43
4	Design of Novel Amphipathic α -Helical Antimicrobial Peptides with No Toxicity as Therapeutics against the Antibiotic-Resistant Gram-Negative Bacterial Pathogen,. <i>Journal of Medicinal Chemistry and Drug Design</i> , 2019, 2, .	0.3	0
5	Role of positively charged residues on the polar and non-polar faces of amphipathic α -helical antimicrobial peptides on specificity and selectivity for Gram-negative pathogens. <i>Chemical Biology and Drug Design</i> , 2018, 91, 75-92.	3.2	27
6	Separation of highly charged (+5 to +10) amphipathic α -helical peptide standards by cation-exchange and reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2018, 1574, 60-70.	3.7	3
7	Two Complementary Approaches for the Controlled Release of Biomolecules Immobilized via Coiled-Coil Interactions: Peptide Core Mutations and Multivalent Presentation. <i>Biomacromolecules</i> , 2017, 18, 965-975.	5.4	16
8	Optimized purification of a fusion protein by reversed-phase high performance liquid chromatography informed by the linear solvent strength model. <i>Journal of Chromatography A</i> , 2017, 1521, 44-52.	3.7	1
9	ProP-ProP and ProP-phospholipid interactions determine the subcellular distribution of osmosensing transporter ProP in <i>Escherichia coli</i> . <i>Molecular Microbiology</i> , 2017, 103, 469-482.	2.5	13
10	Platform technology to generate broadly cross-reactive antibodies to α -helical epitopes in hemagglutinin proteins from influenza A viruses. <i>Biopolymers</i> , 2016, 106, 144-159.	2.4	10
11	Separation of Peptides on HALO Micron Particles. <i>Current Protocols in Protein Science</i> , 2016, 85, 11.6.1-11.6.16.	2.8	4
12	Transmission of Stability Information through the N-domain of Tropomyosin Is Interrupted by a Stabilizing Mutation (A109L) in the Hydrophobic Core of the Stability Control Region (Residues 97-118). <i>Journal of Biological Chemistry</i> , 2014, 289, 4356-4366.	3.4	8
13	Specificity Determinants Improve Therapeutic Indices of Two Antimicrobial Peptides Piscidin 1 and Dermaseptin S4 Against the Gram-negative Pathogens <i>Acinetobacter baumannii</i> and <i>Pseudomonas aeruginosa</i> . <i>Pharmaceuticals</i> , 2014, 7, 366-391.	3.8	52
14	Structure-activity relationships of the antimicrobial peptide gramicidin S and its analogs: Aqueous solubility, self-association, conformation, antimicrobial activity and interaction with model lipid membranes. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2014, 1838, 1420-1429.	2.6	29
15	Bradykinin antagonists and thiazolidinone derivatives as new potential anti-cancer compounds. <i>Bioorganic and Medicinal Chemistry</i> , 2014, 22, 3815-3823.	3.0	27
16	An improved approach to hydrophilic interaction chromatography of peptides: Salt gradients in the presence of high isocratic acetonitrile concentrations. <i>Journal of Chromatography A</i> , 2013, 1277, 15-25.	3.7	16
17	Antiadhesin Peptide-Based Vaccines for <i>Pseudomonas aeruginosa</i> . , 2013, , 563-570.		2
18	Strategies for Designing Peptide Immunogens To Elicit α -Helical Conformation-Specific Antibodies Reactive with Native Proteins. <i>ACS Symposium Series</i> , 2012, , 93-136.	0.5	4

#	ARTICLE	IF	CITATIONS
19	Design of peptide standards with the same composition and minimal sequence variation to monitor performance/selectivity of reversed-phase matrices. <i>Journal of Chromatography A</i> , 2012, 1230, 30-40.	3.7	14
20	Anti-Tuberculosis Activity of α -Helical Antimicrobial Peptides: De Novo Designed L- and D-Enantiomers Versus L- and D-LL37. <i>Protein and Peptide Letters</i> , 2011, 18, 241-252.	0.9	55
21	Rational Design of α -Helical Antimicrobial Peptides to Target Gram-negative Pathogens, <i>Acinetobacter baumannii</i> and <i>Pseudomonas aeruginosa</i> : Utilization of Charge, Specificity Determinants, Total Hydrophobicity, Hydrophobe Type and Location as Design Parameters to Improve the Therapeutic Ratio. <i>Chemical Biology and Drug Design</i> , 2011, 77, 225-240.	3.2	100
22	Specificity determinants dramatically reduce hemolytic activity in amphipathic α -helical antimicrobial peptides: antimicrobial activity against Gram-negative pathogens, <i>Acinetobacter baumannii</i> and <i>Pseudomonas aeruginosa</i> . , 2011, , .		0
23	Reversed-phase HPLC of peptides: Assessing column and solvent selectivity on standard, polar-embedded and polar endcapped columns. <i>Journal of Separation Science</i> , 2010, 33, 3005-3021.	2.5	25
24	Research Article: Synthetic Peptide Vaccine Development: Designing Dual Epitopes into a Single Pilin Peptide Immunogen Generates Antibody Cross-reactivity between Two Strains of <i>Pseudomonas aeruginosa</i> . <i>Chemical Biology and Drug Design</i> , 2010, 76, 293-304.	3.2	11
25	Critical interactions in the stability control region of tropomyosin. <i>Journal of Structural Biology</i> , 2010, 170, 294-306.	2.8	9
26	Intrinsic amino acid side-chain hydrophilicity/hydrophobicity coefficients determined by reversed-phase high-performance liquid chromatography of model peptides: Comparison with other hydrophilicity/hydrophobicity scales. <i>Biopolymers</i> , 2009, 92, 573-595.	2.4	114
27	Advantages of a Synthetic Peptide Immunogen Over a Protein Immunogen in the Development of an Anti-Pilus Vaccine for <i>Pseudomonas aeruginosa</i> . <i>Chemical Biology and Drug Design</i> , 2009, 74, 33-42.	3.2	47
28	Antifreeze protein from shorthorn sculpin: Identification of the ice-binding surface. <i>Protein Science</i> , 2009, 10, 2566-2576.	7.6	53
29	Identification of a Unique Stability Control Region that Controls Protein Stability of Tropomyosin: A Two-stranded α -Helical Coiled-coil. <i>Journal of Molecular Biology</i> , 2009, 392, 747-762.	4.2	25
30	Mixed-mode hydrophilic interaction/cation-exchange chromatography: Separation of complex mixtures of peptides of varying charge and hydrophobicity. <i>Journal of Separation Science</i> , 2008, 31, 1573-1584.	2.5	42
31	Mixed-mode hydrophilic interaction/cation-exchange chromatography (HILIC/CEX) of peptides and proteins. <i>Journal of Separation Science</i> , 2008, 31, 2754-2773.	2.5	74
32	Effects of net charge and the number of positively charged residues on the biological activity of amphipathic α -helical cationic antimicrobial peptides. <i>Biopolymers</i> , 2008, 90, 369-383.	2.4	390
33	Effects of Hydrophobicity on the Antifungal Activity of α -Helical Antimicrobial Peptides. <i>Chemical Biology and Drug Design</i> , 2008, 72, 483-495.	3.2	73
34	Animal Protection and Structural Studies of a Consensus Sequence Vaccine Targeting the Receptor Binding Domain of the Type IV Pilus of <i>Pseudomonas aeruginosa</i> . <i>Journal of Molecular Biology</i> , 2007, 374, 426-442.	4.2	31
35	HPLC Analysis and Purification of Peptides. <i>Methods in Molecular Biology</i> , 2007, 386, 3-55.	0.9	36
36	Role of Peptide Hydrophobicity in the Mechanism of Action of α -Helical Antimicrobial Peptides. <i>Antimicrobial Agents and Chemotherapy</i> , 2007, 51, 1398-1406.	3.2	587

#	ARTICLE	IF	CITATIONS
37	Requirements for prediction of peptide retention time in reversed-phase high-performance liquid chromatography: Hydrophilicity/hydrophobicity of side-chains at the N- and C-termini of peptides are dramatically affected by the end-groups and location. <i>Journal of Chromatography A</i> , 2007, 1141, 212-225.	3.7	50
38	Preparative reversed-phase high-performance liquid chromatography collection efficiency for an antimicrobial peptide on columns of varying diameters (1mm to 9.4mm I.D.). <i>Journal of Chromatography A</i> , 2007, 1140, 112-120.	3.7	43
39	Template-based coiled-coil antigens elicit neutralizing antibodies to the SARS-coronavirus. <i>Journal of Structural Biology</i> , 2006, 155, 176-194.	2.8	38
40	Comparison of Biophysical and Biologic Properties of alpha-Helical Enantiomeric Antimicrobial Peptides. <i>Chemical Biology and Drug Design</i> , 2006, 67, 162-173.	3.2	113
41	Quantitation of the nearest-neighbour effects of amino acid side-chains that restrict conformational freedom of the polypeptide chain using reversed-phase liquid chromatography of synthetic model peptides with l- and d-amino acid substitutions. <i>Journal of Chromatography A</i> , 2006, 1123, 212-224.	3.7	20
42	One-step purification of a recombinant protein from a whole cell extract by reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2006, 1133, 248-253.	3.7	25
43	Determination of intrinsic hydrophilicity/hydrophobicity of amino acid side chains in peptides in the absence of nearest-neighbor or conformational effects. <i>Biopolymers</i> , 2006, 84, 283-297.	2.4	123
44	Rational Design of α -Helical Antimicrobial Peptides with Enhanced Activities and Specificity/Therapeutic Index. <i>Journal of Biological Chemistry</i> , 2005, 280, 12316-12329.	3.4	518
45	Structural Characterization of the SARS-Coronavirus Spike S Fusion Protein Core. <i>Journal of Biological Chemistry</i> , 2004, 279, 20836-20849.	3.4	182
46	Stabilizing and Destabilizing Clusters in the Hydrophobic Core of Long Two-stranded α -Helical Coiled-coils. <i>Journal of Biological Chemistry</i> , 2004, 279, 21576-21588.	3.4	96
47	Defining the minimum size of a hydrophobic cluster in two-stranded α -helical coiled-coils: Effects on protein stability. <i>Protein Science</i> , 2004, 13, 714-726.	7.6	37
48	Effect of chain length on coiled-coil stability: Decreasing stability with increasing chain length. <i>Biopolymers</i> , 2004, 76, 378-390.	2.4	20
49	Synthetic peptide vaccine development: measurement of polyclonal antibody affinity and cross-reactivity using a new peptide capture and release system for surface plasmon resonance spectroscopy. <i>Journal of Molecular Recognition</i> , 2004, 17, 540-557.	2.1	17
50	Unique stabilizing interactions identified in the two-stranded α -helical coiled-coil: Crystal structure of a cortaxillin I/GCN4 hybrid coiled-coil peptide. <i>Protein Science</i> , 2003, 12, 1395-1405.	7.6	33
51	Synthetic peptide vaccine and antibody therapeutic development: Prevention and treatment of <i>Pseudomonas aeruginosa</i> . <i>Biopolymers</i> , 2003, 71, 141-168.	2.4	43
52	Comparison of reversed-phase liquid chromatography and hydrophilic interaction/cation-exchange chromatography for the separation of amphipathic α -helical peptides with l- and d-amino acid substitutions in the hydrophilic face. <i>Journal of Chromatography A</i> , 2003, 1009, 61-71.	3.7	35
53	Temperature profiling of polypeptides in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2003, 1009, 29-43.	3.7	51
54	Temperature profiling of polypeptides in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2003, 1009, 45-59.	3.7	39

#	ARTICLE	IF	CITATIONS
55	Real-Time Monitoring of the Interactions of Two-Strandedde NovoDesigned Coiled-Coils:Â Effect of Chain Length on the Kinetic and Thermodynamic Constants of Bindingâ€. <i>Biochemistry</i> , 2003, 42, 1754-1763.	2.5	120
56	Solution Structure of the C-terminal Antiparallel Coiled-coil Domain from <i>Escherichia coli</i> Osmosensor ProP. <i>Journal of Molecular Biology</i> , 2003, 334, 1063-1076.	4.2	37
57	A Novel Method to Measure Self-association of Small Amphipathic Molecules. <i>Journal of Biological Chemistry</i> , 2003, 278, 22918-22927.	3.4	60
58	Clustering of Large Hydrophobes in the Hydrophobic Core of Two-stranded Î±-Helical Coiled-Coils Controls Protein Folding and Stability. <i>Journal of Biological Chemistry</i> , 2003, 278, 35248-35254.	3.4	39
59	A de Novo Designed Template for Generating Conformation-specific Antibodies That Recognize Î±-Helices in Proteins. <i>Journal of Biological Chemistry</i> , 2002, 277, 23515-23524.	3.4	23
60	Designing Heterodimeric Two-stranded Î±-Helical Coiled-coils. <i>Journal of Biological Chemistry</i> , 2002, 277, 37272-37279.	3.4	271
61	The Role of Unstructured Highly Charged Regions on the Stability and Specificity of Dimerization of Two-Stranded Î±-Helical Coiled-Coils: Analysis of the Neck-Hinge Region of the Kinesin-like Motor Protein Kif3A. <i>Journal of Structural Biology</i> , 2002, 137, 206-219.	2.8	19
62	Helix Capping Interactions Stabilize the N-Terminus of the Kinesin Neck Coiled-Coil. <i>Journal of Structural Biology</i> , 2002, 137, 220-235.	2.8	23
63	Preparative reversed-phase liquid chromatography of peptides. <i>Journal of Chromatography A</i> , 2002, 972, 87-99.	3.7	16
64	Are trigger sequences essential in the folding of two-stranded Î±-helical coiled-coils?11Edited by C. R. Matthews. <i>Journal of Molecular Biology</i> , 2001, 306, 539-553.	4.2	42
65	Peptide Models of the Helical Hydrophobic Transmembrane Segments of Membrane Proteins:Â Interactions of Acetyl-K2-(LA)12-K2-Amide with Phosphatidylethanolamine Bilayer Membranesâ€. <i>Biochemistry</i> , 2001, 40, 474-482.	2.5	36
66	A Polyalanine-Based Peptide Cannot Form a Stable Transmembrane Î±-Helix in Fully Hydrated Phospholipid Bilayersâ€. <i>Biochemistry</i> , 2001, 40, 12103-12111.	2.5	37
67	A Differential Scanning Calorimetric and ³¹ P NMR Spectroscopic Study of the Effect of Transmembrane Î±-Helical Peptides on the Lamellarâˆ“Reversed Hexagonal Phase Transition of Phosphatidylethanolamine Model Membranesâ€. <i>Biochemistry</i> , 2001, 40, 760-768.	2.5	43
68	Structural and functional studies on Troponin I and Troponin C interactions. <i>Journal of Cellular Biochemistry</i> , 2001, 83, 33-46.	2.6	18
69	Characterization of the biologically important interaction between troponin C and the N-terminal region of troponin I. <i>Journal of Cellular Biochemistry</i> , 2001, 83, 99-110.	2.6	8
70	Membrane-bound structure and alignment of the antimicrobial beta-sheet peptide gramicidin S derived from angular and distance constraints by solid state ¹⁹ F-NMR. <i>Journal of Biomolecular NMR</i> , 2001, 21, 191-208.	2.8	116
71	STABLECOIL: An Algorithm Designed to Predict the Location and Relative Stability of Coiled-Coils in Native Protein Sequences. , 2001, , 365-366.		7
72	The role of the carboxyl terminal Î±-helical coiled-coil domain in osmosensing by transporter ProP of <i>Escherichia coli</i> . <i>Journal of Molecular Recognition</i> , 2000, 13, 309-322.	2.1	54

#	ARTICLE	IF	CITATIONS
73	Development of the Structural Basis for Antimicrobial and Hemolytic Activities of Peptides Based on Gramicidin S and Design of Novel Analogs Using NMR Spectroscopy. <i>Journal of Biological Chemistry</i> , 2000, 275, 14287-14294.	3.4	60
74	Effects of side-chain characteristics on stability and oligomerization state of a de Novo -designed model coiled-coil: 20 amino acid substitutions in position α Edited by P. E. Wright. <i>Journal of Molecular Biology</i> , 2000, 300, 377-402.	4.2	238
75	The role of the carboxyl terminal \pm -helical coiled-coil domain in osmosensing by transporter ProP of <i>Escherichia coli</i> . <i>Journal of Molecular Recognition</i> , 2000, 13, 309-322.	2.1	1
76	^{13}C - ^{13}C Rotational resonance in a transmembrane peptide: A comparison of the fluid and gel phases. <i>Physical Review E</i> , 1999, 59, 5945-5957.	2.1	5
77	Dissociation of Antimicrobial and Hemolytic Activities in Cyclic Peptide Diastereomers by Systematic Alterations in Amphipathicity. <i>Journal of Biological Chemistry</i> , 1999, 274, 13181-13192.	3.4	230
78	New ice-binding face for type I antifreeze protein. <i>FEBS Letters</i> , 1999, 463, 87-91.	2.8	164
79	The Measure of Interior Disorder in a Folded Protein and Its Contribution to Stability. <i>Journal of the American Chemical Society</i> , 1999, 121, 8443-8449.	13.7	15
80	Contribution of Translational and Rotational Entropy to the Unfolding of a Dimeric Coiled-Coil. <i>Journal of Physical Chemistry B</i> , 1999, 103, 2270-2278.	2.6	31
81	Alternative Roles for Putative Ice-Binding Residues in Type I Antifreeze Protein. <i>Biochemistry</i> , 1999, 38, 4743-4749.	2.5	43
82	De novo design of a model peptide sequence to examine the effects of single amino acid substitutions in the hydrophobic core on both stability and oligomerization state of coiled-coils 1 Edited by P. Wright. <i>Journal of Molecular Biology</i> , 1999, 285, 785-803.	4.2	81
83	The role of position α in determining the stability and oligomerization state of \pm -helical coiled coils: 20 amino acid stability coefficients in the hydrophobic core of proteins. <i>Protein Science</i> , 1999, 8, 2312-2329.	7.6	149
84	Structural cassette mutagenesis in a de novo designed protein: Proof of a novel concept for examining protein folding and stability. , 1998, 47, 101-123.		7
85	Use of a heterodimeric coiled-coil system for biosensor application and affinity purification. <i>Biomedical Applications</i> , 1998, 715, 307-329.	1.7	85
86	Insights into the mechanism of heterodimerization from the ^1H -NMR solution structure of the c-Myc-Max heterodimeric leucine zipper. <i>Journal of Molecular Biology</i> , 1998, 281, 165-181.	4.2	97
87	Orientation, positional, additivity, and oligomerization-state effects of interhelical ion pairs in \pm -helical coiled-coils. <i>Journal of Molecular Biology</i> , 1998, 283, 993-1012.	4.2	96
88	Breakdown and Release of Myofilament Proteins During Ischemia and Ischemia/Reperfusion in Rat Hearts. <i>Circulation Research</i> , 1998, 82, 261-271.	4.5	218
89	Molecular Organization and Dynamics of 1-Palmitoyl-2-oleoylphosphatidylcholine Bilayers Containing a Transmembrane \pm -Helical Peptide. <i>Biochemistry</i> , 1998, 37, 3156-3164.	2.5	83
90	Effects of lanthanide binding on the stability of de novo designed \pm -helical coiled-coils. <i>Chemical Biology and Drug Design</i> , 1998, 51, 9-18.	1.1	15

#	ARTICLE	IF	CITATIONS
91	Î±-Helical Protein Assembly Motifs. <i>Journal of Biological Chemistry</i> , 1997, 272, 2583-2586.	3.4	169
92	Solution Secondary Structure of a Bacterially Expressed Peptide from the Receptor Binding Domain of <i>Pseudomonas aeruginosa</i> Pili Strain PAK: A Heteronuclear Multidimensional NMR Study. <i>Biochemistry</i> , 1997, 36, 12791-12801.	2.5	22
93	A Diminished Role for Hydrogen Bonds in Antifreeze Protein Binding to Ice. <i>Biochemistry</i> , 1997, 36, 14652-14660.	2.5	204
94	Salt effects on protein stability: two-stranded Î±-helical coiled-coils containing inter- or intrahelical ion pairs. <i>Journal of Molecular Biology</i> , 1997, 267, 1039-1052.	4.2	120
95	Positional dependence of the effects of negatively charged Glu side chains on the stability of two-stranded Î±-helical coiled-coils. <i>Journal of Peptide Science</i> , 1997, 3, 209-223.	1.4	17
96	Pilin C-terminal peptide binds asialoGM1 in liposomes: A 2D NMR study. <i>Protein Science</i> , 1997, 6, 2459-2461.	7.6	1
97	Ion Pairs Significantly Stabilize Coiled-coils in the Absence of Electrolyte. <i>Journal of Molecular Biology</i> , 1996, 255, 367-372.	4.2	68
98	Kinetic Study on the Formation of a de Novo Designed Heterodimeric Coiled-Coil: Use of Surface Plasmon Resonance To Monitor the Association and Dissociation of Polypeptide Chains. <i>Biochemistry</i> , 1996, 35, 12175-12185.	2.5	78
99	Formation of Parallel and Antiparallel Coiled-coils Controlled by the Relative Positions of Alanine Residues in the Hydrophobic Core. <i>Journal of Biological Chemistry</i> , 1996, 271, 3995-4001.	3.4	82
100	Investigation of electrostatic interactions in two-stranded coiled-coils through residue shuffling. <i>Biophysical Chemistry</i> , 1996, 59, 299-314.	2.8	52
101	Use of synthetic peptides to confirm that the <i>Pseudomonas aeruginosa</i> PAK pilus adhesin and the <i>Candida albicans</i> fimbrial adhesin possess a homologous receptor-binding domain. <i>Molecular Microbiology</i> , 1996, 19, 1107-1116.	2.5	25
102	A method for the facile solid-phase synthesis of gramicidin S and its analogs. <i>International Journal of Peptide Research and Therapeutics</i> , 1996, 3, 53-60.	0.1	21
103	A natural variant of type I antifreeze protein with four ice-binding repeats is a particularly potent antifreeze. <i>Protein Science</i> , 1996, 5, 1150-1156.	7.6	62
104	Engineering a de novo-designed coiled-coil heterodimerization domain for the rapid detection, purification and characterization of recombinantly expressed peptides and proteins. <i>Protein Engineering, Design and Selection</i> , 1996, 9, 1029-1042.	2.1	88
105	The relative positions of alanine residues in the hydrophobic core control the formation of two-stranded or four-stranded Î±-helical coiled-coils. <i>Protein Engineering, Design and Selection</i> , 1996, 9, 353-363.	2.1	39
106	De novo design of Î±-helical proteins: basic research to medical applications. <i>Biochemistry and Cell Biology</i> , 1996, 74, 133-154.	2.0	166
107	NMR solution structure of the receptor binding domain of <i>Pseudomonas aeruginosa</i> pilin strain P1. Identification of a Î²-turn. <i>International Journal of Peptide and Protein Research</i> , 1996, 48, 539-552.	0.1	11
108	Gramicidin S is active against both gram-positive and gram-negative bacteria. <i>International Journal of Peptide and Protein Research</i> , 1996, 47, 460-466.	0.1	149

#	ARTICLE	IF	CITATIONS
109	Lactam bridge stabilization of α -helical peptides: Ring size, orientation and positional effects. <i>Journal of Peptide Science</i> , 1995, 1, 274-282.	1.4	78
110	Relationship of sidechain hydrophobicity and α -helical propensity on the stability of the single-stranded amphipathic α -helix. <i>Journal of Peptide Science</i> , 1995, 1, 319-329.	1.4	317
111	The Effects of Interhelical Electrostatic Repulsions between Glutamic Acid Residues in Controlling the Dimerization and Stability of Two-stranded α -Helical Coiled-coils. <i>Journal of Biological Chemistry</i> , 1995, 270, 25495-25506.	3.4	78
112	Comparison of NMR solution structures of the receptor binding domains of <i>Pseudomonas aeruginosa</i> pili strains PAO, KB7, and PAK: implications for receptor binding and synthetic vaccine design. <i>Biochemistry</i> , 1995, 34, 16255-16268.	2.5	50
113	Structure Effects of Double D-Amino Acid Replacements: A Nuclear Magnetic Resonance and Circular Dichroism Study Using Amphipathic Model Helices. <i>Biochemistry</i> , 1995, 34, 12954-12962.	2.5	73
114	Protein destabilization by electrostatic repulsions in the two-stranded α -helical coiled-coil/leucine zipper. <i>Protein Science</i> , 1995, 4, 237-250.	7.6	108
115	Anti-peptide monoclonal antibody imaging of a common binding domain involved in muscle regulation. <i>Protein Science</i> , 1995, 4, 781-790.	7.6	2
116	Effect of trifluoroethanol on the solution structure and flexibility of desmopressin: a two-dimensional NMR study. <i>International Journal of Peptide and Protein Research</i> , 1995, 45, 471-481.	0.1	12
117	The net energetic contribution of interhelical electrostatic attractions to coiled-coil stability. <i>Protein Engineering, Design and Selection</i> , 1994, 7, 1365-1372.	2.1	115
118	Relative stabilities of synthetic peptide homo- and heterodimeric troponin-C domains. <i>Protein Science</i> , 1994, 3, 1010-1019.	7.6	19
119	Protein denaturation with guanidine hydrochloride or urea provides a different estimate of stability depending on the contributions of electrostatic interactions. <i>Protein Science</i> , 1994, 3, 1984-1991.	7.6	310
120	Conformational differences between cis and trans proline isomers of a peptide antigen representing the receptor binding domain of <i>Pseudomonas aeruginosa</i> as studied by $^1\text{H-NMR}$. <i>Biopolymers</i> , 1994, 34, 1221-1230.	2.4	12
121	Reversed-phase chromatography of synthetic amphipathic α -helical peptides as a model for ligand/receptor interactions Effect of changing hydrophobic environment on the relative hydrophilicity/hydrophobicity of amino acid side-chains. <i>Journal of Chromatography A</i> , 1994, 676, 139-153.	3.7	99
122	The Role of Interhelical Ionic Interactions in Controlling Protein Folding and Stability. <i>Journal of Molecular Biology</i> , 1994, 237, 500-512.	4.2	208
123	Electrostatic Interactions Control the Parallel and Antiparallel Orientation of α -Helical Chains in Two-Stranded α -Helical Coiled-Coils. <i>Biochemistry</i> , 1994, 33, 3862-3871.	2.5	186
124	Effect of Chain Length on the Formation and Stability of Synthetic α -Helical Coiled Coils. <i>Biochemistry</i> , 1994, 33, 15501-15510.	2.5	245
125	α -Helical Propensities of Amino Acids in the Hydrophobic Face of an Amphipathic α -Helix. <i>Protein and Peptide Letters</i> , 1994, 1, 114-119.	0.9	74
126	Effect of the α -amino group on peptide retention behaviour in reversed-phase chromatography Determination of the pK_a values of the α -amino group of 19 different N-terminal amino acid residues. <i>Journal of Chromatography A</i> , 1993, 646, 17-30.	3.7	94

#	ARTICLE	IF	CITATIONS
127	Packing and hydrophobicity effects on protein folding and stability: Effects of β -branched amino acids, valine and isoleucine, on the formation and stability of two-stranded α -helical coiled coils/leucine zippers. <i>Protein Science</i> , 1993, 2, 383-394.	7.6	165
128	A synthetic peptide mimics troponin I function in the calcium-dependent regulation of muscle contraction. <i>FEBS Letters</i> , 1993, 323, 223-228.	2.8	32
129	A single-stranded amphipathic α -helix in aqueous solution: Design, structural characterization, and its application for determining α -helical propensities of amino acids. <i>Biochemistry</i> , 1993, 32, 6190-6197.	2.5	80
130	Disulfide bond contribution to protein stability: Positional effects of substitution in the hydrophobic core of the two-stranded α -helical coiled-coil. <i>Biochemistry</i> , 1993, 32, 3178-3187.	2.5	171
131	NMR solution structure and flexibility of a peptide antigen representing the receptor binding domain of <i>Pseudomonas aeruginosa</i> . <i>Biochemistry</i> , 1993, 32, 13432-13440.	2.5	35
132	Synthetic model proteins: the relative contribution of leucine residues at the nonequivalent positions of the 3-4 hydrophobic repeat to the stability of the two-stranded α -helical coiled-coil. <i>Biochemistry</i> , 1992, 31, 5739-5746.	2.5	163
133	Relationship between amide proton chemical shifts and hydrogen bonding in amphipathic α -helical peptides. <i>Journal of the American Chemical Society</i> , 1992, 114, 4320-4326.	13.7	172
134	Role of interchain α -helical hydrophobic interactions in Ca^{2+} affinity, formation, and stability of a two-site domain in troponin C. <i>Protein Science</i> , 1992, 1, 945-955.	7.6	34
135	Antigen-antibody interactions: Elucidation of the epitope and strain-specificity of a monoclonal antibody directed against the pilin protein adherence binding domain of <i>Pseudomonas aeruginosa</i> strain K. <i>Protein Science</i> , 1992, 1, 1308-1318.	7.6	41
136	Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides. <i>Journal of Chromatography A</i> , 1992, 594, 75-86.	3.7	88
137	Unzipping the secrets of coiled-coils. <i>Current Biology</i> , 1992, 2, 122-124.	3.9	66
138	Stoichiometry of calcium binding to a synthetic heterodimeric troponin-C domain. <i>Biopolymers</i> , 1992, 32, 391-397.	2.4	13
139	The two-stranded α -helical coiled-coil is an ideal model for studying protein stability and subunit interactions. <i>Biopolymers</i> , 1992, 32, 419-426.	2.4	131
140	Hydrophilic-interaction chromatography of peptides on hydrophilic and strong cation-exchange columns. <i>Journal of Chromatography A</i> , 1991, 548, 13-24.	3.7	102
141	Multi-column preparative reversed-phase sample displacement chromatography of peptides. <i>Journal of Chromatography A</i> , 1991, 548, 267-280.	3.7	27
142	Reversed-phase chromatographic method development for peptide separations using the computer simulation program prodigest-1c. <i>Journal of Chromatography A</i> , 1989, 485, 365-382.	3.7	27
143	Correlation of protein retention times in reversed-phase chromatography with polypeptide chain length and hydrophobicity. <i>Journal of Chromatography A</i> , 1989, 476, 363-375.	3.7	95
144	Strong cation-exchange high-performance liquid chromatography of peptides. <i>Journal of Chromatography A</i> , 1989, 476, 377-389.	3.7	72

#	ARTICLE	IF	CITATIONS
145	Computer simulation of high-performance liquid chromatographic separations of peptide and protein digests for development of size-exclusion, ion-exchange and reversed-phase chromatographic methods. <i>Journal of Chromatography A</i> , 1988, 458, 147-167.	3.7	34
146	The Duchenne muscular dystrophy gene product is localized in sarcolemma of human skeletal muscle. <i>Nature</i> , 1988, 333, 466-469.	27.8	650
147	Effect of peptide chain length on peptide retention behaviour in reversed-phase chromatography. <i>Journal of Chromatography A</i> , 1988, 458, 193-205.	3.7	107
148	Preparative purification of peptides by reversed-phase chromatography. <i>Journal of Chromatography A</i> , 1988, 444, 349-362.	3.7	43
149	A Novel Approach to Reversed-Phase Preparative High-Performance Liquid Chromatography of Peptides. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1988, 11, 1229-1247.	1.0	21
150	Size-exclusion high-performance liquid chromatography of peptides. <i>Journal of Chromatography A</i> , 1987, 397, 99-112.	3.7	71
151	Effects of ion-pairing reagents on the prediction of peptide retention in reversed-phase high-resolution liquid chromatography. <i>Journal of Chromatography A</i> , 1987, 386, 205-222.	3.7	166
152	Prediction of peptide retention times in reversed-phase high-performance liquid chromatography II. Correlation of observed and predicted peptide retention times factors and influencing the retention times of peptides. <i>Journal of Chromatography A</i> , 1986, 359, 519-532.	3.7	123
153	Studies on the regulatory complex of rabbit skeletal muscle: Contributions of troponin subunits and tropomyosin in the presence and absence of Mg ²⁺ to the acto-S1 ATPase activity. <i>The Protein Journal</i> , 1986, 5, 335-354.	1.1	9
154	Effects of high-performance liquid chromatographic solvents and hydrophobic matrices on the secondary and quaternary structure of a model protein. <i>Journal of Chromatography A</i> , 1984, 317, 129-140.	3.7	145
155	Interaction of neuroleptic drugs with a synthetic calcium-binding peptide analog of site III of rabbit skeletal troponin C. <i>FEBS Letters</i> , 1983, 154, 60-64.	2.8	11
156	Primary sequence analysis and folding behavior of EF hands in relation to the mechanism of action of troponin C and calmodulin. <i>FEBS Letters</i> , 1983, 160, 1-6.	2.8	78
157	A new chromophoric substrate for penicillopepsin and other fungal aspartic proteinases. <i>Biochemical Journal</i> , 1982, 203, 603-610.	3.7	41
158	HPLC Analysis and Purification of Peptides. , 0, , 3-55.		2