

Kiyonori Miura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/189738/publications.pdf>

Version: 2024-02-01

63
papers

1,432
citations

430874

18
h-index

345221

36
g-index

64
all docs

64
docs citations

64
times ranked

1986
citing authors

#	ARTICLE	IF	CITATIONS
1	Guidelines for obstetrical practice in Japan: Society of Obstetrics and Gynecology (JOG) and Japan Association of Obstetricians and Gynecologists (JAOG) 2014 edition. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014, 40, 1469-1499.	1.3	307
2	Identification of Pregnancy-Associated MicroRNAs in Maternal Plasma. <i>Clinical Chemistry</i> , 2010, 56, 1767-1771.	3.2	176
3	Characterization of placenta-specific microRNAs in fetal growth restriction pregnancy. <i>Prenatal Diagnosis</i> , 2013, 33, 214-222.	2.3	135
4	Identification of endometrioid endometrial carcinoma-associated microRNAs in tissue and plasma. <i>Gynecologic Oncology</i> , 2014, 132, 715-721.	1.4	74
5	Microarray comparative genomic hybridization (CGH)-based prenatal diagnosis for chromosome abnormalities using cell-free fetal DNA in amniotic fluid. <i>Journal of Human Genetics</i> , 2006, 51, 412-417.	2.3	45
6	Circulating chromosome 19 miRNA cluster microRNAs in pregnant women with severe pre-eclampsia. <i>Journal of Obstetrics and Gynaecology Research</i> , 2015, 41, 1526-1532.	1.3	45
7	Current status of non-invasive prenatal testing in Japan. <i>Journal of Obstetrics and Gynaecology Research</i> , 2017, 43, 1245-1255.	1.3	40
8	Origin and mechanisms of formation of fetus-in-fetu: Two cases with genotype and methylation analyses. <i>American Journal of Medical Genetics, Part A</i> , 2006, 140A, 1737-1743.	1.2	39
9	Increased level of cell-free placental mRNA in a subgroup of placenta previa that needs hysterectomy. <i>Prenatal Diagnosis</i> , 2008, 28, 805-809.	2.3	38
10	A strong association between human earwax-type and apocrine colostrum secretion from the mammary gland. <i>Human Genetics</i> , 2007, 121, 631-633.	3.8	34
11	Genome-wide association study of HPV-associated cervical cancer in Japanese women. <i>Journal of Medical Virology</i> , 2014, 86, 1153-1158.	5.0	27
12	Pregnancy-associated microRNAs in plasma as potential molecular markers of ectopic pregnancy. <i>Fertility and Sterility</i> , 2015, 103, 1202-1208.e1.	1.0	27
13	Classification of factors involved in nonreportable results of noninvasive prenatal testing (NIPT) and prediction of success rate of second NIPT. <i>Prenatal Diagnosis</i> , 2019, 39, 100-106.	2.3	27
14	Clinical outcome of infants with confined placental mosaicism and intrauterine growth restriction of unknown cause. <i>American Journal of Medical Genetics, Part A</i> , 2006, 140A, 1827-1833.	1.2	26
15	Uniparental disomy analysis in trios using genome-wide SNP array and whole-genome sequencing data imply segmental uniparental isodisomy in general populations. <i>Gene</i> , 2013, 512, 267-274.	2.2	26
16	Placental mRNA in Maternal Plasma and Its Clinical Application to the Evaluation of Placental Status in a Pregnant Woman with Placenta Previa-Percreta. <i>Clinical Chemistry</i> , 2005, 51, 923-925.	3.2	21
17	Identification of Complete Hydatidiform Mole Pregnancy-associated MicroRNAs in Plasma. <i>Clinical Chemistry</i> , 2013, 59, 1410-1412.	3.2	20
18	Retrospective details of false-positive and false-negative results in non-invasive prenatal testing for fetal trisomies 21, 18 and 13. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2021, 256, 75-81.	1.1	20

#	ARTICLE	IF	CITATIONS
19	Clinical application of fetal sex determination using cell-free fetal DNA in pregnant carriers of X-linked genetic disorders. <i>Journal of Human Genetics</i> , 2011, 56, 296-299.	2.3	19
20	Circulating levels of maternal plasma cell-free pregnancy-associated placenta-specific microRNAs are associated with placental weight. <i>Placenta</i> , 2014, 35, 848-851.	1.5	16
21	Effect of labor on plasma concentrations and postpartum clearance of cell-free, pregnancy-associated, placenta-specific microRNAs. <i>Prenatal Diagnosis</i> , 2015, 35, 44-50.	2.3	15
22	Circulating Levels of Pregnancy-Associated, Placenta-Specific microRNAs in Pregnant Women With Placental Abruption. <i>Reproductive Sciences</i> , 2017, 24, 148-155.	2.5	15
23	Impaired early-phase suppression of glucagon secretion after glucose load is associated with insulin requirement during pregnancy in gestational diabetes. <i>Journal of Diabetes Investigation</i> , 2020, 11, 232-240.	2.4	15
24	HTLV-1 targets human placental trophoblasts in seropositive pregnant women. <i>Journal of Clinical Investigation</i> , 2020, 130, 6171-6186.	8.2	15
25	Clinical applications of analysis of plasma circulating complete hydatidiform mole pregnancy-associated miRNAs in gestational trophoblastic neoplasia: A preliminary investigation. <i>Placenta</i> , 2014, 35, 787-789.	1.5	14
26	A significant association between rs8067378 at 17q12 and invasive cervical cancer originally identified by a genome-wide association study in Han Chinese is replicated in a Japanese population. <i>Journal of Human Genetics</i> , 2016, 61, 793-796.	2.3	13
27	Maternal age-specific risk for trisomy 21 based on the clinical performance of NIPT and empirically derived NIPT age-specific positive and negative predictive values in Japan. <i>Journal of Human Genetics</i> , 2018, 63, 1035-1040.	2.3	13
28	The possibility of microarray-based analysis using cell-free placental mRNA in maternal plasma. <i>Prenatal Diagnosis</i> , 2010, 30, 849-861.	2.3	12
29	Epidemiology of human papillomavirus genotypes in pregnant Japanese women. <i>Journal of Human Genetics</i> , 2011, 56, 313-315.	2.3	10
30	Cell-Free DNA Is More Sensitive than Cell-Free mRNA as a Marker for Evaluation of Fetal-Maternal Hemorrhage. <i>Clinical Chemistry</i> , 2006, 52, 2121-2123.	3.2	9
31	Increased Levels of Cell-Free miR-517a and Decreased Levels of Cell-Free miR-518b in Maternal Plasma Samples From Placenta Previa Pregnancies at 32 Weeks of Gestation. <i>Reproductive Sciences</i> , 2015, 22, 1569-1576.	2.5	9
32	Feasibility of placenta-derived mesenchymal stem cells as a tool for studying pregnancy-related disorders. <i>Scientific Reports</i> , 2017, 7, 46220.	3.3	8
33	Reference values for circulating pregnancy-associated microRNAs in maternal plasma and their clinical usefulness in uncomplicated pregnancy and hypertensive disorder of pregnancy. <i>Journal of Obstetrics and Gynaecology Research</i> , 2018, 44, 840-851.	1.3	8
34	Comprehensive immune complexome analysis detects disease-specific immune complex antigens in seminal plasma and follicular fluids derived from infertile men and women. <i>Clinica Chimica Acta</i> , 2019, 495, 545-551.	1.1	8
35	Genital human papilloma virus infection in mentally-institutionalized virgins. <i>Gynecologic Oncology</i> , 2007, 106, 488-489.	1.4	7
36	Does increased nuchal translucency indicate a fetal abnormality? A retrospective study to clarify the clinical significance of nuchal translucency in Japan. <i>Journal of Human Genetics</i> , 2008, 53, 688-693.	2.3	7

#	ARTICLE	IF	CITATIONS
37	Predominantly placenta-expressed mRNAs in maternal plasma as predictive markers for twin-twin transfusion syndrome. <i>Prenatal Diagnosis</i> , 2014, 34, 345-349.	2.3	7
38	Phase 2 single-arm study on the safety of maintenance niraparib in Japanese patients with platinum-sensitive relapsed ovarian cancer. <i>Journal of Gynecologic Oncology</i> , 2021, 32, e21.	2.2	7
39	Clinical Applications of Plasma Circulating mRNA Analysis in Cases of Gestational Trophoblastic Disease. <i>Clinical Chemistry</i> , 2005, 51, 1261-1263.	3.2	6
40	A case of Kallmann syndrome carrying a missense mutation in alternatively spliced exon 8A encoding the immunoglobulin-like domain IIIb of fibroblast growth factor receptor 1. <i>Human Reproduction</i> , 2010, 25, 1076-1080.	0.9	6
41	Pre-vaccination epidemiology of human papillomavirus infections in Japanese women with abnormal cytology. <i>Journal of Obstetrics and Gynaecology Research</i> , 2011, 37, 1666-1670.	1.3	6
42	Adult T-cell leukemia-lymphoma in a pregnant woman diagnosed as a human T-cell lymphotropic virus type 1 carrier. <i>Journal of Obstetrics and Gynaecology Research</i> , 2016, 42, 336-340.	1.3	6
43	Circulating levels of maternal plasma cell-free miR-21 are associated with maternal body mass index and neonatal birth weight. <i>Prenatal Diagnosis</i> , 2015, 35, 509-511.	2.3	5
44	The Effects of Endometriosis on Ovarian Functions. <i>Endocrines</i> , 2021, 2, 142-149.	1.0	5
45	Mesenchymal stem cell-derived extracellular vesicles as probable triggers of radiation-induced heart disease. <i>Stem Cell Research and Therapy</i> , 2021, 12, 422.	5.5	5
46	Retained products of conception (RPOC) following delivery without placenta previa: Which patients with RPOC show postpartum hemorrhage?. <i>Placenta</i> , 2022, 124, 12-17.	1.5	5
47	Circulating Cell-Free Placental mRNA in the Maternal Plasma as a Predictive Marker for Twin-Twin Transfusion Syndrome. <i>Clinical Chemistry</i> , 2007, 53, 1167-1168.	3.2	4
48	Huge uterine fibroid arising from primary uterine cervical diverticulum: a case report and review of the literatures. <i>Journal of Obstetrics and Gynaecology</i> , 2019, 39, 1186-1187.	0.9	4
49	Malignant transformation from mature cystic teratoma of the ovary. <i>Journal of Obstetrics and Gynaecology Research</i> , 2019, 45, 1957-1960.	1.3	4
50	Paediatric-onset haploinsufficiency of A20 associated with a novel and de novo nonsense TNFAIP3 mutation. <i>Rheumatology</i> , 2020, 59, e85-e87.	1.9	3
51	Challenges facing workstyle reform for Japanese obstetricians and gynecologists revealed from time studies. <i>Journal of Obstetrics and Gynaecology Research</i> , 2022, 48, 1580-1590.	1.3	3
52	Single human papillomavirus 16 or 52 infection and later cytological findings in Japanese women with NILM or ASC-US. <i>Journal of Human Genetics</i> , 2014, 59, 251-255.	2.3	2
53	Fetiform teratoma was a parthenogenetic tumor arising from a mature ovum. <i>Journal of Human Genetics</i> , 2017, 62, 803-808.	2.3	2
54	Bihormonal dysregulation of insulin and glucagon contributes to glucose intolerance development at one year post-delivery in women with gestational diabetes: a prospective cohort study using an early postpartum 75-g glucose tolerance test. <i>Endocrine Journal</i> , 2021, 68, 919-931.	1.6	2

#	ARTICLE	IF	CITATIONS
55	Evaluation of the clinical performance of noninvasive prenatal testing at a Japanese laboratory. <i>Journal of Obstetrics and Gynaecology Research</i> , 2021, 47, 3437-3446.	1.3	2
56	Long-term culture of rat hepatocytes using human amniotic membrane as a culture substrate. <i>Regenerative Therapy</i> , 2021, 18, 384-390.	3.0	2
57	Prevention of Human T-Cell Leukemia Virus Type 1 (HTLV-1) Mother-to-Child Transmission. , 2017, , 157-169.		2
58	Initial viral load in cases of single human papillomavirus 16 or 52 persistent infection is associated with progression of later cytopathological findings in the uterine cervix. <i>Journal of Medical Virology</i> , 2013, 85, 2093-2100.	5.0	1
59	Large deletion in 6q containing the TNFAIP3 gene associated with autoimmune lymphoproliferative syndrome. <i>Clinical Immunology</i> , 2022, 235, 108853.	3.2	1
60	Prevalence of common aneuploidy in twin pregnancies. <i>Journal of Human Genetics</i> , 2022, 67, 261-265.	2.3	1
61	Biological Differences Between Ovarian Cancer-associated Fibroblasts and Contralateral Normal Ovary-derived Mesenchymal Stem Cells. <i>Anticancer Research</i> , 2022, 42, 1729-1737.	1.1	1
62	Gene Disorders and Genetic Counseling. <i>Comprehensive Gynecology and Obstetrics</i> , 2021, , 297-305.	0.0	0
63	Port-site hernia after laparoscopic gynecological surgery. <i>Japanese Journal of Gynecologic and Obstetric Endoscopy</i> , 2020, 36, 119-124.	0.0	0