## **Gregor Liegl**

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1951855/publications.pdf

Version: 2024-02-01

22	791	13	526166
papers	citations	h-index	g-index
29	29	29	1533
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	General population normative data for the EORTC QLQ-C30 health-related quality of life questionnaire based on 15,386 persons across 13 European countries, Canada and the Unites States. European Journal of Cancer, 2019, 107, 153-163.	1.3	259
2	The effect of sleep deprivation on pain perception in healthy subjects: a meta-analysis. Sleep Medicine, 2015, 16, 1313-1320.	0.8	118
3	Neural correlates of somatoform disorders from a meta-analytic perspective on neuroimaging studies. NeuroImage: Clinical, 2016, 11, 606-613.	1.4	50
4	Neural correlates of conversion disorder: overview and meta-analysis of neuroimaging studies on motor conversion disorder. BMC Psychiatry, 2016, 16, 195.	1.1	43
5	Establishing the European Norm for the health-related quality of life domains of the computer-adaptive test EORTC CAT Core. European Journal of Cancer, 2019, 107, 133-141.	1.3	38
6	Updated EORTC QLQ-C30 general population norm data for Germany. European Journal of Cancer, 2020, 137, 161-170.	1.3	32
7	Patients' Perceptions of Risky Developments During Psychotherapy. Journal of Contemporary Psychotherapy, 2013, 43, 95-105.	0.7	31
8	A Modified Version of the Transactional Stress Concept According to Lazarus and Folkman Was Confirmed in a Psychosomatic Inpatient Sample. Frontiers in Psychology, 2021, 12, 584333.	1.1	25
9	Guided self-help interventions for irritable bowel syndrome. European Journal of Gastroenterology and Hepatology, 2015, 27, 1209-1221.	0.8	21
10	An initial psychometric evaluation of the German PROMIS v1.2 Physical Function item bank in patients with a wide range of health conditions. Clinical Rehabilitation, 2018, 32, 84-93.	1.0	20
11	Normative data for the EORTC QLQ-C30 from the Austrian general population. Health and Quality of Life Outcomes, 2020, 18, 275.	1.0	20
12	Using Patient Health Questionnaire-9 item parameters of a common metric resulted in similar depression scores compared to independent item response theory model reestimation. Journal of Clinical Epidemiology, 2016, 71, 25-34.	2.4	17
13	Languageâ€related differential item functioning between English and German PROMIS Depression items is negligible. International Journal of Methods in Psychiatric Research, 2017, 26, .	1.1	14
14	Varying the item format improved the range of measurement in patient-reported outcome measures assessing physical function. Arthritis Research and Therapy, 2017, 19, 66.	1.6	11
15	General Spanish population normative data analysis for the EORTC QLQ-C30 by sex, age, and health condition. Health and Quality of Life Outcomes, 2021, 19, 208.	1.0	11
16	<p>A Step Towards a Better Understanding of Pain Phenotypes: Latent Class Analysis in Chronic Pain Patients Receiving Multimodal Inpatient Treatment</p> . Journal of Pain Research, 2020, Volume 13, 1023-1038.	0.8	10
17	A metaâ€analytic review of brief guided selfâ€help education for chronic pain. European Journal of Pain, 2016, 20, 1551-1562.	1.4	9
18	A 67-item stress resilience item bank showing high content validity wasÂdeveloped in a psychosomatic sample. Journal of Clinical Epidemiology, 2018, 100, 1-12.	2.4	7

#	Article	IF	CITATIONS
19	Applying Item Response Theory to the OPD Structure Questionnaire: Identification of a Unidimensional Core Construct and Feasibility of Computer Adaptive Testing. Journal of Personality Assessment, 2021, 103, 645-658.	1.3	5
20	Using subdomain-specific item sets affected PROMIS physical function scores differently in cardiology and rheumatology patients. Journal of Clinical Epidemiology, 2020, 127, 151-160.	2.4	5
21	Frequently Used Patient-Reported Outcome Measures of General Physical Function Were Highly Correlated With a Multitask Performance Outcome Test Battery. Value in Health, 2022, 25, 1752-1759.	0.1	3
22	Hand-related physical function in rheumatic hand conditions: a protocol for developing a patient-reported outcome measurement instrument: TableÂ1. BMJ Open, 2016, 6, e011174.	0.8	2