

# Xing Li

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1986055/publications.pdf>

Version: 2024-02-01

40  
papers

1,276  
citations

430874

18  
h-index

377865

34  
g-index

46  
all docs

46  
docs citations

46  
times ranked

2789  
citing authors

#	ARTICLE	IF	CITATIONS
1	Associations of Short-Term and Long-Term Exposure to Ambient Air Pollutants With Hypertension. <i>Hypertension</i> , 2016, 68, 62-70.	2.7	239
2	Transitory presence of myeloid-derived suppressor cells in neonates is critical for control of inflammation. <i>Nature Medicine</i> , 2018, 24, 224-231.	30.7	150
3	Endoplasmic reticulum stress induced LOX <sup>+</sup> CD <sup>15</sup> <sup>+</sup> polymorphonuclear myeloid-derived suppressor cells in hepatocellular carcinoma. <i>Immunology</i> , 2018, 154, 144-155.	4.4	92
4	Platelet-to-lymphocyte ratio acts as a prognostic factor for patients with advanced hepatocellular carcinoma. <i>Tumor Biology</i> , 2015, 36, 2263-2269.	1.8	91
5	Comparison of Face-Touching Behaviors Before and During the Coronavirus Disease 2019 Pandemic. <i>JAMA Network Open</i> , 2020, 3, e2016924.	5.9	69
6	ICAM-1 controls development and function of ILC2. <i>Journal of Experimental Medicine</i> , 2018, 215, 2157-2174.	8.5	62
7	Lactoferrin-induced myeloid-derived suppressor cell therapy attenuates pathologic inflammatory conditions in newborn mice. <i>Journal of Clinical Investigation</i> , 2019, 129, 4261-4275.	8.2	59
8	Neutrophil count is associated with myeloid derived suppressor cell level and presents prognostic value for hepatocellular carcinoma patients. <i>Oncotarget</i> , 2017, 8, 24380-24388.	1.8	41
9	Neutrophil-to-lymphocyte ratio acts as a prognostic factor for patients with advanced hepatocellular carcinoma. <i>Tumor Biology</i> , 2014, 35, 11057-11063.	1.8	37
10	Atorvastatin promotes the expansion of myeloid-derived suppressor cells and attenuates murine colitis. <i>Immunology</i> , 2016, 149, 432-446.	4.4	34
11	Identification of abnormally expressed lncRNAs induced by PM2.5 in human bronchial epithelial cells. <i>Bioscience Reports</i> , 2018, 38, .	2.4	34
12	Combination of ULK1 and LC3B improve prognosis assessment of hepatocellular carcinoma. <i>Biomedicine and Pharmacotherapy</i> , 2018, 97, 195-202.	5.6	33
13	Identification of ceRNA network based on a RNA-seq shows prognostic lncRNA biomarkers in human lung adenocarcinoma. <i>Oncology Letters</i> , 2018, 16, 5697-5708.	1.8	30
14	<p></p>MicroRNA-126 Inhibit Viability of Colorectal Cancer Cell by Repressing mTOR Induced Apoptosis and Autophagy<p>. <i>OncoTargets and Therapy</i> , 2020, Volume 13, 2459-2468.	2.0	28
15	IRF-7 Is a Critical Regulator of Type 2 Innate Lymphoid Cells in Allergic Airway Inflammation. <i>Cell Reports</i> , 2019, 29, 2718-2730.e6.	6.4	27
16	Circular RNA circCDK13 suppresses cell proliferation, migration and invasion by modulating the JAK/STAT and PI3K/AKT pathways in liver cancer. <i>International Journal of Oncology</i> , 2018, 53, 246-256.	3.3	26
17	Expansion of polymorphonuclear myeloid-derived suppressor cells in patients with end-stage renal disease may lead to infectious complications. <i>Kidney International</i> , 2017, 91, 1236-1242.	5.2	22
18	Coenzyme Q10 Prevents the Interleukin-1 Beta Induced Inflammatory Response via Inhibition of MAPK Signaling Pathways in Rat Articular Chondrocytes. <i>Drug Development Research</i> , 2017, 78, 403-410.	2.9	19

#	ARTICLE	IF	CITATIONS
19	Brazilin induces FOXO3A-dependent autophagic cell death by disturbing calcium homeostasis in osteosarcoma cells. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 479-491.	2.3	18
20	Aberrant expression of enhancer of zeste homologue 2, correlated with HIF-1 $\pm$ , refines relapse risk and predicts poor outcome for breast cancer. <i>Oncology Reports</i> , 2014, 32, 1101-1107.	2.6	17
21	Expansion of monocytic myeloid-derived suppressor cells in endometriosis patients: A pilot study. <i>International Immunopharmacology</i> , 2017, 47, 150-158.	3.8	17
22	Frequency of asthma exacerbation in children during the coronavirus disease pandemic with strict mitigative countermeasures. <i>Pediatric Pulmonology</i> , 2021, 56, 1455-1463.	2.0	16
23	Hypoxemia is an independent predictor of bronchiolitis obliterans following respiratory adenoviral infection in children. <i>SpringerPlus</i> , 2016, 5, 1622.	1.2	15
24	Prognostic value of serum HIF-1 $\pm$ change following transarterial chemoembolization in hepatocellular carcinoma. <i>Clinical and Experimental Medicine</i> , 2021, 21, 109-120.	3.6	12
25	Selection of glucocorticoid-sensitive patients in interstitial lung disease secondary to connective tissue diseases population by radiomics. <i>Therapeutics and Clinical Risk Management</i> , 2018, Volume 14, 1975-1986.	2.0	11
26	Comparison of five staging systems in predicting the survival rate of patients with hepatocellular carcinoma undergoing transarterial chemoembolization therapy. <i>Oncology Letters</i> , 2017, 15, 855-862.	1.8	9
27	miR-30b regulates chondrogenic differentiation of mouse embryo-derived stem cells by targeting SOX9. <i>Experimental and Therapeutic Medicine</i> , 2017, 14, 6131-6137.	1.8	8
28	Interferon Regulatory Factor 4 Regulates the Development of Polymorphonuclear Myeloid-Derived Suppressor Cells Through the Transcription of c-Myc in Cancer. <i>Frontiers in Immunology</i> , 2021, 12, 627072.	4.8	8
29	Neutrophil-to-Apolipoprotein A1 Ratio Predicted Overall Survival in Hepatocellular Carcinoma Receiving Transarterial Chemoembolization. <i>Oncologist</i> , 2021, 26, e1434-e1444.	3.7	8
30	Angiotensin II enhances group 2 innate lymphoid cell responses via AT1a during airway inflammation. <i>Journal of Experimental Medicine</i> , 2022, 219, .	8.5	8
31	Serum Golgi protein 73 is a prognostic rather than diagnostic marker in hepatocellular carcinoma. <i>Oncology Letters</i> , 2017, 14, 6277-6284.	1.8	6
32	Alanine aminotransferase to hemoglobin ratio is an indicator for disease progression for hepatocellular carcinoma patients receiving transcatheter arterial chemoembolization. <i>Tumor Biology</i> , 2016, 37, 2951-2959.	1.8	5
33	Different patient satisfaction levels between the first and second knee in the early stage after simultaneous bilateral total knee arthroplasty (TKA): a comparison between subjective and objective outcome assessments. <i>Journal of Orthopaedic Surgery and Research</i> , 2017, 12, 121.	2.3	5
34	&lt;p>&gt;Neutrophil-To-Lymphocyte Ratio Predicted Long-Term Chemotherapy Benefits In Stage IIIB-IV Non-Squamous Non-Small Cell Lung Cancer Patients Without Sensitive Mutations&lt;/p>&lt;p>&gt;. <i>OncoTargets and Therapy</i> , 2019, Volume 12, 8779-8787.	2.0	5
35	Expansion of Group 2 Innate Lymphoid Cells in Patients with End-Stage Renal Disease and Their Clinical Significance. <i>Journal of Immunology</i> , 2020, 205, 36-44.	0.8	4
36	Accumulation of LOX-1+ PMN-MDSCs in nasopharyngeal carcinoma survivors with chronic hepatitis B might permit immune tolerance to epstein-bar virus and relate to tumor recurrence. <i>Aging</i> , 2021, 13, 437-449.	3.1	3

#	ARTICLE	IF	CITATIONS
37	A 2-year step-down withdrawal from inhaled corticosteroids in asthmatic children receiving immunotherapy. <i>World Journal of Pediatrics</i> , 2017, 13, 560-565.	1.8	2
38	Timing of regular egg intake for prevention of egg allergy. <i>Journal of Allergy and Clinical Immunology</i> , 2018, 141, 459-460.	2.9	2
39	Dexamethasone and lactoferrin induced PMN-MDSCs relieved inflammatory adverse events of anti-cancer therapy without tumor promotion. <i>Communications Biology</i> , 2021, 4, 252.	4.4	2
40	Expansion of Monocytic Myeloid-Derived Suppressor Cells in Patients Under Hemodialysis Might Lead to Cardiovascular and Cerebrovascular Events. <i>Frontiers in Immunology</i> , 2020, 11, 577253.	4.8	2