

Mafalda Oliveira

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1989630/publications.pdf>

Version: 2024-02-01

123
papers

6,136
citations

136950

32
h-index

76900

74
g-index

126
all docs

126
docs citations

126
times ranked

6973
citing authors

#	ARTICLE	IF	CITATIONS
1	Tucatinib, Trastuzumab, and Capecitabine for HER2-Positive Metastatic Breast Cancer. <i>New England Journal of Medicine</i> , 2020, 382, 597-609.	27.0	789
2	Cerebrospinal fluid-derived circulating tumour DNA better represents the genomic alterations of brain tumours than plasma. <i>Nature Communications</i> , 2015, 6, 8839.	12.8	605
3	Sacituzumab Govitecan in Metastatic Triple-Negative Breast Cancer. <i>New England Journal of Medicine</i> , 2021, 384, 1529-1541.	27.0	601
4	Ipatasertib plus paclitaxel versus placebo plus paclitaxel as first-line therapy for metastatic triple-negative breast cancer (LOTUS): a multicentre, randomised, double-blind, placebo-controlled, phase 2 trial. <i>Lancet Oncology</i> , The, 2017, 18, 1360-1372.	10.7	377
5	Neratinib Plus Capecitabine Versus Lapatinib Plus Capecitabine in HER2-Positive Metastatic Breast Cancer Previously Treated With 2 HER2-Directed Regimens: Phase III NALA Trial. <i>Journal of Clinical Oncology</i> , 2020, 38, 3138-3149.	1.6	355
6	Intracranial Efficacy and Survival With Tucatinib Plus Trastuzumab and Capecitabine for Previously Treated HER2-Positive Breast Cancer With Brain Metastases in the HER2CLIMB Trial. <i>Journal of Clinical Oncology</i> , 2020, 38, 2610-2619.	1.6	331
7	Capturing intra-tumor genetic heterogeneity by de novo mutation profiling of circulating cell-free tumor DNA: a proof-of-principle. <i>Annals of Oncology</i> , 2014, 25, 1729-1735.	1.2	308
8	HER2-enriched subtype as a predictor of pathological complete response following trastuzumab and lapatinib without chemotherapy in early-stage HER2-positive breast cancer (PAMELA): an open-label, single-group, multicentre, phase 2 trial. <i>Lancet Oncology</i> , The, 2017, 18, 545-554.	10.7	250
9	Biomarker analyses in the phase III ASCENT study of sacituzumab govitecan versus chemotherapy in patients with metastatic triple-negative breast cancer. <i>Annals of Oncology</i> , 2021, 32, 1148-1156.	1.2	146
10	A First-in-Human Phase I Study of the ATP-Competitive AKT Inhibitor Ipatasertib Demonstrates Robust and Safe Targeting of AKT in Patients with Solid Tumors. <i>Cancer Discovery</i> , 2017, 7, 102-113.	9.4	136
11	Ribociclib plus letrozole versus chemotherapy for postmenopausal women with hormone receptor-positive, HER2-negative, luminal B breast cancer (CORALLEEN): an open-label, multicentre, randomised, phase 2 trial. <i>Lancet Oncology</i> , The, 2020, 21, 33-43.	10.7	105
12	Early ctDNA dynamics as a surrogate for progression-free survival in advanced breast cancer in the BEECH trial. <i>Annals of Oncology</i> , 2019, 30, 945-952.	1.2	103
13	Primary results of LORELEI: A phase II randomized, double-blind study of neoadjuvant letrozole (LET) plus tasiselisib versus LET plus placebo (PLA) in postmenopausal patients (pts) with ER+/HER2-negative early breast cancer (EBC). <i>Annals of Oncology</i> , 2017, 28, v605.	1.2	103
14	FAIRLANE, a double-blind placebo-controlled randomized phase II trial of neoadjuvant ipatasertib plus paclitaxel for early triple-negative breast cancer. <i>Annals of Oncology</i> , 2019, 30, 1289-1297.	1.2	97
15	HER2-Enriched Subtype and ERBB2 Expression in HER2-Positive Breast Cancer Treated with Dual HER2 Blockade. <i>Journal of the National Cancer Institute</i> , 2020, 112, 46-54.	6.3	97
16	A predictive model of pathologic response based on tumor cellularity and tumor-infiltrating lymphocytes (CeTIL) in HER2-positive breast cancer treated with chemo-free dual HER2 blockade. <i>Annals of Oncology</i> , 2018, 29, 170-177.	1.2	84
17	Next Generation-Targeted Amplicon Sequencing (NG-TAS): an optimised protocol and computational pipeline for cost-effective profiling of circulating tumour DNA. <i>Genome Medicine</i> , 2019, 11, 1.	8.2	84
18	Genomic and Transcriptomic Analyses of Breast Cancer Primaries and Matched Metastases in AURORA, the Breast International Group (BIG) Molecular Screening Initiative. <i>Cancer Discovery</i> , 2021, 11, 2796-2811.	9.4	79

#	ARTICLE	IF	CITATIONS
19	Neoadjuvant letrozole plus taselesib versus letrozole plus placebo in postmenopausal women with oestrogen receptor-positive, HER2-negative, early-stage breast cancer (LORELEI): a multicentre, randomised, double-blind, placebo-controlled, phase 2 trial. <i>Lancet Oncology</i> , The, 2019, 20, 1226-1238.	10.7	76
20	Neratinib + capecitabine versus lapatinib + capecitabine in patients with HER2+ metastatic breast cancer previously treated with 2 HER2-directed regimens: Findings from the multinational, randomized, phase III NALA trial.. <i>Journal of Clinical Oncology</i> , 2019, 37, 1002-1002.	1.6	71
21	Capturing Hyperprogressive Disease with Immune-Checkpoint Inhibitors Using RECIST 1.1 Criteria. <i>Clinical Cancer Research</i> , 2020, 26, 1846-1855.	7.0	70
22	Palbociclib and Trastuzumab in HER2-Positive Advanced Breast Cancer: Results from the Phase II SOLTI-1303 PATRICIA Trial. <i>Clinical Cancer Research</i> , 2020, 26, 5820-5829.	7.0	68
23	Phenotypic changes of HER2-positive breast cancer during and after dual HER2 blockade. <i>Nature Communications</i> , 2020, 11, 385.	12.8	67
24	Complete response in HER2+leptomeningeal carcinomatosis from breast cancer with intrathecal trastuzumab. <i>Breast Cancer Research and Treatment</i> , 2011, 127, 841-844.	2.5	59
25	BEECH: a dose-finding run-in followed by a randomised phase II study assessing the efficacy of AKT inhibitor capivasertib (AZD5363) combined with paclitaxel in patients with estrogen receptor-positive advanced or metastatic breast cancer, and in a PIK3CA mutant sub-population. <i>Annals of Oncology</i> , 2019, 30, 774-780.	1.2	57
26	Capivasertib, an AKT Kinase Inhibitor, as Monotherapy or in Combination with Fulvestrant in Patients with AKT1 E17K-Mutant, ER-Positive Metastatic Breast Cancer. <i>Clinical Cancer Research</i> , 2020, 26, 3947-3957.	7.0	54
27	Brain Metastases in HER2-Positive Breast Cancer: Current and Novel Treatment Strategies. <i>Cancers</i> , 2021, 13, 2927.	3.7	54
28	A multivariable prognostic score to guide systemic therapy in early-stage HER2-positive breast cancer: a retrospective study with an external evaluation. <i>Lancet Oncology</i> , The, 2020, 21, 1455-1464.	10.7	52
29	Phase II Study of Taselesib (GDC-0032) in Combination with Fulvestrant in Patients with HER2-Negative, Hormone Receptor-Positive Advanced Breast Cancer. <i>Clinical Cancer Research</i> , 2018, 24, 4380-4387.	7.0	49
30	Preclinical In Vivo Validation of the RAD51 Test for Identification of Homologous Recombination-Deficient Tumors and Patient Stratification. <i>Cancer Research</i> , 2022, 82, 1646-1657.	0.9	40
31	Metabolic Imaging Detects Resistance to PI3K± Inhibition Mediated by Persistent FOXM1 Expression in ER+ Breast Cancer. <i>Cancer Cell</i> , 2020, 38, 516-533.e9.	16.8	38
32	Final results of the double-blind placebo-controlled randomized phase 2 LOTUS trial of first-line ipatasertib plus paclitaxel for inoperable locally advanced/metastatic triple-negative breast cancer. <i>Breast Cancer Research and Treatment</i> , 2021, 189, 377-386.	2.5	38
33	Genetic heterogeneity and actionable mutations in HER2-positive primary breast cancers and their brain metastases. <i>Oncotarget</i> , 2018, 9, 20617-20630.	1.8	36
34	Prognostic value of ctDNA detection in patients with early breast cancer undergoing neoadjuvant therapy: A systematic review and meta-analysis. <i>Cancer Treatment Reviews</i> , 2022, 104, 102362.	7.7	33
35	Ipatasertib plus paclitaxel for PIK3CA/AKT1/PTEN-altered hormone receptor-positive HER2-negative advanced breast cancer: primary results from cohort B of the IPATunity130 randomized phase 3 trial. <i>Breast Cancer Research and Treatment</i> , 2022, 191, 565-576.	2.5	32
36	Efficacy of Neratinib Plus Capecitabine in the Subgroup of Patients with Central Nervous System Involvement from the NALA Trial. <i>Oncologist</i> , 2021, 26, e1327-e1338.	3.7	31

#	ARTICLE	IF	CITATIONS
37	Safety, activity, and molecular heterogeneity following neoadjuvant non-pegylated liposomal doxorubicin, paclitaxel, trastuzumab, and pertuzumab in HER2-positive breast cancer (Opti-HER HEART): an open-label, single-group, multicenter, phase 2 trial. <i>BMC Medicine</i> , 2019, 17, 8.	5.5	28
38	Implication of breast cancer phenotype for patients with leptomeningeal carcinomatosis. <i>Breast</i> , 2013, 22, 19-23.	2.2	27
39	Immune microenvironment characterisation and dynamics during anti-HER2-based neoadjuvant treatment in HER2-positive breast cancer. <i>Npj Precision Oncology</i> , 2021, 5, 23.	5.4	26
40	SEOM clinical guidelines in metastatic breast cancer 2015. <i>Clinical and Translational Oncology</i> , 2015, 17, 946-955.	2.4	25
41	A phase I dose escalation and expansion study of the next generation oral SERD AZD9833 in women with ER-positive, HER2-negative advanced breast cancer.. <i>Journal of Clinical Oncology</i> , 2020, 38, 1024-1024.	1.6	25
42	Overall survival (OS) update of the double-blind placebo (PBO)-controlled randomized phase 2 LOTUS trial of first-line ipatasertib (IPAT) + paclitaxel (PAC) for locally advanced/metastatic triple-negative breast cancer (mTNBC).. <i>Journal of Clinical Oncology</i> , 2018, 36, 1008-1008.	1.6	24
43	Genetic Alterations in the PI3K/AKT Pathway and Baseline AKT Activity Define AKT Inhibitor Sensitivity in Breast Cancer Patient-derived Xenografts. <i>Clinical Cancer Research</i> , 2020, 26, 3720-3731.	7.0	21
44	A phase II study of the PI3K inhibitor taselisib (GDC-0032) combined with fulvestrant (F) in patients (pts) with HER2-negative (HER2-), hormone receptor-positive (HR+) advanced breast cancer (BC).. <i>Journal of Clinical Oncology</i> , 2016, 34, 520-520.	1.6	21
45	Functional Mapping of AKT Signaling and Biomarkers of Response from the FAIRLANE Trial of Neoadjuvant Ipatasertib plus Paclitaxel for Triple-Negative Breast Cancer. <i>Clinical Cancer Research</i> , 2022, 28, 993-1003.	7.0	21
46	POSEIDON Trial Phase 1b Results: Safety, Efficacy and Circulating Tumor DNA Response of the Beta Isoform-Sparing PI3K Inhibitor Taselisib (GDC-0032) Combined with Tamoxifen in Hormone Receptor Positive Metastatic Breast Cancer Patients. <i>Clinical Cancer Research</i> , 2019, 25, 6598-6605.	7.0	17
47	SOLTI-1805 TOT-HER3 Study Concept: A Window-of-Opportunity Trial of Patritumab Deruxtecan, a HER3 Directed Antibody Drug Conjugate, in Patients With Early Breast Cancer. <i>Frontiers in Oncology</i> , 2021, 11, 638482.	2.8	16
48	IPATunity130: A pivotal randomized phase III trial evaluating ipatasertib (IPAT) + paclitaxel (PAC) for PIK3CA/AKT1/PTEN-altered advanced triple-negative (TN) or hormone receptor-positive HER2-negative (HR+/HER2-) breast cancer (BC).. <i>Journal of Clinical Oncology</i> , 2018, 36, TPS1117-TPS1117.	1.6	16
49	SEOM clinical guidelines in advanced and recurrent breast cancer (2018). <i>Clinical and Translational Oncology</i> , 2019, 21, 31-45.	2.4	14
50	Biomarker Analysis of the Phase III NALA Study of Neratinib + Capecitabine versus Lapatinib + Capecitabine in Patients with Previously Treated Metastatic Breast Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 5818-5827.	7.0	14
51	PI3K activation promotes resistance to eribulin in HER2-negative breast cancer. <i>British Journal of Cancer</i> , 2021, 124, 1581-1591.	6.4	12
52	High FGFR1 mRNA Expression Levels Correlate with Response to Selective FGFR Inhibitors in Breast Cancer. <i>Clinical Cancer Research</i> , 2022, 28, 137-149.	7.0	12
53	Circulating Tumor DNA and Biomarker Analyses From the LOTUS Randomized Trial of First-Line Ipatasertib and Paclitaxel for Metastatic Triple-Negative Breast Cancer. <i>JCO Precision Oncology</i> , 2020, 4, 1012-1024.	3.0	11
54	Phase Ib Dose-escalation/Expansion Trial of Ribociclib in Combination With Everolimus and Exemestane in Postmenopausal Women with HR+, HER2+ Advanced Breast Cancer. <i>Clinical Cancer Research</i> , 2020, 26, 6417-6428.	7.0	11

#	ARTICLE	IF	CITATIONS
55	Abstract PD5-2: Ph1b study of the PI3K inhibitor taselisib (GDC-0032) in combination with letrozole in patients with hormone receptor-positive advanced breast cancer. <i>Cancer Research</i> , 2015, 75, PD5-2-PD5-2.	0.9	11
56	Evolving Landscape of Molecular Prescreening Strategies for Oncology Early Clinical Trials. <i>JCO Precision Oncology</i> , 2020, 4, 505-513.	3.0	10
57	HER2-enriched subtype and ERBB2 mRNA as predictors of pathological complete response following trastuzumab and lapatinib without chemotherapy in early-stage HER2-positive breast cancer: A combined analysis of TBCRC006/023 and PAMELA trials.. <i>Journal of Clinical Oncology</i> , 2018, 36, 509-509.	1.6	10
58	Management of the axilla in early breast cancer patients in the genomic era. <i>Annals of Oncology</i> , 2013, 24, 1163-1170.	1.2	9
59	1O Neratinib + capecitabine vs lapatinib + capecitabine in HER2+ metastatic breast cancer previously treated with 2 HER2-directed regimens: Exploratory biomarker analyses from phase III NALA trial. <i>Annals of Oncology</i> , 2020, 31, S15.	1.2	9
60	Independent Validation of the PAM50-Based Chemo-Endocrine Score (CES) in Hormone Receptor-Positive HER2-Positive Breast Cancer Treated with Neoadjuvant Anti-HER2-Based Therapy. <i>Clinical Cancer Research</i> , 2021, 27, 3116-3125.	7.0	9
61	Neoadjuvant eribulin in HER2-negative early-stage breast cancer (SOLTI-1007-NeoEribulin): a multicenter, two-cohort, non-randomized phase II trial. <i>Npj Breast Cancer</i> , 2021, 7, 145.	5.2	9
62	The AURORA pilot study for molecular screening of patients with advanced breast cancer—a study of the breast international group. <i>Npj Breast Cancer</i> , 2017, 3, 23.	5.2	8
63	SOLTI-1503 PROMETEO TRIAL: combination of talimogene laherparepvec with atezolizumab in early breast cancer. <i>Future Oncology</i> , 2020, 16, 1801-1813.	2.4	8
64	Tucatinib versus placebo added to trastuzumab and capecitabine for patients with previously treated HER2+ metastatic breast cancer with brain metastases (HER2CLIMB).. <i>Journal of Clinical Oncology</i> , 2020, 38, 1005-1005.	1.6	8
65	Oestrogen receptor activity in hormone-dependent breast cancer during chemotherapy. <i>EBioMedicine</i> , 2021, 69, 103451.	6.1	7
66	Concordance of genomic alterations (GA) in synchronous tumor biopsies (tBx) and circulating tumor (ct) DNA from metastatic breast cancer (MBC) patients (pts).. <i>Journal of Clinical Oncology</i> , 2018, 36, 1073-1073.	1.6	7
67	Sudden death during adjuvant trastuzumab therapy of breast cancer. <i>Annals of Oncology</i> , 2010, 21, 901.	1.2	6
68	Neratinib plus capecitabine for the treatment of advanced HER2-positive breast cancer. <i>Expert Review of Anticancer Therapy</i> , 2020, 20, 731-741.	2.4	6
69	POSEIDON trial phase 1b results: Safety and preliminary efficacy of the isoform selective PI3K inhibitor taselisib (GDC-0032) combined with tamoxifen in hormone receptor (HR) positive, HER2-negative metastatic breast cancer (MBC) patients (pts) - including response monitoring by plasma circulating tumor (ct) DNA.. <i>Journal of Clinical Oncology</i> , 2016, 34, 2520-2520.	1.6	6
70	Phase 2 Study of Trabectedin in Patients With Hormone Receptor-Positive, HER-2-Negative, Advanced Breast Carcinoma According to Expression of Xeroderma Pigmentosum G Gene. <i>Clinical Breast Cancer</i> , 2016, 16, 364-371.	2.4	5
71	SOLTI-1303 PATRICIA: A phase II study of palbociclib and trastuzumab (HR+ with or without letrozole) in trastuzumab-pretreated, postmenopausal patients with HER2-positive metastatic breast cancer.. <i>Journal of Clinical Oncology</i> , 2018, 36, TPS1101-TPS1101.	1.6	5
72	A phase Ib, open-label, dose-escalation study of the safety and pharmacology of taselisib (GDC-0032) in combination with either docetaxel or paclitaxel in patients with HER2-negative, locally advanced, or metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019, 178, 121-133.	2.5	4

#	ARTICLE	IF	CITATIONS
73	Palbociclib combined with endocrine therapy in heavily pretreated HR+/HER2- advanced breast cancer patients: Results from the compassionate use program in Spain (PALBOCOMP). <i>Breast</i> , 2020, 54, 286-292.	2.2	4
74	Abstract CT041: Primary results from FAIRLANE (NCT02301988), a double-blind placebo (PBO)-controlled randomized phase II trial of neoadjuvant ipatasertib (IPAT) + paclitaxel (PAC) for early triple-negative breast cancer (eTNBC). , 2018, , .		4
75	Abstract CT046: A phase I basket study of the PI3K inhibitor taselelisib (GDC-0032) in <i>PIK3CA</i>-mutated locally advanced or metastatic solid tumors. <i>Cancer Research</i> , 2018, 78, CT046-CT046.	0.9	4
76	PI3K pathway (PI3Kp) dysregulation and response to pan-PI3K/AKT/mTOR/dual PI3K-mTOR inhibitors (PI3Kpi) in metastatic breast cancer (MBC) patients (pts).. <i>Journal of Clinical Oncology</i> , 2012, 30, 509-509.	1.6	3
77	LOTUS: A randomized, phase II, multicenter, placebo-controlled study of ipatasertib (Ipat, GDC-0068), an inhibitor of Akt, in combination with paclitaxel (Pac) as front-line treatment for patients (pts) with metastatic triple-negative breast cancer (TNBC).. <i>Journal of Clinical Oncology</i> , 2015, 33, TPS1111-TPS1111.	1.6	3
78	LORELEI: A Phase II randomized, double-blind study of neoadjuvant letrozole plus taselelisib (GDC-0032) versus letrozole plus placebo in postmenopausal women with ER-positive/HER2-negative, early-stage breast cancer.. <i>Journal of Clinical Oncology</i> , 2016, 34, TPS613-TPS613.	1.6	3
79	LOTUS (NCT02162719): A double-blind placebo (PBO)-controlled randomized phase II trial of first-line ipatasertib (IPAT) + paclitaxel (P) for metastatic triple-negative breast cancer (TNBC).. <i>Journal of Clinical Oncology</i> , 2017, 35, 1009-1009.	1.6	3
80	A phase I/II dose escalation and expansion study to investigate the safety, pharmacokinetics, pharmacodynamics and clinical activity of GSK525762 in combination with fulvestrant in subjects with ER+ breast cancer.. <i>Journal of Clinical Oncology</i> , 2017, 35, TPS1114-TPS1114.	1.6	3
81	Clonality of PIK3CA mutations (mut) and efficacy of PI3K/AKT/mTOR inhibitors (PAMi) in patients (pts) with metastatic breast cancer (MBC).. <i>Journal of Clinical Oncology</i> , 2016, 34, 528-528.	1.6	3
82	First Nationwide Molecular Screening Program in Spain for Patients With Advanced Breast Cancer: Results From the AGATA SOLTI-1301 Study. <i>Frontiers in Oncology</i> , 2021, 11, 744112.	2.8	3
83	Tackling the Biological Diversity in Early Triple-Negative Breast Cancer. <i>Breast Care</i> , 2020, 15, 205-207.	1.4	2
84	Neratinib+capecitabine sustains health-related quality of life in patients with HER2-positive metastatic breast cancer and2 prior HER2-directed regimens. <i>Breast Cancer Research and Treatment</i> , 2021, 182, 449-458.		2
85	Abstract CT331: a phase I/II study of the AKT inhibitor AZD5363 combined with paclitaxel in patients with advanced or metastatic breast cancer: results from the dose-finding study, including quantitative assessment of circulating tumor DNA as a s. , 2015, , .		2
86	Matching degree between PI3K/AKT/mTOR (PAM) pathway mutations (mut) and therapy (ttx) as predictor of clinical benefit (ClinBen) in early trials.. <i>Journal of Clinical Oncology</i> , 2016, 34, 2572-2572.	1.6	2
87	Prognostic and therapeutic implications of fibroblast growth factor receptors (FGFRs) 1 and 2 gene amplifications in patients (pts) with advanced breast cancer (ABC).. <i>Journal of Clinical Oncology</i> , 2016, 34, 537-537.	1.6	2
88	Determinants of concordance in clinically relevant genes (CRG) from synchronously acquired tumor biopsies (tBx) and ctDNA in metastatic breast cancer (MBC).. <i>Journal of Clinical Oncology</i> , 2019, 37, 1075-1075.	1.6	2
89	CONTESSA TRIO: A multinational, multicenter, phase (P) II study of tasetaxel (T) plus three different PD-(L)1 inhibitors in patients (Pts) with metastatic triple-negative breast cancer (TNBC) and tasetaxel monotherapy in elderly pts with HER2-metastatic breast cancer (MBC).. <i>Journal of Clinical Oncology</i> , 2020, 38, TPS1111-TPS1111.	1.6	2
90	Abstract 930: Analysis of cell-free tumor DNA in cerebrospinal fluid to characterize and monitor the genetic alterations of brain tumors. <i>Cancer Research</i> , 2015, 75, 930-930.	0.9	2

#	ARTICLE	IF	CITATIONS
91	P200 Neoadjuvant therapy in HER2+ breast cancer: Opti-HER Heart run-in phase safety data (SOLTI-1002). <i>Breast</i> , 2015, 24, S93.	2.2	1
92	Web Accessibility for Elderly. , 2016, , .		1
93	Reply to T. J. A. Dekker, D.-C. Mo et al, and A. Seidman et al. <i>Journal of Clinical Oncology</i> , 2021, 39, 254-255.	1.6	1
94	Abstract OT-09-02: A randomized, open-label, parallel-group, multicenter phase 2 study comparing the efficacy and safety of oral AZD9833 versus fulvestrant in women with advanced ER-positive HER2-negative breast cancer (SERENA-2). , 2021, , .		1
95	Abstract OT-09-08: Solti-1502 aRIANNA: Targeting PAM50 HER2-enriched intrinsic subtype with enzalutamide in hormone receptor-positive/HER2-negative metastatic breast cancer. , 2021, , .		1
96	P5-13-01: Survival Outcome with Bevacizumab: Activation of the Phosphatidylinositol-3 Kinase (PI3K) Pathway Due to PIK3CA Mutations or PTEN Loss Makes a Difference.. , 2011, , .		1
97	Abstract 3596: Biomarkers of response to CDK4/6 inhibitor (CDK4/6i) in hormone receptor (HR) positive and HER2-positive breast cancer (BC) patient-derived xenografts (PDX). , 2018, , .		1
98	A phase II trial of trabectedin (T) in patients with hormone receptor-positive, HER2-negative advanced breast cancer, according to xeroderma pigmentosum gene (XPG) expression.. <i>Journal of Clinical Oncology</i> , 2012, 30, TPS652-TPS652.	1.6	1
99	FAIRLANE: A phase II randomized, double-blind, study of the Akt inhibitor ipatasertib (Ipat, GDC-0068) in combination with paclitaxel (Pac) as neoadjuvant treatment for early stage triple-negative breast cancer (TNBC).. <i>Journal of Clinical Oncology</i> , 2015, 33, TPS1112-TPS1112.	1.6	1
100	PATRICIA: A phase II study of palbociclib and trastuzumab with or without letrozole in previously treated, postmenopausal patients with HER2-positive metastatic breast cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, TPS642-TPS642.	1.6	1
101	Final results of a phase II trial of trabectedin (T) in patients with hormone receptor-positive, HER2-negative advanced breast cancer, according to xeroderma pigmentosum gene (XPG) expression.. <i>Journal of Clinical Oncology</i> , 2013, 31, 547-547.	1.6	1
102	Abstract 2964: On-treatment changes in circulating tumor DNA (ctDNA) level as an early predictor of clinical outcome in the LOTUS randomized phase 2 trial of 1st-line ipatasertib (IPAT) + paclitaxel (PAC) for metastatic triple-negative breast cancer (mTNBC). <i>Cancer Research</i> , 2018, 78, 2964-2964.	0.9	1
103	Abstract P2-14-13: Talimogene laherparepvec (T-VEC) + atezolizumab combination in early breast cancer (SOLTI-1503 PROMETEO): Safety and efficacy interim analysis. <i>Cancer Research</i> , 2022, 82, P2-14-13-P2-14-13.	0.9	1
104	882 Evaluation of Synergy Between Novel PI3K-pathway Inhibitors and Microtubule-targeting Agents in HER2-negative Breast Cancer. <i>European Journal of Cancer</i> , 2012, 48, S213.	2.8	0
105	Intrathecal Trastuzumab in the Treatment of Leptomeningeal Metastases from Her2-Positive Cancer. <i>Annals of Oncology</i> , 2012, 23, ix141.	1.2	0
106	Triplet Combination of Endocrine Therapy with CDK 4/6 Inhibitor, Ribociclib, and MTOR Inhibitor, Everolimus in HR+, HER2-ABC: Results from the Dose-Expansion Cohort. <i>Breast</i> , 2017, 36, S46-S47.	2.2	0
107	Prognostic estimates of Ki-67 percentage drop after neoadjuvant chemotherapy (NAC) in luminal B (lumB) and triple negative breast cancer (TNBC). <i>Annals of Oncology</i> , 2017, 28, v68.	1.2	0
108	Primary results of the first nationwide molecular screening program in Spain for patients with advanced breast cancer (AGATA SOLTI-1301 study). <i>Annals of Oncology</i> , 2018, 29, viii90.	1.2	0

#	ARTICLE	IF	CITATIONS
109	ItRECIST adapted efficacy assessment in solid tumors treated with intratumoral immunotherapy.. Journal of Clinical Oncology, 2021, 39, 2557-2557.	1.6	0
110	Late toxicity and quality of life in oral cavity and oropharyngeal cancer survivors treated with chemoradiotherapy.. Journal of Clinical Oncology, 2010, 28, 9092-9092.	1.6	0
111	Prognostic significance of PI3K pathway (PI3Kp) dysregulation in metastatic breast cancer (MBC) patients (pts).. Journal of Clinical Oncology, 2012, 30, 566-566.	1.6	0
112	Analysis of the intratumoral heterogeneity of PIK3CA mutant alleles in breast cancer (BC): Implications for the luminal (LUM) phenotype.. Journal of Clinical Oncology, 2012, 30, 10511-10511.	1.6	0
113	Abstract P6-13-03: Symptomatic bone marrow involvement (BMinv) in breast cancer (BC): Clinical presentation, treatment and prognosis according to BC subtype and Zoledronic acid (ZA) use. A single institution review. , 2012, , .		0
114	PAM50 HER2-enriched (HER2E) phenotype as a predictor of early-response to neoadjuvant lapatinib plus trastuzumab in stage I to IIIA HER2-positive breast cancer.. Journal of Clinical Oncology, 2013, 31, TPS665-TPS665.	1.6	0
115	Abstract OT1-1-01: LORELEI: A Phase II randomized, double-blind study of neoadjuvant letrozole plus tasisib (GDC-0032) versus letrozole plus placebo in postmenopausal women with ER-positive/HER2-negative, early stage breast cancer. , 2015, , .		0
116	Abstract OT1-1-16: A randomized, multicenter, phase II study of ipatasertib (Ipat, GDC-0068), an inhibitor of Akt, in combination with paclitaxel (Pac) as front-line treatment for patients (pts) with metastatic triple-negative breast cancer (TNBC). , 2015, , .		0
117	Abstract 2399: Circulating tumor DNA (ctDNA) analysis of PIK3CA and AKT1 mutations in patients enrolled onto the Phase 1b study of the PI3K inhibitor tasisib (GDC-0032) in solid malignancies. , 2015, , .		0
118	Phase I evaluation of the PI3 kinase (PI3K) inhibitor tasisib (GDC-0032) in multiple locally advanced or metastatic <i>PIK3CA</i> mutant solid tumor types.. Journal of Clinical Oncology, 2016, 34, TPS11621-TPS11621.	1.6	0
119	FAIRLANE: A phase II randomized, double-blind, study of the Akt inhibitor ipatasertib (GDC-0068) in combination with paclitaxel as neoadjuvant treatment for early stage triple-negative breast cancer.. Journal of Clinical Oncology, 2016, 34, TPS1105-TPS1105.	1.6	0
120	Abstract 2825: Identification of CDK4/6-response biomarkers using estrogen receptor-positive breast cancer patient-derived xenografts (PDX). , 2016, , .		0
121	CORALLEEN: A phase 2 clinical trial of chemotherapy or letrozole plus ribociclib as neoadjuvant treatment for postmenopausal patients with luminal B/HER2-negative breast cancer.. Journal of Clinical Oncology, 2017, 35, TPS594-TPS594.	1.6	0
122	Abstract 150: Identification of determinants of sensitivity to AKT inhibition using breast cancer (BC) patient-derived tumor xenografts (PDX). , 2017, , .		0
123	Abstract 3129: Patient-derived tumor xenografts (PDXs) recapitulate the antitumor activity of novel therapies in metastatic breast cancer (MBC) patients (pts). , 2017, , .		0