

Thomas J Baum

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2031753/publications.pdf>

Version: 2024-02-01

102
papers

9,800
citations

41627

51
h-index

42259

96
g-index

108
all docs

108
docs citations

108
times ranked

5594
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Recognition and Response in Plant–Nematode Interactions. Annual Review of Phytopathology, 2022, 60, 143-162. | 3.5 | 23 |
| 2 | miR778 mediates gene expression, histone modification, and DNA methylation during cyst nematode parasitism. Plant Physiology, 2022, 189, 2432-2453. | 2.3 | 4 |
| 3 | Phytonematode peptide effectors exploit a host post-translational trafficking mechanism to the ER using a novel translocation signal. New Phytologist, 2021, 229, 563-574. | 3.5 | 24 |
| 4 | Toward genetic modification of plant-parasitic nematodes: delivery of macromolecules to adults and expression of exogenous mRNA in second stage juveniles. G3: Genes, Genomes, Genetics, 2021, 11, . | 0.8 | 9 |
| 5 | A chromosomal assembly of the soybean cyst nematode genome. Molecular Ecology Resources, 2021, 21, 2407-2422. | 2.2 | 10 |
| 6 | Esophageal Gland RNA-Seq Resource of a Virulent and Avirulent Population of the Soybean Cyst Nematode <i>Heterodera glycines</i> . Molecular Plant-Microbe Interactions, 2021, 34, 1084-1087. | 1.4 | 7 |
| 7 | Targeted transcriptomics reveals signatures of large-scale independent origins and concerted regulation of effector genes in <i>Radopholus similis</i> . PLoS Pathogens, 2021, 17, e1010036. | 2.1 | 2 |
| 8 | A role for Arabidopsis growth-regulating factors 1 and 3 in growth–stress antagonism. Journal of Experimental Botany, 2020, 71, 1402-1417. | 2.4 | 32 |
| 9 | Large tandem duplications affect gene expression, 3D organization, and plant–pathogen response. Genome Research, 2020, 30, 1583-1592. | 2.4 | 31 |
| 10 | Screening soybean cyst nematode effectors for their ability to suppress plant immunity. Molecular Plant Pathology, 2020, 21, 1240-1247. | 2.0 | 24 |
| 11 | Targeted suppression of soybean BAG6-induced cell death in yeast by soybean cyst nematode effectors. Molecular Plant Pathology, 2020, 21, 1227-1239. | 2.0 | 9 |
| 12 | A new esophageal gland transcriptome reveals signatures of large scale de novo effector birth in the root lesion nematode <i>Pratylenchus penetrans</i> . BMC Genomics, 2020, 21, 738. | 1.2 | 15 |
| 13 | SCNBase: a genomics portal for the soybean cyst nematode (<i>Heterodera glycines</i>). Database: the Journal of Biological Databases and Curation, 2019, 2019, . | 1.4 | 9 |
| 14 | <i>Heterodera glycines</i> utilizes promiscuous spliced leaders and demonstrates a unique preference for a species-specific spliced leader over <i>C. elegans</i> SL1. Scientific Reports, 2019, 9, 1356. | 1.6 | 5 |
| 15 | Homeostasis in the soybean miRNA396–GRF network is essential for productive soybean cyst nematode infections. Journal of Experimental Botany, 2019, 70, 1653-1668. | 2.4 | 27 |
| 16 | The genome of the soybean cyst nematode (<i>Heterodera glycines</i>) reveals complex patterns of duplications involved in the evolution of parasitism genes. BMC Genomics, 2019, 20, 119. | 1.2 | 55 |
| 17 | Re-targeting of a plant defense protease by a cyst nematode effector. Plant Journal, 2019, 98, 1000-1014. | 2.8 | 30 |
| 18 | Novel global effector mining from the transcriptome of early life stages of the soybean cyst nematode <i>Heterodera glycines</i> . Scientific Reports, 2018, 8, 2505. | 1.6 | 31 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Identification of candidate effector genes of <i>Pratylenchus penetrans</i> . <i>Molecular Plant Pathology</i> , 2018, 19, 1887-1907. | 2.0 | 36 |
| 20 | Suppression or Activation of Immune Responses by Predicted Secreted Proteins of the Soybean Rust Pathogen <i>Phakopsora pachyrhizi</i> . <i>Molecular Plant-Microbe Interactions</i> , 2018, 31, 163-174. | 1.4 | 54 |
| 21 | An Effector from the Cyst Nematode <i>Heterodera schachtii</i> Derepresses Host rRNA Genes by Altering Histone Acetylation. <i>Plant Cell</i> , 2018, 30, 2795-2812. | 3.1 | 30 |
| 22 | The plant-parasitic cyst nematode effector GLAND4 is a DNA-binding protein. <i>Molecular Plant Pathology</i> , 2018, 19, 2263-2276. | 2.0 | 31 |
| 23 | STATAWAARS: a promoter motif associated with spatial expression in the major effector-producing tissues of the plant-parasitic nematode <i>Bursaphelenchus xylophilus</i> . <i>BMC Genomics</i> , 2018, 19, 553. | 1.2 | 26 |
| 24 | The novel cyst nematode effector protein 30D08 targets host nuclear functions to alter gene expression in feeding sites. <i>New Phytologist</i> , 2018, 219, 697-713. | 3.5 | 38 |
| 25 | Research into <i>Heterodera</i> parasitism. <i>PLoS Pathogens</i> , 2018, 14, e1006791. | 2.1 | 26 |
| 26 | Cooperative Regulatory Functions of miR858 and MYB83 during Cyst Nematode Parasitism. <i>Plant Physiology</i> , 2017, 174, 1897-1912. | 2.3 | 46 |
| 27 | A <i>Plasmodium</i> -like virulence effector of the soybean cyst nematode suppresses plant innate immunity. <i>New Phytologist</i> , 2016, 212, 444-460. | 3.5 | 47 |
| 28 | <i>Arabidopsis</i> miR827 mediates posttranscriptional gene silencing of its ubiquitin E3 ligase target gene in the syncytium of the cyst nematode <i>Heterodera schachtii</i> to enhance susceptibility. <i>Plant Journal</i> , 2016, 88, 179-192. | 2.8 | 65 |
| 29 | A cyst nematode effector binds to diverse plant proteins, increases nematode susceptibility and affects root morphology. <i>Molecular Plant Pathology</i> , 2016, 17, 832-844. | 2.0 | 32 |
| 30 | Horizontal gene transfer of acetyltransferases, invertases and chorismate mutases from different bacteria to diverse recipients. <i>BMC Evolutionary Biology</i> , 2016, 16, 74. | 3.2 | 19 |
| 31 | A Small Cysteine-Rich Protein from the Asian Soybean Rust Fungus, <i>Phakopsora pachyrhizi</i> , Suppresses Plant Immunity. <i>PLoS Pathogens</i> , 2016, 12, e1005827. | 2.1 | 79 |
| 32 | The Cyst Nematode Effector Protein 10A07 Targets and Recruits Host Posttranslational Machinery to Mediate Its Nuclear Trafficking and to Promote Parasitism in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2015, 27, 891-907. | 3.1 | 84 |
| 33 | Gene Silencing in Nematode Feeding Sites. <i>Advances in Botanical Research</i> , 2015, 73, 221-239. | 0.5 | 12 |
| 34 | Eighteen New Candidate Effectors of the Phytonematode <i>Heterodera glycines</i> Produced Specifically in the Secretory Esophageal Gland Cells During Parasitism. <i>Phytopathology</i> , 2015, 105, 1362-1372. | 1.1 | 57 |
| 35 | Sequence and Spatiotemporal Expression Analysis of CLE-Motif Containing Genes from the Reniform Nematode (<i>Rotylenchulus reniformis</i> Linford & Oliveira). <i>Journal of Nematology</i> , 2015, 47, 159-65. | 0.4 | 13 |
| 36 | Synchronization of Developmental Processes and Defense Signaling by Growth Regulating Transcription Factors. <i>PLoS ONE</i> , 2014, 9, e98477. | 1.1 | 76 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 37 | A virus-induced gene silencing method to study soybean cyst nematode parasitism in <i>Glycine max.</i> BMC Research Notes, 2013, 6, 255. | 0.6 | 28 |
| 38 | Transcriptome analyses and virus induced gene silencing identify genes in the Rpp4-mediated Asian soybean rust resistance pathway. Functional Plant Biology, 2013, 40, 1029. | 1.1 | 57 |
| 39 | Manipulation of Plant Cells by Cyst and Root-Knot Nematode Effectors. Molecular Plant-Microbe Interactions, 2013, 26, 9-16. | 1.4 | 184 |
| 40 | Isolation of Whole Esophageal Gland Cells from Plant-Parasitic Nematodes for Transcriptome Analyses and Effector Identification. Molecular Plant-Microbe Interactions, 2013, 26, 31-35. | 1.4 | 56 |
| 41 | The <i>8D05</i> Parasitism Gene of <i>Meloidogyne incognita</i> Is Required for Successful Infection of Host Roots. Phytopathology, 2013, 103, 175-181. | 1.1 | 86 |
| 42 | Nematode effector proteins: an emerging paradigm of parasitism. New Phytologist, 2013, 199, 879-894. | 3.5 | 269 |
| 43 | A ubiquitin carboxyl extension protein secreted from a plant-parasitic nematode <i>Globodera rostochiensis</i> is cleaved <i>in planta</i> to promote plant parasitism. Plant Journal, 2013, 74, 185-196. | 2.8 | 98 |
| 44 | Complex feedback regulations govern the expression of miRNA396 and its GRF target genes. Plant Signaling and Behavior, 2012, 7, 749-751. | 1.2 | 52 |
| 45 | The Arabidopsis MicroRNA396-GRF1/GRF3 Regulatory Module Acts as a Developmental Regulator in the Reprogramming of Root Cells during Cyst Nematode Infection. Plant Physiology, 2012, 159, 321-335. | 2.3 | 214 |
| 46 | The interaction of the novel 30C02 cyst nematode effector protein with a plant β -1,3-endoglucanase may suppress host defence to promote parasitism. Journal of Experimental Botany, 2012, 63, 3683-3695. | 2.4 | 80 |
| 47 | A soybean cyst nematode resistance gene points to a new mechanism of plant resistance to pathogens. Nature, 2012, 492, 256-260. | 13.7 | 332 |
| 48 | Temporal and spatial <i>Bean pod mottle virus</i> -induced gene silencing in soybean. Molecular Plant Pathology, 2012, 13, 1140-1148. | 2.0 | 19 |
| 49 | The Arabidopsis bHLH25 and bHLH27 transcription factors contribute to susceptibility to the cyst nematode <i>Heterodera schachtii</i> . Plant Journal, 2011, 65, 319-328. | 2.8 | 40 |
| 50 | Nematode CLE signaling in Arabidopsis requires CLAVATA2 and CORYNE. Plant Journal, 2011, 65, 430-440. | 2.8 | 108 |
| 51 | Identification of potential host plant mimics of CLAVATA3/ESR (CLE)-like peptides from the plant-parasitic nematode <i>Heterodera schachtii</i> . Molecular Plant Pathology, 2011, 12, 177-186. | 2.0 | 95 |
| 52 | Arabidopsis peroxidase AtPRX53 influences cell elongation and susceptibility to <i>Heterodera schachtii</i> . Plant Signaling and Behavior, 2011, 6, 1778-1786. | 1.2 | 30 |
| 53 | The Novel Cyst Nematode Effector Protein 19C07 Interacts with the Arabidopsis Auxin Influx Transporter LAX3 to Control Feeding Site Development. Plant Physiology, 2011, 155, 866-880. | 2.3 | 141 |
| 54 | The Soybean <i>Rhg1</i> Locus for Resistance to the Soybean Cyst Nematode <i>Heterodera glycines</i> Regulates the Expression of a Large Number of Stress- and Defense-Related Genes in Degenerating Feeding Cells. Plant Physiology, 2011, 155, 1960-1975. | 2.3 | 102 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 55 | Dual roles for the variable domain in protein trafficking and host-specific recognition of <i>Heterodera glycines</i> CLE effector proteins. <i>New Phytologist</i> , 2010, 187, 1003-1017. | 3.5 | 116 |
| 56 | <i>Arabidopsis</i> Spermidine Synthase Is Targeted by an Effector Protein of the Cyst Nematode <i>Heterodera schachtii</i> . <i>Plant Physiology</i> , 2010, 152, 968-984. | 2.3 | 189 |
| 57 | A nematode effector protein similar to annexins in host plants. <i>Journal of Experimental Botany</i> , 2010, 61, 235-248. | 2.4 | 114 |
| 58 | Sequence divergences between cyst nematode effector protein orthologs may contribute to host specificity. <i>Plant Signaling and Behavior</i> , 2010, 5, 187-189. | 1.2 | 5 |
| 59 | Effective and specific in planta RNAi in cyst nematodes: expression interference of four parasitism genes reduces parasitic success. <i>Journal of Experimental Botany</i> , 2009, 60, 315-324. | 2.4 | 144 |
| 60 | Parasitism Genes: What They Reveal about Parasitism. <i>Plant Cell Monographs</i> , 2009, , 15-44. | 0.4 | 12 |
| 61 | Sequence mining and transcript profiling to explore cyst nematode parasitism. <i>BMC Genomics</i> , 2009, 10, 58. | 1.2 | 43 |
| 62 | Genome sequence of the metazoan plant-parasitic nematode <i>Meloidogyne incognita</i> . <i>Nature Biotechnology</i> , 2008, 26, 909-915. | 9.4 | 1,012 |
| 63 | Parasitism proteins in nematode-plant interactions. <i>Current Opinion in Plant Biology</i> , 2008, 11, 360-366. | 3.5 | 223 |
| 64 | Cellulose Binding Protein from the Parasitic Nematode <i>Heterodera schachtii</i> Interacts with <i>Arabidopsis</i> Pectin Methyltransferase: Cooperative Cell Wall Modification during Parasitism. <i>Plant Cell</i> , 2008, 20, 3080-3093. | 3.1 | 201 |
| 65 | <i>Arabidopsis</i> Small RNAs and Their Targets During Cyst Nematode Parasitism. <i>Molecular Plant-Microbe Interactions</i> , 2008, 21, 1622-1634. | 1.4 | 124 |
| 66 | Genomics of the Soybean Cyst Nematode-Soybean Interaction. , 2008, , 321-341. | | 7 |
| 67 | GmERE1BP1 Is a Transcription Factor Activating Defense Genes in Soybean and <i>Arabidopsis</i> . <i>Molecular Plant-Microbe Interactions</i> , 2007, 20, 107-119. | 1.4 | 78 |
| 68 | Parallel Genome-Wide Expression Profiling of Host and Pathogen During Soybean Cyst Nematode Infection of Soybean. <i>Molecular Plant-Microbe Interactions</i> , 2007, 20, 293-305. | 1.4 | 197 |
| 69 | Developmental Transcript Profiling of Cyst Nematode Feeding Cells in Soybean Roots. <i>Molecular Plant-Microbe Interactions</i> , 2007, 20, 510-525. | 1.4 | 240 |
| 70 | Divergent evolution of arrested development in the dauer stage of <i>Caenorhabditis elegans</i> and the infective stage of <i>Heterodera glycines</i> . <i>Genome Biology</i> , 2007, 8, R211. | 13.9 | 40 |
| 71 | Root-Knot and Cyst Nematode Parasitism Genes: The Molecular Basis of Plant Parasitism. , 2007, 28, 17-43. | | 49 |
| 72 | Active uptake of cyst nematode parasitism proteins into the plant cell nucleus. <i>International Journal for Parasitology</i> , 2007, 37, 1269-1279. | 1.3 | 73 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Quantitative Detection of Double-Stranded RNA-Mediated Gene Silencing of Parasitism Genes in <i>Heterodera glycines</i> . <i>Journal of Nematology</i> , 2007, 39, 145-52. | 0.4 | 25 |
| 74 | Engineering broad root-knot resistance in transgenic plants by RNAi silencing of a conserved and essential root-knot nematode parasitism gene. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 14302-14306. | 3.3 | 543 |
| 75 | A Root-Knot Nematode Secretory Peptide Functions as a Ligand for a Plant Transcription Factor. <i>Molecular Plant-Microbe Interactions</i> , 2006, 19, 463-470. | 1.4 | 189 |
| 76 | Two chorismate mutase genes from the root-knot nematode <i>Meloidogyne incognita</i> . <i>Molecular Plant Pathology</i> , 2005, 6, 23-30. | 2.0 | 66 |
| 77 | A parasitism gene from a plant-parasitic nematode with function similar to <i>CLAVATA3/ESR (CLE)</i> of <i>Arabidopsis thaliana</i> . <i>Molecular Plant Pathology</i> , 2005, 6, 187-191. | 2.0 | 215 |
| 78 | Developmental expression and molecular analysis of two <i>Meloidogyne incognita</i> pectate lyase genes. <i>International Journal for Parasitology</i> , 2005, 35, 685-692. | 1.3 | 63 |
| 79 | Mutation of a UDP-glucose-4-epimerase alters nematode susceptibility and ethylene responses in <i>Arabidopsis</i> roots. <i>Plant Journal</i> , 2004, 40, 712-724. | 2.8 | 47 |
| 80 | Use of solid-phase subtractive hybridization for the identification of parasitism gene candidates from the root-knot nematode <i>Meloidogyne incognita</i> . <i>Molecular Plant Pathology</i> , 2004, 5, 217-222. | 2.0 | 48 |
| 81 | Homologous soybean and <i>Arabidopsis</i> genes share responsiveness to cyst nematode infection. <i>Molecular Plant Pathology</i> , 2004, 5, 409-423. | 2.0 | 16 |
| 82 | Getting to the roots of parasitism by nematodes. <i>Trends in Parasitology</i> , 2004, 20, 134-141. | 1.5 | 273 |
| 83 | Molecular characterisation and developmental expression of a cellulose-binding protein gene in the soybean cyst nematode <i>Heterodera glycines</i> . <i>International Journal for Parasitology</i> , 2004, 34, 1377-1383. | 1.3 | 40 |
| 84 | Expression of an <i>Arabidopsis</i> phosphoglycerate mutase homologue is localized to apical meristems, regulated by hormones, and induced by sedentary plant-parasitic nematodes. <i>Plant Molecular Biology</i> , 2003, 53, 513-530. | 2.0 | 85 |
| 85 | <i>Arabidopsis</i> gene expression changes during cyst nematode parasitism revealed by statistical analyses of microarray expression profiles. <i>Plant Journal</i> , 2003, 33, 911-921. | 2.8 | 180 |
| 86 | The Parasitome of the Phytonematode <i>Heterodera glycines</i> . <i>Molecular Plant-Microbe Interactions</i> , 2003, 16, 720-726. | 1.4 | 257 |
| 87 | A Profile of Putative Parasitism Genes Expressed in the Esophageal Gland Cells of the Root-knot Nematode <i>Meloidogyne incognita</i> . <i>Molecular Plant-Microbe Interactions</i> , 2003, 16, 376-381. | 1.4 | 211 |
| 88 | Identification and Characterization of a Soybean Ethylene-Responsive Element-Binding Protein Gene Whose mRNA Expression Changes During Soybean Cyst Nematode Infection. <i>Molecular Plant-Microbe Interactions</i> , 2002, 15, 577-586. | 1.4 | 64 |
| 89 | Secrets in secretions: genes that control nematode parasitism of plants. <i>Brazilian Journal of Plant Physiology</i> , 2002, 14, 183-194. | 0.5 | 59 |
| 90 | Characterisation and developmental expression of a chitinase gene in <i>Heterodera glycines</i> . <i>International Journal for Parasitology</i> , 2002, 32, 1293-1300. | 1.3 | 43 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | The use of DNA microarrays for the developmental expression analysis of cDNAs from the oesophageal gland cell region of <i>Heterodera glycines</i> . <i>Molecular Plant Pathology</i> , 2002, 3, 261-270. | 2.0 | 25 |
| 92 | Susceptibility to the Sugar Beet Cyst Nematode Is Modulated by Ethylene Signal Transduction in <i>Arabidopsis thaliana</i> . <i>Molecular Plant-Microbe Interactions</i> , 2001, 14, 1206-1212. | 1.4 | 134 |
| 93 | Identification of Putative Parasitism Genes Expressed in the Esophageal Gland Cells of the Soybean Cyst Nematode <i>Heterodera glycines</i> . <i>Molecular Plant-Microbe Interactions</i> , 2001, 14, 1247-1254. | 1.4 | 107 |
| 94 | Signal Peptide-Selection of cDNA Cloned Directly from the Esophageal Gland Cells of the Soybean Cyst Nematode <i>Heterodera glycines</i> . <i>Molecular Plant-Microbe Interactions</i> , 2001, 14, 536-544. | 1.4 | 156 |
| 95 | Molecular characterisation and expression of two venom allergen-like protein genes in <i>Heterodera glycines</i> . <i>International Journal for Parasitology</i> , 2001, 31, 1617-1625. | 1.3 | 75 |
| 96 | Changes in mRNA Abundance within <i>Heterodera schachtii</i> -Infected Roots of <i>Arabidopsis thaliana</i> . <i>Molecular Plant-Microbe Interactions</i> , 2000, 13, 309-315. | 1.4 | 46 |
| 97 | Nematode Parasitism Genes. <i>Annual Review of Phytopathology</i> , 2000, 38, 365-396. | 3.5 | 270 |
| 98 | Isolation of a cDNA Encoding a β -1,4-endoglucanase in the Root-Knot Nematode <i>Meloidogyne incognita</i> and Expression Analysis During Plant Parasitism. <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 585-591. | 1.4 | 188 |
| 99 | Developmental Expression of Secretory β -1,4-endoglucanases in the Subventral Esophageal Glands of <i>Heterodera glycines</i> . <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 663-669. | 1.4 | 87 |
| 100 | In Planta Localization of a β -1,4-Endoglucanase Secreted by <i>Heterodera glycines</i> . <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 64-67. | 1.4 | 113 |
| 101 | Genomic organization of four β -1,4-endoglucanase genes in plant-parasitic cyst nematodes and its evolutionary implications. <i>Gene</i> , 1998, 220, 61-70. | 1.0 | 128 |
| 102 | Differential Display Analysis of the Early Compatible Interaction Between Soybean and the Soybean Cyst Nematode. <i>Molecular Plant-Microbe Interactions</i> , 1998, 11, 1258-1263. | 1.4 | 52 |