## Jan Skupien

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2057688/publications.pdf

Version: 2024-02-01

		361413	377865
35	2,515	20	34
papers	citations	h-index	g-index
36	36	36	3304
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Circulating TNF Receptors 1 and 2 Predict ESRD in Type 2 Diabetes. Journal of the American Society of Nephrology: JASN, 2012, 23, 507-515.	6.1	388
2	Circulating TNF Receptors 1 and 2 Predict Stage 3 CKD in Type 1 Diabetes. Journal of the American Society of Nephrology: JASN, 2012, 23, 516-524.	6.1	307
3	A signature of circulating inflammatory proteins and development of end-stage renal disease in diabetes. Nature Medicine, 2019, 25, 805-813.	30.7	260
4	Early Progressive Renal Decline Precedes the Onset of Microalbuminuria and Its Progression to Macroalbuminuria. Diabetes Care, 2014, 37, 226-234.	8.6	219
5	Uremic solutes and risk of end-stage renal disease in type 2 diabetes: metabolomic study. Kidney International, 2014, 85, 1214-1224.	5.2	182
6	Risk for ESRD in Type 1 Diabetes Remains High Despite Renoprotection. Journal of the American Society of Nephrology: JASN, 2011, 22, 545-553.	6.1	166
7	Fast renal decline to end-stage renal disease: an unrecognized feature of nephropathy in diabetes. Kidney International, 2017, 91, 1300-1311.	5.2	159
8	The early decline in renal function in patients with type 1 diabetes and proteinuria predicts the risk of end-stage renal disease. Kidney International, 2012, 82, 589-597.	5.2	120
9	Increased plasma kidney injury molecule-1 suggests early progressive renal decline in non-proteinuric patients with type 1 diabetes. Kidney International, 2016, 89, 459-467.	5.2	101
10	Markers of early progressive renal decline in typeÂ2Âdiabetes suggest different implications forÂetiological studies and prognostic testsÂdevelopment. Kidney International, 2018, 93, 1198-1206.	5.2	88
11	Risk of macrosomia remains glucose-dependent in a cohort of women with pregestational type 1 diabetes and good glycemic control. Endocrine, 2017, 55, 447-455.	2.3	58
12	Patterns of Estimated Glomerular Filtration Rate Decline Leading to End-Stage Renal Disease in Type 1 Diabetes. Diabetes Care, 2016, 39, 2262-2269.	8.6	46
13	Synergism Between Circulating Tumor Necrosis Factor Receptor 2 and HbA1c in Determining Renal Decline During 5–18 Years of Follow-up in Patients With Type 1 Diabetes and Proteinuria. Diabetes Care, 2014, 37, 2601-2608.	8.6	43
14	Improved Glycemic Control and Risk of ESRD in Patients with Type 1 Diabetes and Proteinuria. Journal of the American Society of Nephrology: JASN, 2014, 25, 2916-2925.	6.1	39
15	Improved clinical trial enrollment criterion toÂidentify patients with diabetes at risk of end-stage renal disease. Kidney International, 2017, 92, 258-266.	5.2	38
16	Variations in Risk of End-Stage Renal Disease and Risk of Mortality in an International Study of Patients With Type 1 Diabetes and Advanced Nephropathy. Diabetes Care, 2019, 42, 93-101.	8.6	37
17	Evaluation of Apolipoprotein M Serum Concentration as a Biomarker of HNF-1alpha MODY. Review of Diabetic Studies, 2007, 4, 231-235.	1.3	30
18	Clinical Application of 1,5-Anhydroglucitol Measurements in Patients with Hepatocyte Nuclear Factor-1α Maturity-Onset Diabetes of the Young. Diabetes Care, 2008, 31, 1496-1501.	8.6	29

#	Article	IF	Citations
19	Mortality Following Clostridioides difficile Infection in Europe: A Retrospective Multicenter Case-Control Study. Antibiotics, 2021, 10, 299.	3.7	23
20	Profibrotic Circulating Proteins and Risk of Early Progressive Renal Decline in Patients With Type 2 Diabetes With and Without Albuminuria. Diabetes Care, 2020, 43, 2760-2767.	8.6	21
21	Genetic testing for monogenic diabetes using targeted next-generation sequencing in patients with maturity-onset diabetes of the young. Polish Archives of Internal Medicine, 2015, 125, 845-851.	0.4	21
22	Diabetic pregnancy. Current Opinion in Obstetrics and Gynecology, 2014, 26, 431-437.	2.0	20
23	A family with the Arg103Pro mutation in the NEUROD1 gene detected by next-generation sequencing – Clinical characteristics of mutation carriers. European Journal of Medical Genetics, 2016, 59, 75-79.	1.3	19
24	Circulating ghrelin level is higher in HNF1A–MODY and GCK–MODY than in polygenic forms of diabetes mellitus. Endocrine, 2015, 50, 643-649.	2.3	18
25	Circulating proteins protect against renal decline and progression to end-stage renal disease in patients with diabetes. Science Translational Medicine, 2021, 13, .	12.4	18
26	Hypoglycemic episodes are associated with inflammatory status in patients with type 1 diabetes mellitus. Atherosclerosis, 2016, 251, 334-338.	0.8	17
27	A decision algorithm to identify patients with high probability of monogenic diabetes due to HNF1A mutations. Endocrine, 2019, 64, 75-81.	2.3	12
28	Assessment of Insulin Sensitivity in Adults with Permanent Neonatal Diabetes Mellitus due to Mutations in the KCNJ11 Gene Encoding Kir6.2. Review of Diabetic Studies, 2006, 3, 17-17.	1.3	10
29	Analysis of gene expression to predict dynamics of future hypertension incidence in type 2 diabetic patients. BMC Proceedings, 2016, 10, 113-117.	1.6	9
30	Increased Incidence of Type 1 Diabetes in Children and No Change in the Age of Diagnosis and BMI-SDS at the Onset - is the Accelerator Hypothesis not Working?. JCRPE Journal of Clinical Research in Pediatric Endocrinology, 2020, 12, 281-286.	0.9	8
31	Risk factors of hypoglycaemia in type 1 diabetes individuals during intensive sport exercise—Data from the SPORTGIVECHANCE event. International Journal of Clinical Practice, 2019, 73, e13411.	1.7	3
32	Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase p22phox subunit polymorphisms, systemic oxidative stress, endothelial dysfunction, and atherosclerosis in type 2 diabetes mellitus. Polish Archives of Internal Medicine, 2021, 131, 447-454.	0.4	3
33	Increased Incidence of Type 1 Diabetes in Children and No Change in the Age of Diagnosis and BMI-SDS at the Onset - is the Accelerator Hypothesis not Working?. JCRPE Journal of Clinical Research in Pediatric Endocrinology, 2020, 12, 281-286.	0.9	2
34	COVID-19 Infection Negative in Nasopharyngeal Swabs but Suspected in Computed Tomography and Confirmed in Bronchoalveolar Lavage Material. Case Reports in Infectious Diseases, 2021, 2021, 1-3.	0.5	1
35	Analytical Evaluation of the Xpress and Hospital Blood Glucose Monitoring Systems. Journal of Diabetes Science and Technology, 2014, 8, 434-435.	2.2	0