

Hideki Wanibuchi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2089634/publications.pdf>

Version: 2024-02-01

124
papers

3,165
citations

147801

31
h-index

189892

50
g-index

126
all docs

126
docs citations

126
times ranked

2994
citing authors

#	ARTICLE	IF	CITATIONS
1	FOXP3 and CXCR4-positive regulatory T cells in the tumor stroma as indicators of tumor immunity in the conjunctival squamous cell carcinoma microenvironment. PLoS ONE, 2022, 17, e0263895.	2.5	1
2	The carbonic anhydrase inhibitor acetazolamide inhibits urinary bladder cancers via suppression of β -catenin signaling. Cancer Science, 2022, 113, 2642-2653.	3.9	3
3	Cache Domain Containing 1 Is a Novel Marker of Non-Alcoholic Steatohepatitis-Associated Hepatocarcinogenesis. Cancers, 2021, 13, 1216.	3.7	5
4	Expression of thrombospondin-1 in conjunctival squamous cell carcinoma is correlated to the Ki67 index and associated with progression-free survival. Graefe's Archive for Clinical and Experimental Ophthalmology, 2021, 259, 3127-3136.	1.9	1
5	ER membrane protein complex 1 interacts with STIM1 and regulates store-operated Ca^{2+} entry. Journal of Biochemistry, 2021, 170, 483-488.	1.7	4
6	Canopy Homolog 2 as a Novel Molecular Target in Hepatocarcinogenesis. Cancers, 2021, 13, 3613.	3.7	4
7	Accumulation of 8-hydroxydeoxyguanosine, L-arginine and Glucose Metabolites by Liver Tumor Cells Are the Important Characteristic Features of Metabolic Syndrome and Non-Alcoholic Steatohepatitis-Associated Hepatocarcinogenesis. International Journal of Molecular Sciences, 2020, 21, 7746.	4.1	17
8	Expression, intracellular localization, and mutation of EGFR in conjunctival squamous cell carcinoma and the association with prognosis and treatment. PLoS ONE, 2020, 15, e0238120.	2.5	6
9	Myeloid-derived suppressor cells are essential partners for immune checkpoint inhibitors in the treatment of cisplatin-resistant bladder cancer. Cancer Letters, 2020, 479, 89-99.	7.2	36
10	Dimethylarsinic acid (DMA) enhanced lung carcinogenesis via histone H3K9 modification in a transplacental mouse model. Archives of Toxicology, 2020, 94, 927-937.	4.2	12
11	Comprehensive analysis of DNA adducts (DNA adductome analysis) in the liver of rats treated with 1,4-dioxane. Proceedings of the Japan Academy Series B: Physical and Biological Sciences, 2020, 96, 180-187.	3.8	14
12	Title is missing!. , 2020, 15, e0238120.		0
13	Title is missing!. , 2020, 15, e0238120.		0
14	Title is missing!. , 2020, 15, e0238120.		0
15	Title is missing!. , 2020, 15, e0238120.		0
16	Title is missing!. , 2020, 15, e0238120.		0
17	Title is missing!. , 2020, 15, e0238120.		0
18	PITX1 protein interacts with ZCCHC10 to regulate hTERT mRNA transcription. PLoS ONE, 2019, 14, e0217605.	2.5	21

#	ARTICLE	IF	CITATIONS
19	Promotion effects of acetoaceto-o-toluidide on N-butyl-N-(4-hydroxybutyl)nitrosamine-induced bladder carcinogenesis in rats. Archives of Toxicology, 2019, 93, 3617-3631.	4.2	4
20	A chronic toxicity study of diphenylarsinic acid in the drinking water of C57BL/6J mice for 52 weeks. Journal of Toxicologic Pathology, 2019, 32, 127-134.	0.7	2
21	Acetoaceto-o-Toluidide Enhances Cellular Proliferative Activity in the Urinary Bladder of Rats. Toxicological Sciences, 2019, 169, 456-464.	3.1	7
22	mTOR Activation in Liver Tumors Is Associated with Metabolic Syndrome and Non-Alcoholic Steatohepatitis in Both Mouse Models and Humans. Cancers, 2018, 10, 465.	3.7	12
23	Steroid sulfatase promotes invasion through epithelialâ€mesenchymal transition and predicts the progression of bladder cancer. Experimental and Therapeutic Medicine, 2018, 16, 4463-4470.	1.8	1
24	Chronic dietary toxicity and carcinogenicity studies of dammar resin in F344 rats. Archives of Toxicology, 2018, 92, 3565-3583.	4.2	1
25	Generation of Rat Monoclonal Antibodies Against a Deubiquitinase, Ovarian Tumor Domain-Containing Protein 1. Monoclonal Antibodies in Immunodiagnosis and Immunotherapy, 2018, 37, 180-184.	1.6	2
26	Hypoxia-inducible factor-2 alpha up-regulates CD70 under hypoxia and enhances anchorage-independent growth and aggressiveness in cancer cells. Oncotarget, 2018, 9, 19123-19135.	1.8	21
27	Carbonic anhydrase 2 is a novel invasionâ€associated factor in urinary bladder cancers. Cancer Science, 2017, 108, 331-337.	3.9	12
28	PIK3CA mutation as a distinctive genetic feature of non-small cell lung cancer with chronic obstructive pulmonary disease: A comprehensive mutational analysis from a multi-institutional cohort. Lung Cancer, 2017, 112, 96-101.	2.0	17
29	A chronic toxicity study of diphenylarsinic acid in F344 rats in drinking water for 52 weeks. Experimental and Toxicologic Pathology, 2017, 69, 1-7.	2.1	5
30	Enhanced Susceptibility of Ogg1 Mutant Mice to Multiorgan Carcinogenesis. International Journal of Molecular Sciences, 2017, 18, 1801.	4.1	16
31	Proteome Characteristics of Non-Alcoholic Steatohepatitis Liver Tissue and Associated Hepatocellular Carcinomas. International Journal of Molecular Sciences, 2017, 18, 434.	4.1	20
32	Progression of Hepatic Adenoma to Carcinoma in <i>Ogg1</i> Mutant Mice Induced by Phenobarbital. Oxidative Medicine and Cellular Longevity, 2017, 2017, 1-16.	4.0	9
33	A carcinogenicity study of diphenylarsinic acid in F344 rats in drinking water for 104 weeks. Journal of Toxicological Sciences, 2017, 42, 475-483.	1.5	2
34	Chemopreventive Action by Ethanol-extracted Brazilian Green Propolis on Post-initiation Phase of Inflammation-associated Rat Colon Tumorigenesis. In Vivo, 2017, 31, 187-198.	1.3	24
35	Pueraria mirifica Exerts Estrogenic Effects in the Mammary Gland and Uterus and Promotes Mammary Carcinogenesis in Donryu Rats. Toxins, 2016, 8, 275.	3.4	9
36	Antiâ€PDâ€L1 treatment enhances antitumor effect of everolimus in a mouse model of renal cell carcinoma. Cancer Science, 2016, 107, 1736-1744.	3.9	56

#	ARTICLE	IF	CITATIONS
37	Examination of in vivo mutagenicity of sodium arsenite and dimethylarsinic acid in gpt delta rats. <i>Journal of Environmental Sciences</i> , 2016, 49, 125-130.	6.1	6
38	Alteration of Esophageal Microbiome by Antibiotic Treatment Does Not Affect Incidence of Rat Esophageal Adenocarcinoma. <i>Digestive Diseases and Sciences</i> , 2016, 61, 3161-3168.	2.3	19
39	Role of deltaNp63 pos CD 44v pos cells in the development of N-nitrosoethylurea-induced peripheral-type mouse lung squamous cell carcinomas. <i>Cancer Science</i> , 2016, 107, 123-132.	3.9	27
40	<scp>CD</scp>44 variant 9 is a potential biomarker of tumor initiating cells predicting survival outcome in hepatitis C virus-positive patients with resected hepatocellular carcinoma. <i>Cancer Science</i> , 2016, 107, 609-618.	3.9	34
41	Detection of non-genotoxic hepatocarcinogens and prediction of their mechanism of action in rats using gene marker sets. <i>Journal of Toxicological Sciences</i> , 2016, 41, 281-292.	1.5	8
42	Qualitative and Quantitative Assessments on Low-Dose Carcinogenicity of Genotoxic Hepatocarcinogens. , 2016, , 1-17.		10
43	Ethanol-Extracted Brazilian Propolis Exerts Protective Effects on Tumorigenesis in Wistar Hannover Rats. <i>PLoS ONE</i> , 2016, 11, e0158654.	2.5	17
44	Modifying effects of 1,2-dichloropropane on N-nitrosobis(2-oxopropyl)amine-induced cholangiocarcinogenesis in male Syrian hamsters. <i>Journal of Toxicological Sciences</i> , 2015, 40, 647-656.	1.5	5
45	Induction of cell proliferation in the rat liver by the short-term administration of ethyl & tertiary-butyl ether. <i>Journal of Toxicologic Pathology</i> , 2015, 28, 27-32.	0.7	8
46	Gene-modified embryonic stem cell test to characterize chemical risks. <i>Environmental Science and Pollution Research</i> , 2015, 22, 18252-18259.	5.3	0
47	Integrative analyses of miRNA and proteomics identify potential biological pathways associated with onset of pulmonary fibrosis in the bleomycin rat model. <i>Toxicology and Applied Pharmacology</i> , 2015, 286, 188-197.	2.8	14
48	Ethanol-extracted propolis enhances BBN-initiated urinary bladder carcinogenesis via non-mutagenic mechanisms in rats. <i>Food and Chemical Toxicology</i> , 2015, 83, 193-200.	3.6	7
49	Determination of Hepatotoxicity and Its Underlying Metabolic Basis of 1,2-Dichloropropane in Male Syrian Hamsters and B6C3F1 Mice. <i>Toxicological Sciences</i> , 2015, 145, 196-208.	3.1	9
50	Roles of Leucine and Isoleucine in Experimental Models of Bladder Carcinogenesis. <i>Food Safety (Tokyo)</i> , Tj ETQq0 0,0rgBT /Overlock 10	1.8	1
51	Isoleucine, Leucine and Their Role in Experimental Models of Bladder Carcinogenesis. , 2015, , 253-260.		1
52	Valerian Inhibits Rat Hepatocarcinogenesis by Activating GABA(A) Receptor-Mediated Signaling. <i>PLoS ONE</i> , 2014, 9, e113610.	2.5	11
53	Comparative Proteomics Analysis of Gastric Cancer Stem Cells. <i>PLoS ONE</i> , 2014, 9, e110736.	2.5	39
54	Inhibitory effect of raphanobrassica on Helicobacter pylori-induced gastritis in Mongolian gerbils. <i>Food and Chemical Toxicology</i> , 2014, 70, 107-113.	3.6	8

#	ARTICLE	IF	CITATIONS
55	l-Leucine and l-Isoleucine enhance growth of BBN-induced urothelial tumors in the rat bladder by modulating expression of amino acid transporters and tumorigenesis-associated genes. <i>Food and Chemical Toxicology</i> , 2013, 59, 137-144.	3.6	19
56	Diphenylarsinic acid, a chemical warfare-related neurotoxicant, promotes liver carcinogenesis via activation of aryl hydrocarbon receptor signaling and consequent induction of oxidative DNA damage in rats. <i>Toxicology and Applied Pharmacology</i> , 2013, 273, 1-9.	2.8	11
57	Myristoylated alanine-rich C-kinase substrate as a prognostic biomarker in human primary lung squamous cell carcinoma. <i>Cancer Biomarkers</i> , 2013, 13, 289-298.	1.7	30
58	Mode of action of ethyl tertiary-butyl ether hepatotumorigenicity in the rat: Evidence for a role of oxidative stress via activation of CAR, PXR and PPAR signaling pathways. <i>Toxicology and Applied Pharmacology</i> , 2013, 273, 390-400.	2.8	29
59	Oxidative Stress in the Carcinogenicity of Chemical Carcinogens. <i>Cancers</i> , 2013, 5, 1332-1354.	3.7	39
60	Evaluation of the Modifying Effect of Inhalation of Mainstream Cigarette Smoke on Mouse Bladder Carcinogenesis. <i>Journal of Toxicologic Pathology</i> , 2013, 26, 447-451.	0.7	2
61	Novel medium-term carcinogenesis model for lung squamous cell carcinoma induced by N-nitroso-N-ethylurea in mice. <i>Cancer Science</i> , 2013, 104, 1560-1566.	3.9	10
62	2-Amino-3-Methylimidazo[4,5-f]Quinoline (IQ) Promotes Mouse Hepatocarcinogenesis by Activating Transforming Growth Factor- α and Wnt/ β -Catenin Signaling Pathways. <i>Toxicological Sciences</i> , 2012, 125, 392-400.	3.1	4
63	Hormonally Active Doses of Isoflavone Aglycones Promote Mammary and Endometrial Carcinogenesis and Alter the Molecular Tumor Environment in Donryu Rats. <i>Toxicological Sciences</i> , 2012, 126, 39-51.	3.1	23
64	Long-term treatment with l-Isoleucine or l-Leucine in AIN-93G diet has promoting effects on rat bladder carcinogenesis. <i>Food and Chemical Toxicology</i> , 2012, 50, 3934-3940.	3.6	14
65	Dammar resin, a non-mutagen, induces oxidative stress and metabolic enzymes in the liver of gpt delta transgenic mouse which is different from a mutagen, 2-amino-3-methylimidazo[4,5-f]quinoline. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2012, 748, 29-35.	1.7	7
66	DDX39 acts as a suppressor of invasion for bladder cancer. <i>Cancer Science</i> , 2012, 103, 1363-1369.	3.9	27
67	Low-dose carcinogenicity of 2-amino-3-methylimidazo[4,5-f]quinoline in rats: Evidence for the existence of no-effect levels and a mechanism involving p21 ^{Cip1} /p27 ^{Swaf1} . <i>Cancer Science</i> , 2011, 102, 88-94.	3.9	19
68	Mitochondrial Prohibitins and Septin 9 Are Implicated in the Onset of Rat Hepatocarcinogenesis. <i>Toxicological Sciences</i> , 2011, 119, 61-72.	3.1	44
69	Targeted Proteomics of Isolated Glomeruli from the Kidneys of Diabetic Rats: Sorbin and SH3 Domain Containing 2 Is a Novel Protein Associated with Diabetic Nephropathy. <i>Experimental Diabetes Research</i> , 2011, 2011, 1-11.	3.8	28
70	Enhanced Urinary Bladder, Liver and Colon Carcinogenesis in Zucker Diabetic Fatty Rats in a Multiorgan Carcinogenesis Bioassay: Evidence for Mechanisms Involving Activation of PI3K Signaling and Impairment of P53 on Urinary Bladder Carcinogenesis. <i>Journal of Toxicologic Pathology</i> , 2011, 24, 25-36.	0.7	12
71	Sensitive quantitative assay for point mutations in the rat H-ras gene based on single nucleotide primer extension. <i>Experimental and Therapeutic Medicine</i> , 2010, 1, 657-661.	1.8	5
72	Cytokeratin 8/18 as a new marker of mouse liver preneoplastic lesions. <i>Toxicology and Applied Pharmacology</i> , 2010, 242, 47-55.	2.8	29

#	ARTICLE	IF	CITATIONS
73	Chemopreventive effects of 13 β ,14 β -epoxy-3 β -methoxyserratane-21 β -ol (PJ-34), a serratane-type triterpenoid, in a rat multi-organ carcinogenesis bioassay. <i>Cancer Letters</i> , 2010, 289, 161-169.	7.2	8
74	Potassium Bromate Enhances N-Ethyl-N-Hydroxyethylnitrosamine-Induced Kidney Carcinogenesis Only at High Doses in Wistar Rats: Indication of the Existence of an Enhancement Threshold. <i>Toxicologic Pathology</i> , 2009, 37, 983-991.	1.8	14
75	Enhancement of preneoplastic lesion yield by Chios Mastic Gum in a rat liver medium-term carcinogenesis bioassay. <i>Toxicology and Applied Pharmacology</i> , 2009, 234, 135-142.	2.8	22
76	Cytokeratin 8/18 overexpression and complex formation as an indicator of GST-P positive foci transformation into hepatocellular carcinomas. <i>Toxicology and Applied Pharmacology</i> , 2009, 238, 71-79.	2.8	32
77	Evaluation of initiation activity of dimethylarsinic acid: Initiation potential of rat hepatocarcinogenesis. <i>Toxicological and Environmental Chemistry</i> , 2009, 91, 1339-1351.	1.2	0
78	Existence of a Threshold for the Genotoxic Carcinogens: Evidence from Mechanism-based Carcinogenicity Studies. <i>Genes and Environment</i> , 2009, 31, 33-36.	2.1	5
79	Elevated oxidative stress and DNA damage and repair levels in urinary bladder carcinomas associated with schistosomiasis. <i>International Journal of Cancer</i> , 2008, 123, 601-608.	5.1	47
80	Lack of mutagenic and toxic effects of low dose potassium bromate on kidneys in the Big Blue rat. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2008, 652, 1-11.	1.7	32
81	Chemopreventive effects of a serratane-type triterpenoid, 3 β -methoxyserrat-14-en-21 β -ol (PJ-1), against rat lung carcinogenesis. <i>Food and Chemical Toxicology</i> , 2008, 46, 1882-1888.	3.6	12
82	Possible Involvement of Adaptation Mechanisms in the Achievement of an Ineffective Dose Range for the Carcinogenicity of Genotoxic Carcinogens. <i>Genes and Environment</i> , 2008, 30, 125-131.	2.1	4
83	Evaluation of the toxicity of mastic gum with 13 weeks dietary administration to F344 rats. <i>Food and Chemical Toxicology</i> , 2007, 45, 494-501.	3.6	24
84	Oral administration of diphenylarsinic acid, a degradation product of chemical warfare agents, induces oxidative and nitrosative stress in cerebellar Purkinje cells. <i>Life Sciences</i> , 2007, 81, 1518-1525.	4.3	37
85	Altered Gene Expression in Rat Colonic Adenocarcinomas Induced in an Azoxymethane plus 2-Amino-1-Methyl-6-Phenylimidazo[4,5-b]pyridine Initiation-Promotion Model. <i>Oncology</i> , 2007, 73, 252-260.	1.9	6
86	Carcinogenicity of dimethylarsinic acid in Ogg1-deficient mice. <i>Cancer Science</i> , 2007, 98, 803-814.	3.9	41
87	Elevation of 8-hydroxydeoxyguanosine and cell proliferation via generation of oxidative stress by organic arsenicals contributes to their carcinogenicity in the rat liver and bladder. <i>Toxicology and Applied Pharmacology</i> , 2007, 221, 295-305.	2.8	53
88	Alpha-benzene hexachloride exerts hormesis in preneoplastic lesion formation of rat hepatocarcinogenesis with the possible role for hepatic detoxifying enzymes. <i>Cancer Letters</i> , 2006, 240, 102-113.	7.2	32
89	A comparative study of the sub-chronic toxic effects of three organic arsenical compounds on the urothelium in F344 rats; gender-based differences in response. <i>Toxicology and Applied Pharmacology</i> , 2006, 210, 171-180.	2.8	35
90	Ingestion of Hijiki seaweed and risk of arsenic poisoning. <i>Applied Organometallic Chemistry</i> , 2006, 20, 557-564.	3.5	22

#	ARTICLE	IF	CITATIONS
91	Existence of No Hepatocarcinogenic Effect Levels of 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline with or without Coadministration with Ethanol. <i>Toxicologic Pathology</i> , 2006, 34, 232-236.	1.8	8
92	Inhibition of rat urinary bladder carcinogenesis by the antiangiogenic drug TNP-470. <i>Asian Pacific Journal of Cancer Prevention</i> , 2006, 7, 101-7.	1.2	4
93	Effects of cessation of alcohol exposure on rat hepatocarcinogenesis. <i>Asian Pacific Journal of Cancer Prevention</i> , 2006, 7, 122-6.	1.2	1
94	Lack of promoting effects of phenobarbital at low dose on diethylnitrosamine-induced hepatocarcinogenesis in TGF- α transgenic mice. <i>Asian Pacific Journal of Cancer Prevention</i> , 2006, 7, 274-8.	1.2	4
95	Current and emerging challenges in toxicopathology: Carcinogenic threshold of phenobarbital and proof of arsenic carcinogenicity using rat medium-term bioassays for carcinogens. <i>Toxicology and Applied Pharmacology</i> , 2005, 207, 225-229.	2.8	13
96	Low dose DDT inhibition of hepatocarcinogenesis initiated by diethylnitrosamine in male rats: Possible mechanisms. <i>Toxicology and Applied Pharmacology</i> , 2005, 208, 285-294.	2.8	25
97	Lack of large intestinal carcinogenicity of 2-amino-6-methylphenylimidazo[4,5-b]pyridine at low doses in rats initiated with azoxymethane. <i>International Journal of Cancer</i> , 2005, 115, 870-878.	5.1	20
98	Hormesis and dose-response-mediated mechanisms in carcinogenesis: evidence for a threshold in carcinogenicity of non-genotoxic carcinogens. <i>Carcinogenesis</i> , 2005, 26, 1835-1845.	2.8	90
99	Lack of potential of low dose N-nitrosodimethylamine to induce preneoplastic lesions, glutathione S-transferase placental form-positive foci, in rat liver. <i>Cancer Letters</i> , 2005, 222, 11-15.	7.2	21
100	No-Observed Effect Levels for Carcinogenicity and for in vivo Mutagenicity of a Genotoxic Carcinogen. <i>Toxicological Sciences</i> , 2004, 81, 273-279.	3.1	49
101	Understanding arsenic carcinogenicity by the use of animal models. <i>Toxicology and Applied Pharmacology</i> , 2004, 198, 366-376.	2.8	77
102	Revised rat multi-organ carcinogenesis bioassay for whole-body detection of chemopreventive agents: modifying potential of S-methylcysteine. <i>Cancer Letters</i> , 2004, 206, 15-26.	7.2	7
103	Induction of glutathione S-transferase placental form positive foci in liver and epithelial hyperplasia in urinary bladder, but no tumor development in male Fischer 344 rats treated with monomethylarsonic acid for 104 weeks. <i>Toxicology and Applied Pharmacology</i> , 2003, 193, 335-345.	2.8	24
104	Lack of initiation activity in rat liver of low doses of 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline. <i>Cancer Letters</i> , 2003, 191, 35-40.	7.2	33
105	Enhancing risk of ethanol on MeIQx-induced rat hepatocarcinogenesis is accompanied with increased levels of cellular proliferation and oxidative stress. <i>Cancer Letters</i> , 2003, 192, 37-47.	7.2	10
106	Liver tumorigenicity of trimethylarsine oxide in male Fischer 344 rats-association with oxidative DNA damage and enhanced cell proliferation. <i>Carcinogenesis</i> , 2003, 24, 1827-1835.	2.8	55
107	Value of GST-P Positive Preneoplastic Hepatic Foci in Dose-Response Studies of Hepatocarcinogenesis: Evidence for Practical Thresholds with Both Genotoxic and Nongenotoxic Carcinogens. A Review of Recent Work. <i>Toxicologic Pathology</i> , 2003, 31, 80-86.	1.8	69
108	Carcinogenicity of dimethylarsinic acid in p53 heterozygous knockout and wild-type C57BL/6J mice. <i>Carcinogenesis</i> , 2003, 24, 335-342.	2.8	60

#	ARTICLE	IF	CITATIONS
109	Phenobarbital at low dose exerts hormesis in rat hepatocarcinogenesis by reducing oxidative DNA damage, altering cell proliferation, apoptosis and gene expression. <i>Carcinogenesis</i> , 2003, 24, 1389-1399.	2.8	75
110	Value of GST-P Positive Preneoplastic Hepatic Foci in Dose-Response Studies of Hepatocarcinogenesis: Evidence for Practical Thresholds with Both Genotoxic and Nongenotoxic Carcinogens. A Review of Recent Work. <i>Toxicologic Pathology</i> , 2003, 31, 80-86.	1.8	68
111	Carcinogenicity of dimethylarsinic acid in male F344 rats and genetic alterations in induced urinary bladder tumors. <i>Carcinogenesis</i> , 2002, 23, 1387-1397.	2.8	207
112	Formation of 8-hydroxydeoxyguanosine and cell-cycle arrest in the rat liver via generation of oxidative stress by phenobarbital: association with expression profiles of p21WAF1/Cip1, cyclin D1 and Ogg1. <i>Carcinogenesis</i> , 2002, 23, 341-349.	2.8	94
113	Lack of promoting effect due to oral administration of dimethylarsinic acid on rat lung carcinogenesis initiated with N-bis(2-hydroxypropyl)nitrosamine. <i>Cancer Letters</i> , 2002, 175, 113-119.	7.2	15
114	Detailed low-dose study of 1,1-b?IS(p-chlorophenyl)-2,2,2- trichloroethane carcinogenesis suggests the possibility of a hormetic effect. <i>International Journal of Cancer</i> , 2002, 99, 112-118.	5.1	47
115	Promoting effects of monomethylarsonic acid, dimethylarsinic acid and trimethylarsine oxide on induction of rat liver preneoplastic glutathione S-transferase placental form positive foci: A possible reactive oxygen species mechanism. <i>International Journal of Cancer</i> , 2002, 100, 136-139.	5.1	80
116	Lack of a Dose-response Relationship for Carcinogenicity in the Rat Liver with Low Doses of 2-Amino-3,8-dimethylimidazo[4,5-f]quinoxaline or N-Nitrosodiethylamine. <i>Japanese Journal of Cancer Research</i> , 2002, 93, 1076-1082.	1.7	66
117	Presence of a no-observed effect level for enhancing effects of development of the Î±-isomer of benzene hexachloride (Î±-BHC) on diethylnitrosamine-initiated hepatic foci in rats. <i>Cancer Letters</i> , 2001, 163, 179-185.	7.2	25
118	Promotion of Skin Carcinogenesis by Dimethylarsinic Acid in Keratin (K6)/ODCTransgenic Mice. <i>Japanese Journal of Cancer Research</i> , 2000, 91, 579-581.	1.7	41
119	Urinary bladder carcinogenicity of dimethylarsinic acid in male F344 rats. <i>Carcinogenesis</i> , 1999, 20, 1873-1876.	2.8	184
120	Lack of inhibitory effects of the ju-myo protein on development of glutathione S-transferase placental form-positive foci in the male F344 rat liver.. <i>Journal of Toxicological Sciences</i> , 1999, 24, 27-31.	1.5	1
121	Promotion of NCI-Black-Reiter male rat bladder carcinogenesis by dimethylarsinic acid an organic arsenic compound. <i>Cancer Letters</i> , 1998, 134, 29-36.	7.2	49
122	Promotion of Rat Hepatocarcinogenesis by Dimethylarsinic Acid: Association with Elevated Ornithine Decarboxylase Activity and Formation of 8-Hydroxydeoxyguanosine in the Liver. <i>Japanese Journal of Cancer Research</i> , 1997, 88, 1149-1154.	1.7	71
123	Significance of cyclin D1 overexpression in transitional cell carcinomas of the urinary bladder and its correlation with histopathologic features. , 1997, 79, 780-789.		88
124	Promoting effects of dimethylarsinic acid on N-butyl-N-(4-hydroxybutyl)nitrosamine-induced urinary bladder carcinogenesis in rats. <i>Carcinogenesis</i> , 1996, 17, 2435-4239.	2.8	130