
## Baoli Zhu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2111716/publications.pdf Version: 2024-02-01



Βλου ΖΗΙ

#	Article	IF	CITATIONS
1	Co-infection with respiratory pathogens among COVID-2019 cases. Virus Research, 2020, 285, 198005.	2.2	419
2	Toilets dominate environmental detection of severe acute respiratory syndrome coronavirus 2 in a hospital. Science of the Total Environment, 2021, 753, 141710.	8.0	114
3	Association of viral load with serum biomakers among COVID-19 cases. Virology, 2020, 546, 122-126.	2.4	68
4	Effects of glyphosate exposure on sperm concentration in rodents: A systematic review and meta-analysis. Environmental Toxicology and Pharmacology, 2017, 55, 148-155.	4.0	47
5	A Functional Ser326Cys Polymorphism in hOGG1 Is Associated with Noise-Induced Hearing Loss in a Chinese Population. PLoS ONE, 2014, 9, e89662.	2.5	37
6	Accurately Differentiating Between Patients With COVID-19, Patients With Other Viral Infections, and Healthy Individuals: Multimodal Late Fusion Learning Approach. Journal of Medical Internet Research, 2021, 23, e25535.	4.3	30
7	A reverse-transcription recombinase-aided amplification assay for the rapid detection of N gene of severe acute respiratory syndrome coronavirus 2(SARS-CoV-2). Virology, 2020, 549, 1-4.	2.4	29
8	A Multimodality Machine Learning Approach to Differentiate Severe and Nonsevere COVID-19: Model Development and Validation. Journal of Medical Internet Research, 2021, 23, e23948.	4.3	27
9	Plasma microRNAs expression profile in female workers occupationally exposed to mercury. Journal of Thoracic Disease, 2016, 8, 833-841.	1.4	23
10	Characteristics and Trends of Pneumoconiosis in the Jiangsu Province, China, 2006–2017. International Journal of Environmental Research and Public Health, 2019, 16, 437.	2.6	20
11	Genetic variation in APE1 gene promoter is associated with noise-induced hearing loss in a Chinese population. International Archives of Occupational and Environmental Health, 2016, 89, 621-628.	2.3	14
12	PON2 and ATP2B2 gene polymorphisms with noise-induced hearing loss. Journal of Thoracic Disease, 2016, 8, 430-438.	1.4	13
13	Effects of occupational exposure to dust on chest radiograph, pulmonary function, blood pressure and electrocardiogram among coal miners in an eastern province, China. BMC Public Health, 2019, 19, 1229.	2.9	13
14	Genetic Variations in the Promoter of the APE1 Gene Are Associated with DMF-Induced Abnormal Liver Function: A Case-Control Study in a Chinese Population. International Journal of Environmental Research and Public Health, 2016, 13, 752.	2.6	10
15	Evaluating the Effectiveness of Earplugs in Preventing Noise-Induced Hearing Loss in an Auto Parts Factory in China. International Journal of Environmental Research and Public Health, 2021, 18, 7190.	2.6	9
16	Correlation between CAT polymorphism and susceptibility to DMAc-induced abnormal liver function: a case-control study of Chinese population. Biomarkers, 2018, 23, 147-153.	1.9	7
17	THE FREQUENCIES OF X-RAY EXAMINATIONS AND CT SCANS: FINDINGS FROM A SAMPLE INVESTIGATION IN JIANGSU, CHINA. Radiation Protection Dosimetry, 2020, 190, 38-44.	0.8	5
18	Verifying earplug attenuation and evaluating the effectiveness of oneâ€onâ€one training along with earplug fit testing at nine facilities in China. American Journal of Industrial Medicine, 2021, 64, 771-780.	2.1	5

Baoli Zhu

#	Article	IF	CITATIONS
19	Association between single nucleotide polymorphism (rs4252424) in TRPV5 calcium channel gene and lead poisoning in Chinese workers. Molecular Genetics & Genomic Medicine, 2019, 7, e562.	1.2	4
20	Polyurethane foam sampling for the determination of acetochlor in the air of workplace-gas chromatography. Chemical Papers, 2022, 76, 2375-2384.	2.2	4
21	Analysis of Polymorphisms Associated with Base Excision Repair in Patients Susceptible and Resistant to Noise-Induced Hearing Loss. Disease Markers, 2019, 2019, 1-8.	1.3	3
22	Salt-Inducible Kinase 3 Haplotypes Associated with Noise-Induced Hearing Loss in Chinese Workers. Audiology and Neuro-Otology, 2020, 25, 200-208.	1.3	3
23	A missense variant rs2585405 in clock gene PER1 is associated with the increased risk of noise-induced hearing loss in a Chinese occupational population. BMC Medical Genomics, 2021, 14, 221.	1.5	3
24	Lead and noise exposures at eight Chinese registered electronics recycling facilities. International Journal of Hygiene and Environmental Health, 2020, 230, 113611.	4.3	2
25	Variations in the potassium voltage-gated channel subfamily E regulatory subunit 1 gene associated with noise-induced hearing loss in the Chinese population. Environmental Science and Pollution Research, 2020, 27, 18822-18830.	5.3	2